

## **EFFECT OF SUCROSE AND GROWTH REGULATORS ON THE MICROPROPAGATION OF SUGARCANE CLONES**

**IMTIAZ AHMED KHAN, \*M. UMAR DAHOT, SHAFQAT YASMIN,  
ABDULLAH KHATRI, NIGHAT SEEMA AND MAZHAR H. NAQVI**

*Agriculture Biotechnology Division,  
Nuclear Institute of Agriculture, Tando Jam, 70060, Sindh, Pakistan*  
*\*Institute of Biotechnology and Genetic Engineering,  
Sindh University, Jamshoro, Pakistan*

### **Abstract**

The rapid multiplication of early, mid and late maturing sugarcane clones were obtained under different concentrations of commercial sugar and plant growth regulators in this study. Late maturing sugarcane clone AEC82-223 did not produce any auxillary shoot at 6% commercial sugar concentration. In contrast multiplication rate was higher in early and mid maturing sugarcane clones NIA-2004, BL4 and NIA-98, respectively. Apical meristems were cultured on modified MS medium containing different concentrations of auxins and cytokinins. An optimal multiplication was observed on M4 (1.0 mg/l IAA + 1.0 mg/l BAP + 0.1mg/l Kinetin) and M5 (0.1 mg/l kinetin + 1.5mg/l BAP + 1.5mg/l IAA). Maximum numbers of shoot were observed in BL4 followed by NIA-2004 when 1.0 mg/l of BAP and IAA were applied with low concentration of kinetin (0.1 mg/l). Shoot elongation and multiplication was improved on media devoid of BAP (MS+ 2 mg/l kinetin + 2 mg/l IBA + 2 mg/l IAA). Best rooting was observed on media containing MS+ 1 mg/l IBA + 6% sucrose.

### **Introduction**

Sugarcane is the second largest cash crop of Pakistan (Naz, 2003) with the country's sugar industries solely dependent on the fate of this crop. Although Pakistan is the fifth largest sugarcane growing country in terms of production area, its per hectare yield is the lowest among all sugarcane producing countries (Anon., 2003; Khan *et al.*, 2004). Uneconomical yield and lower sugar recovery cause very high production cost which makes Pakistan the least competitive in domestic and international sugar markets (Khan *et al.*, 2005).

There are many causes of low yield, one of which is the lack of a rapid seed multiplication procedure. Once a desired clone is identified, it usually takes 6-7 years to produce sufficient quality of improved seed material. This long duration causes a major bottleneck in breeding programmes (Siddiqui *et al.*, 1994). Another important reason for low yield in sugarcane is its susceptibility to attacks by pathogens such as fungi, virus, bacteria and mycoplasma which cause up to 70% in yields reduction (Xue & Chen., 1994; Oropez *et al.*, 1995; Bhavan & Gautam, 2002). For instance, sugarcane mosaic virus (SCMV) is found in almost all the cultivars grown in the sub-continent (Naz, 2003). A significant part of the yield (39-40%) is lost each year due to SCMV (Malik & Munir, 1990). As sugarcane is mostly propagated by vegetative means, once a plant becomes infected by a pathogen it can easily transfer the pathogen from one generation to another. For this reason, sugarcane seed (seed cane) production through micropropagation is a suitable and effective method for rapid propagation in comparison to conventional methods.

All the correspondence may kindly be made to Imtiaz Ahmed Khan e-mail: imtiaz\_khan2@hotmail.com

Initial attempts to regenerate plants through *In vitro* techniques were conducted on sugarcane by Nickell (1964) and Heinz & Mee (1969). Protocols for *In vitro* plant regeneration of sugarcane through callus culture, axillary bud and shoot tip culture have been developed by many authors (Lee 1986, 1987; Hu & Wang 1983; Hendre *et al.*, 1983; Nagai, 1987; Milton & Alien 1995; Baksha *et al.*, 2002). One of the major obstacles to the *In vitro* micropropagation of plants is the genotype / media interaction and rooting of the plantlet. Sugarcane is a highly heterozygous, polyploid and aneuploid crop (Jannoo *et al.*, 1999) and as a consequence the frequency of shoot differentiation from apical shoots in most sugarcane varieties varies greatly in number (Siddiqui *et al.*, 1994). Mulleegadoo & Dookun (1999) examined the effect of explant and genotype on growth of sugarcane under *In vitro* conditions. Although considerable advancement in tissue culture systems for micropropagation has been achieved, there is little information regarding how stalk sucrose concentration influences *In vitro* propagation of the plant. All commercial clones are divided into 3 groups on the basis of stalk sucrose concentration (Javed *et al.*, 2000). The present investigation has been undertaken to determine the propagation rate of different genotypes under different auxin and sugar concentrations. Commercial sugar was used rather than AnalaR grade sucrose, making this technique economically viable and technically feasible.

### Materials and Methods

Four sugarcane clones (*Saccharum* spp. hybrid) were selected on the basis of their maturity viz., NIA-2004, BL4 (early maturing), NIA-98 (mid maturing) and AEC82-223 (late maturing) were used in this study. Maturity parameter was set on the basis of their sucrose concentration in cane stalk (Khan *et al.*, 2005). Ten explants containing apical meristems were taken from each genotype, sterilized by a standard procedure (Siddiqui *et al.*, 1994) and cultured on modified MS medium (Murashige & Skoog, 1962) supplemented with different concentrations of growth regulators (Table 1&2). Two different concentrations of commercial sugar (4% and 6%) were used in order to study the effect of sucrose on initial multiplication. The explants were kept in the dark for 15, 20, 25, and 30 days to check the effect of darkness on phenol production in the explant. Data on shoot initiation were recorded and shootlets were then subcultured on shoot elongation and multiplication media S1) MS+ 2 mg/l kinetin + 2 mg/l IBA + 2 mg/l IAA and (S2) MS+ 4.5 mg/l BAP having two different sugar concentration i.e., 4 and 6%. Plantlets of different height (3, 4, 5, 6, 7 cm) were subjected to rooting by transferring them on three different media i) MS + 1 mg/l IBA + 6% sugar, ii) MS + 1 mg/l NAA + 6% sugar and iii) MS + 1 mg/l IBA+1mg/l NAA + 6% sugar. All these operations were carried out under aseptic conditions and cultures were incubated at  $28 \pm 2^{\circ}\text{C}$  with a 16 hours photoperiod. Media was solidified with 0.2% gelrite. Commercial sugar was used instead of AnalaR grade sucrose as a carbon source in the medium. Rooted plantlets were acclimatized and transplanted to the field. The mean and standard deviations were computed from each treatment. Data was analysed by using Duncan Multiple range test.

### Results and Discussions

Significant effect of different combination of Kinetin, BAP and IAA were observed. Apical meristems elongated a few days after incubation (Fig 1a). It was observed that shoot initiation in most of the clones started 15 days after explantation. Excessive phenol

**Table 1. Effect of different concentration of Kinetin, BAP and IAA 4% sucrose concentration on shoot regeneration of different sugarcane varieties after one month of incubation.**

Medium (mg/l) with 4% sucrose	Average number of shoots observed after one month of incubation			
	NIA-98	NIA-2004	BL4	AEC82-223
M1 = 0.1 + 0.1 + 0.0	4.25 ± 0.50b	7.50 ± 0.57b	4.75 ± 0.50b	6.75 ± 0.50b
M2 = 0.1 + 0.5 + 0.0	3.50 ± 0.57c	3.00 ± 0.81c	4.00 ± 0.81b	5.25 ± 0.50c
M3 = 0.1 + 1.0 + 0.0	4.25 ± 0.52b	3.50 ± 0.57c	4.00 ± 0.81b	4.00 ± 0.81d
M4 = 0.1 + 1.0 + 1.0	7.75 ± 0.51a	9.25 ± 0.50a	8.25 ± 0.95a	11.50 ± 0.57a
M5 = 0.1 + 1.5 + 1.0	4.75 ± 0.49b	7.50 ± 0.57b	7.25 ± 0.95a	3.75 ± 0.95d

DMR Test: Means denoted by similar letter showed non significant difference among the treatments

**Table 2. Effect of different concentration of Kinetin, BAP and IAA at 6% sucrose concentration on shoot regeneration of different sugarcane varieties after one month of incubation.**

Medium (mg/l) with 6% sucrose	Average number of shoots observed after one month of incubation			
	NIA-98	NIA-2004	BL4	AEC82-223
M1 = 0.1 + 0.5 + 0.0	6.75 ± 0.50c	8.50 ± 0.57c	8.25 ± 0.50d	-
M2 = 0.1 + 1.0 + 0.0	6.50 ± 0.57c	5.25 ± 0.50d	6.00 ± 0.81e	-
M3 = 0.1 + 1.5 + 0.0	9.25 ± 0.95a	10.00 ± 0.81b	9.50 ± 0.57c	-
M4 = 0.1 + 1.0 + 1.0	7.50 ± 0.57bc	11.75 ± 0.50a	10.50 ± 0.57b	-
M5 = 0.1 + 1.5 + 1.0	8.25 ± 0.95a	12.00 ± 0.81a	11.75 ± 0.95a	-

DMR Test: Means denoted by similar letter showed non significant difference among the treatments




Fig. 1. a- Explant of sugarcane; b- Excessives phenol production due to no dark treatment; c- Shoot initiation; d & e- Shoot multiplication; f- Sugarcane lanlfet in jiffy pot; g- Plantlets in the field for evaluation.

production was observed in cases when explants were not kept in darkness (Fig. 1b), with 20 days of darkness treatment showing good result. When 1.0 mg/l of BAP and IAA were applied along with 0.1 mg/l kinetin at a 4% sugar concentration, better shoot regeneration was observed in all clones when compared to other hormonal combinations (Table 1, Fig. 1c). Clone AEC82-223, which is a late maturing clonal line, showed a variable response according to sugar concentrations. It showed significantly higher shoot multiplication at 4% sugar concentration. In contrast, clones NIA-2004 and BL4, which are early maturing sugarcane lines, and NIA-98, mid maturing sugarcane line, all exhibited better multiplication at 6% sugar concentration. These results showed that in addition to hormonal concentration, sugar plays a vital role in sugarcane shoot multiplication under *In-vitro* conditions (Table 1 & 2). Chengalrayan & Meagher (2001) reported that auxins and cytokinin concentration influence shoot regeneration. Siddiqui *et al.*, (1994), reported the positive effect of lower concentrations of BAP and Kin on shoot proliferation of sugarcane. Sorory & Hosien (2000) also reported that the use of 6% sucrose concentration enhanced shoot regeneration in sugarcane.

Higher shoot multiplication was observed in media containing kinetin, IBA and IAA (Table 3, Fig. 1d & 1e). Clone NIA-98 showed a higher shoot multiplication on medium containing high concentration of BAP (4.5 mg/l) at both sugar concentrations (Table 3 & 4). Statistically non-significant differences were observed between NIA-2004, BL4 and AEC82-223. However, maximum shoot multiplication was observed in clone AEC82-223 and the lowest in NIA-98 at 4% sugar concentration. At 6% sugar concentration maximum shoot multiplication was observed in NIA-2004. The result suggests that shoot multiplication in sugarcane is dependent on the genotype/media interaction and sugar concentration also play significant role in shoot multiplication. Serving as a carbohydrate source, it also regulates the osmolarity of the culture media and plays a role during morphogenesis (Sopory, 1979). Nagai (1988) reported that high concentration of BAP suppressed the shoot proliferation in sugarcane. Geetha & Padmanadhan (2001) reported that the combination of BAP with Kin gave the maximum response in most varieties. In our study we found that addition of IAA along with IBA and Kin enhances the production of *In vitro* sugarcane plantlets. Rajesh *et al.*, (1994) studied *In vitro* clonal propagation of sugarcane with modified MS media supplemented with IAA, BAP and Kin with 0.5 mg/l of each for optimal growth. Pawar *et al.*, (2002) also obtained similar results with low concentration of IAA + BAP + Kin for mass multiplication of sugarcane.

Roots grow from the nodal primordial when the plantlets are well developed (Khan *et al.*, 1998). Rooting was highly influenced by the different types and concentrations of auxin used (Table 5). Appropriate amounts of auxin in the rooting medium are crucial for root induction. Among three auxins concentrations, IBA at 1 mg/l produced the highest percentage of rooting. These results confirm the previous findings of Khatri *et al.*, (2002), whereas, Lal & Singh (1994) who reported that the most efficient auxin for root initiation was NAA. Shenk & Hildebrandt (1972) have also reported a requirement of high concentration of auxin for rooting in sugarcane. A more vigorous root development was achieved when the plantlets were separated, the leaves trimmed and plantlets cultured on a root induction MS medium containing 1 mg/l IBA and 6% sucrose (Fig. 5). It was also observed that plantlets of less than 5 cm height did not produce good roots. Plantlets with well developed shoots and roots were transferred to jiffy pots containing sterilized perlite (Fig. 6). After acclimatization, plantlets were initially transferred to earthen pots for hardening and subsequently to the field (Fig 7). These plantlets are currently being evaluated for desired agronomic traits.

**Table 3. Effect of phytohormone on shoot multiplication of different sugarcane varieties after one month of incubation.**

Medium (mg/1) with 4% sucrose	Average number of shoots multiplied after one month			
	NIA-98	NIA-2004	BL4	AEC82-223
MS1-2.0+ 0.0+2.0+2.0	7.50 ±0.57b	10.50 ±0.57a	10.0 ±0.81a	11.00 ±0.81a
MS2-0.0 + 4.5 + 0.0 + 0.0	8.25 ±0.95b	9.50 ±0.57a	8.25 ±0.95b	8.50 ±0.57b

DMR Test: Means denoted by similar letter showed non significant difference among the treatments

**Table 4. Effect of phytohormone on shoot multiplication of different sugarcane varieties after one month of incubation.**

Medium (mg/1) with 6% sucrose	Average number of shoots multiplied after one month			
	NIA-98	NIA-2004	BL4	AEC82-223
MS1-2.0+ 0.0+2.0+2.0	6.75 ±0.95b	10.75 ±0.96a	10.50±0.58a	-
MS2-0.0 + 4.5 + 0.0 + 0.0	8.25 ±0.95a	9.50 ±0.58b	8.75 ±0.50b	-

DMR Test: Means denoted by similar letter showed non significant difference among the treatments

**Table 5. Effect of phytohormone on root induction of different sugarcane varieties after one month of incubation.**

Medium (mg/1) with 6% sucrose	Average number of shoots multiplied after one month				
	NIA-98	NIA-2004	BL4	AEC82-223	Thatta-10
MS +IBA +NAA					
MS + 1.0 + 0.0	+++	+++	+++	+++	+++
MS + 0.0 + 1.0	+	+	+	+	+
MS + 1.0 + 1.0	-	-	-	-	-

- , No rooting, +, week rooting, +++, Profuse rooting

The results of the experiments with different concentrations of sucrose showed that the rates of sugarcane micro shoots obtained from micropropagule were greatly influenced by the concentration of sugar in the medium. Of the two concentrations tested, 4% commercial sugar appeared to be optimum for shoot regeneration and the same can be used for multiplication, whereas 6% commercial sugar was recommended for rooting. The present study would suggest an efficient and easy to handle protocol for micropropagation of sugarcane. This protocol provides a successful technique that can be used for rapid propagation. The application of this protocol can help minimize the variety assessment period and contribute to the rapid propagation of high yielding sugarcane cultivars.

#### Acknowledgments

The authors are thankful to Dr. Mazhar H. Naqvi, Director, Nuclear Institute of Agriculture, Tando Jam, Sindh, Pakistan for the support during the research work. Thanks also due to Dr. J. L. Noyer, CIRAD, France, for critical review of the paper.

#### References

- Anonymous. 2003. *Agricultural Statistics of Pakistan*, 2002-03. Govt. Pakistan, 2003; 27-28; 106.
- Baksha, R., R. Alam, M.Z. Karim, S.K. Paul, M.A. Hossain, M.A.S. Miah and A.B.M.M. Rahman. 2002. *In vitro* Shoot Tip Culture of Sugarcane (*Saccharum officinarum*) Variety Isd 28. *Int.Quarterly J. Biotechnol.*, 1: 67-72.

- Bhavan, K. and G. Gautam. 2002. Micropropagation technology through tissue culture. *Indian Council for Agric. Res.*, 24-25.
- Chengalrayan, K. and G. Meagher. 2001. Effect of various growth regulators on shoot regeneration. *In vitro Cellular and Dev. Biol.*, 37: 434-439.
- Geetha, S. and D. Padmanadhan. 2001. Effect of hormones on direct somatic embryogenesis in sugarcane. *Sugar Tech.*, 3: 120-121.
- Heinz, D.J. and G.W. Mee. 1969. Plant differentiation from callus issue of *Saccharum* species. *Crop Sci.*, 9: 346-348.
- Hendre, R.R., R.S. Lyer, M. Kotwal, S.S. Khuspe and A.F. Mascavenhas. 1983. Rapid multiplication of sugarcane by tissue culture. *Sugarcane*, 1: 5-8.
- Hu, C.Y. and P.J. Wang. 1983. Meristem, shoot tip; bud culture. In: *Plant Cell Culture*, 1: 177-227. (Ed.): Evans *et al.*, New York Macmillan.
- Jannoo, N., L. Grivet, M. Segiun, F. Paulet, R. Domaingue, P.S. Rao, A. Dookun, A., D'Hont and J.C. Glaszmann. 1999. Molecular investigation of the genetic base of sugarcane cultivars. *Theor. Appl. Genet.*, 99: 171-184.
- Javed, M.A., A. Khatri, I.A. Khan, M. Ashraf, M.A. Siddiqui, N.A. Dahar, M.H. Khanzada and R. Khan. 2000. Prospects of enhancement / improvement of sugarcane productivity in Pakistan. *Pak. Sugar J.*, 15: 14-19.
- Khan I.A., A. Khatri, M. Ahmed, S.H. Siddiqui, G.S. Nizamani, M.H. Khanzada, N.A. Dahar and R. Khan. 1998. *In vitro* mutagenesis in sugarcane. *Pak. J. Bot.*, 30: 253-261.
- Khan, I.A., A. Khatri, G.S. Nizamani, M.A. Siddiqui, M.H. Khanzada, N.A. Dahar, N. Seema and M.H. Naqvi. 2004. *In vitro* studies in sugarcane. *Pak. J. Biotech.*, 1: 6-10.
- Khan, I.A., A. Khatri, M.A. Siddiqui, G.S. Nizamani, S. Raza, M.H. Khanzada, N.A. Dahar and R. Khan. 2005. Effect of NPK fertilizers on the growth of sugarcane clone AEC86-347 developed at NIA, Tando Jam. *Pak. J. Bot.*, 37: 355-360.
- Khatri, A., I.A. Khan, M.A. Javed, M.A. Siddiqui, M.K.R. Khan, M.H. Khanzada, N.A. Dahar and R. Khan. 2002. Studies on callusing and regeneration potential of indigenous and exotic sugarcane clones. *Asian J. Plant Sci.*, 1: 41-43.
- Lal, N. and N.H. Sing. 1994. Rapid clonal multiplication of sugarcane through tissue culture: *Plant Tissue Cult.*, 4: 1-7.
- Lee, T.S.G. 1986. Multiplication of sugarcane by apex culture. *Tumalba*, 36: 231-235.
- Lee, T.S.G. 1987. Micropropagation of sugarcane (*Saccharum* spp.). *Plant Cell, Tissue and Organ Culture*, 10: 47-55.
- Malik, K.B. and A. Munir. 1990. Sugarcane disease in Pakistan. *Pak. Sugar J.*, 4: 3-10.
- Milton, J. and D. Alien. 1995. *Breeding of field crops*. 4<sup>th</sup> Edition, 134-135.
- Mulleegadoo, K.D. and A. Dookun. 1999. Effect of explant source and genotype on growth of sugarcane *in vitro*. *Revue Agricole et Sucriere de l'Île Maurice*, 78: 35-39.
- Murashige, T. and F. Skoog. 1962. A revised medium for rapid growth and bioassays with tobacco tissue culture. *Physiol. Planta.*, 15: 473-497.
- Nagai, C. 1986. Micropropagation of H75-8776. *Annual Report 1986*. Experimental Station, Hawaiian Sugar Planters' Association. 4-5.
- Nagai, C. 1988. Micropropagation of sugarcane. Laboratory methodology: *Annual Report 1988*. Experimental Station, Hawaiian Sugar Planters' Association pp. A34-A37.
- Naz, S. 2003. Micropropagation of promising varieties of sugarcane and their acclimatization response. Activities on Sugar Crops in Pakistan. In: *Proc. Fourth Workshop Res. & Dev.*, 1-9.
- Nickell, L.G. 1964. Tissue culture of sugarcane, another research tool. *Hawaiian Planters Res.*, 57: 223-229.
- Oropez, M.P., R. Guevara and J.I. Ramiez. 1995. Identification of somaclonal variants of sugarcane resistant to sugarcane mosaic virus via RAPD markers. *Plant Mol. Biol. Rep.*, 13: 182-191.
- Pawar, S.V., S.C. Patil, V.M. Jambhale, R.M. Naik and S.S. Mehete. 2002. Rapid multiplication of commercial sugarcane varieties through tissue culture. *Indian Sugar*, 11: 183-186.
- Rajesh, S., A.Q. Khan and G.K. Gang. 1994. *In vitro* clonal propagation of sugarcane: Optimization of media and hardening of plants. *Indian Sugar*, 44: 113-116.

- Shenk, R.U. and A.C. Hildebrandt. 1972. Medium and techniques for induction and growth of monocotyledonous and dicotyledonous plant cell cultures. *Can. J. Bot.*, 50: 199-204.
- Siddiqui, S.H., I.A. Khan, A. Khatri and G.S. Nizamani. 1994. Rapid multiplication of sugarcane through micropropagation. *Pak. J. Agri. Res.*, 15: 134-136.
- Sopory S.K. 1979. Effect of sucrose, hormones and metabolic inhibitors on the development of pollen embryoids in anther cultures of dihaploid *Solanum tuberosum*. *Can. J. Bot.*, 57: 2691-2694.
- Sorory, A. and Z.A. Hosien. 2000. Effect of genotype, media and explant on callus initiation and plant regeneration in sugarcane. *Iranian J. Agri. Sci.*, 31: 211-220.
- Xue, L.P. and R.K. Chen. 1994. Elimination of sugarcane mosaic virus by callus tissue culture and apical culture. *J. Fujian Agric. Univ.*, 23: 253-256.

(Received for publication 30 December 2004)