

SEED DRESSING WITH BIOCONTROL AGENTS AND NEMATICIDES FOR THE CONTROL OF ROOT KNOT NEMATODE ON SUNFLOWER AND OKRA

SHAHNAZ DAWAR, A. SATTAR AND M.J. ZAKI

Department of Botany, University of Karachi,
Karachi-75270, Pakistan

Abstract

Application of biocontrol agents viz., *Rhizobium meliloti*, *Bacillus thuringiensis*, *Aspergillus niger*, *Trichoderma harzianum* and nematicides viz., fertinemakil, vermox as seed dressing for the reduction of *Meloidogyne javanica* infection on sunflower and okra was examined. *In vitro* experiments, maximum reduction in hatching of *M. javanica* egg was observed in *R. meliloti* aqueous extract whereas *T. harzianum* showed significant mortality of second stage juveniles of *M. javanica*. *In vivo* experiment the biocontrol agents and nematicides coated with sugar, mollasses, glucose and gum arabic used @ 1 and 2% significantly reduced the infection of root knot nematode on okra and sunflower roots. An increase in concentration of coating materials significantly enhanced the germination and growth parameters in terms of shoot length, shoot weight, root length and root weight on both okra and sunflower plants. Of the different microbial antagonists and nematicides used, *T. harzianum* and fertinemakil were found more effective followed by *R. meliloti*, *B. thuringiensis*, vermox and *A. niger* in the control of root knot nematode.

Introduction

The diseases by plant parasitic nematodes reduce the yield of the world's 40 major food and cash crop by an average of 12.3% for the 20 life sustaining crops that serves as men primary food source with an estimated annual yield loss of 10.7% for the 20 crops not considering to be life sustaining (Sasser & Freekman, 1987). Root knot nematodes are plant parasitic nematodes from the genus *Meloidogyne*. The members of the genus have a wide host range of plant species (Goodey *et al.*, 1965). Upto 70 *Meloidogyne* species have been identified so far (Luc *et al.*, 1988). *M. javanica* (Treb) Chitwood is the most common in subtropical and tropical regions (Schneider, 1991). About 2000 plants are susceptible to infection by root knot nematodes and they cause approximately 5% of global crop losses. Root knot nematode larvae infect plant root causing the development of root knot galls that drains the plant's photosynthetic and nutrient. In addition these nematodes have the ability to interact synergistically with other plant pathogen and cause up to 5-34% yield losses in vegetables in tropical climates (Eisenback & Triantaphyllous, 1991).

The biology, ecology and potential of biological control agents for the control of nematodes have been extensively reviewed in recent years (Kerry, 1987; Stirling, 1991; Sayre & Walter, 1991; Sikora, 1992; Baoyin Tian *et al.*, 2007). Nematologists have identified natural enemies with a range of modes of action similar to those currently studied by plant pathologists for the control of soil-borne diseases. *Rhizobium* spp., have a beneficial effect on plants including biological control of soil borne pathogens, induce systematic resistance to plant pathogen and improvement of nutrient uptake of plant (Seuk Bae *et al.*, 2000). The damage caused by nematode was more in unbacterized plants than in bacterized ones. Nematode multiplication reduced in the presence of *Rhizobium* (Siddiqui & Mahmood, 1995). *B. thuringiensis* toxins have also been shown to be somewhat active

towards species of *Meloidogyne* (Devidas & Rehberger, 1992). There are reports where use of *T. harzianum* significantly suppressed root knot diseases in maize (Windhum *et al.*, 1989). *T. harzianum* has also been found as an egg parasite of *M. javanica* race-3 killing 53% of eggs *In vitro* (Santos *et al.*, 1992). Present research was undertaken to control root knot nematode on okra (*Abelmoschus esculentus* L.) and sunflower (*Helianthus annuus* L.) by using different biological antagonists and nematicides.

Materials and Methods

Microbial antagonists viz., *Bacillus thuringiensis* (Bt 10), *Rhizobium meliloti* (R5), *Aspergillus niger* (An 20) and *Trichoderma harzianum* (KUCC 65) obtained from Karachi University culture collection (KUCC) and nematicides fertinmakil and vermax obtained from local market were used.

***In vitro* studies:** For hatching test, eggs of *M. javanica* were obtained from the roots of egg plant (*Solanum melongena* L.) collected by the method of Hussey & Barker (1973). Eggs suspension was prepared in distilled water and 2 ml suspension containing 20-40 eggs were poured in each cavity block with microbial antagonists and nematicides and kept at room temperature (34-38°C). Cavity blocks without suspension served as control. Each treatment was replicated three times. The numbers of juveniles were counted at 24, 48 and 72 hrs intervals.

For mortality test, freshly hatched second stage juveniles of *M. javanica* were suspended in sterile distilled water and 2 ml of this suspension containing 15-20 larvae/ml was placed in each cavity block. Cavity blocks without extract of microbial antagonists and nematicides served as control. There were three replicates of each treatment. The number of juveniles that were killed at 24, 48 and 72 h intervals was recorded using a stereoscope.

***In vivo* experiments:** The roots infested with *M. javanica* root knot nematode were collected from the experimental plot of Department of Botany, University of Karachi. The roots were washed under running tap water and cut into small pieces then dipped in 100 ml of 1% Ca (OCl)₂ in a bottle and the mouth was tightly closed then vigorously shake by hands for 5 min and content was poured on to a 100 or 200 mesh sieve fitted over a 400 mesh sieve, and the roots were washed under running tap water for 1 min. The residues from 400 mesh sieve were transferred into 250 ml beaker. Number of eggs and larvae/ml of suspension were determined with the help of counting dish (Hussey & Barker, 1973).

Five seeds each of okra (*Abelmoschus esculentus* L.) and sunflower (*Helianthus annuus* L.) were surface sterilized with 1% Ca(OCl)₂ for three minutes, rinsed thoroughly in running water and dried aseptically. The seeds were treated with microbial antagonists viz, *B. thuringiensis*, *R. meliloti*, *A. niger*, *T. harzianum* separately by using 1% and 2% sugar, molasses, glucose and gum arabic solution as a sticker. Ten seeds after treatment with suspension of microbial antagonists were transferred in test tube containing 9ml sterilized distilled water. The test tube was shaken and dilution series was made. One ml suspension was poured on PDA and cells/seed of bacteria and number of conidia/seed of fungi was calculated by using the formula:

$$\frac{\text{Number of cells}}{\text{Seed and conidia}} \times \text{dilution factor}$$

Table 1. Population of bacteria and fungi on seeds of okra and sunflower after seed treatment.

Treatments	Cfu/Seed			
	Okra		Sunflower	
	1%	2%	1%	2%
<i>A. niger</i>				
Sugar	16x10 ⁵	13 x10 ⁵	30 x10 ⁵	24 x10 ⁵
Mollases	14 x10 ⁵	14 x10 ⁵	16 x10 ⁵	21 x10 ⁵
Glucose	19 x10 ⁵	12 x10 ⁵	31 x10 ⁵	34 x10 ⁵
Gum arabic	13 x10 ⁵	15 x10 ⁵	15 x10 ⁵	15 x10 ⁵
<i>T. harzianum</i>				
Sugar	11 x10 ⁵	10 x10 ⁵	15 x10 ⁵	13 x10 ⁵
Mollases	18 x10 ⁵	17 x10 ⁵	17 x10 ⁵	14 x10 ⁵
Glucose	12 x10 ⁵	8 x10 ⁵	22 x10 ⁵	20 x10 ⁵
Gum arabic	10 x10 ⁵	5 x10 ⁵	10 x10 ⁵	7 x10 ⁵
<i>B. thuringiensis</i>				
Sugar	10 x10 ⁵	8 x10 ⁵	9 x10 ⁵	16 x10 ⁵
Mollases	38 x10 ⁵	7 x10 ⁵	9 x10 ⁵	12 x10 ⁵
Glucose	18 x10 ⁵	10 x10 ⁵	10 x10 ⁵	12 x10 ⁵
Gum arabic	14 x10 ⁵	5 x10 ⁵	12 x10 ⁵	6 x10 ⁵
<i>R. meliloti</i>				
Sugar	53 x10 ⁵	76 x10 ⁵	60 x10 ⁵	30 x10 ⁵
Mollases	83 x10 ⁵	40 x10 ⁵	72 x10 ⁵	48 x10 ⁵
Glucose	35 x10 ⁵	41 x10 ⁵	35 x10 ⁵	29 x10 ⁵
Gum arabic	40 x10 ⁵	70 x10 ⁵	45 x10 ⁵	32 x10 ⁵

In seed dressing, seeds of okra and sunflower coated with 48 hrs old cultures of *B. thuringiensis*, *R. meliloti*, *A. niger*, *T. harzianum*, fertinmakil and vermax using 1 and 2% gum arabic, sugar, mollases and glucose solution as sticker were sown in 8cm, diam., plastic pots, each pot containing 300gm soil. Pots were kept randomized on screen house bench at the Department of Botany, University of Karachi, where soil was kept at 40% MHC (Keen & Raczkowski, 1922). After two weeks of plant growth, the plants were inoculated with 2000 freshly hatched second stage juveniles by introducing holes around each plant. Pots without microbial antagonists served as control. Treatments and control were replicated thrice. After 60 days of growth, plants were uprooted and number of root knots was determined.

Data were analyzed and subjected to analysis of variance (ANOVA) using procedure given by Sokal & Rohlf (1995).

Results and Discussion

Population of bacteria and fungi after seed treatment was counted by serial dilution technique (Table 1). *In vitro* studies, aqueous extract of all biocontrol agents and nematicides showed reduction in *M. javanica* egg hatching. *R. meliloti* was found more effective in reducing egg hatching followed by *B. thuringiensis*, *A. niger*, *T. harzianum*, vermax and fertinmakil. Hatching of eggs was reduced considerably with the increase in time period (Table 2).

Table 2. Effect of aqueous extract of microbial antagonists and nematicides on hatching and mortality % of *Meloidogyne javanica*.

Treatments	Time (hrs)					
	Hatching %			Mortality %		
	24	48	72	24	48	72
Control	5	21	35	2	5	7
<i>A. niger</i>	3	5	8	10	17	34
<i>B. thuringiensis</i>	3	5	7	7	13	20
Fertinemakil	8	15	21	8	11	22
<i>R. meliloti</i>	2	2	3	4	12	24
<i>T. harzianum</i>	0	3	6	19	26	52
Vermox	3	7	15	3	9	17
	LSD0.05 _{treatment} = 2.76, LSD0.05 _{time} = 1.81			LSD0.05 _{treatment} = 4.37, LSD0.05 _{time} = 2.86		

Results showed that aqueous extract of biocontrol agents and nematicides caused appreciable mortality of *M. javanica* juveniles due to the reasons that they contain some compounds which are toxic to *M. javanica* and produce lethal effect. *T. harzianum* was found to be more effective as compared to other biocontrol agents and nematicides for mortality of *M. javanica* (Table 2).

An increase in germination of okra seeds and significant increase in shoot length, shoot weight, root length and root weight were observed. Significant reduction of number of knots ($p < 0.001$) was observed in okra plant when seeds were treated with microbial antagonists viz., *B. thuringiensis*, *R. meliloti*, *Aspergillus niger*, *T. harzianum* using 1 and 2% of sugar, molasses, glucose and gum arabic as stickers. There was significant increase in shoot length, shoot weight, root length and root weight ($p < 0.001$) and significant reduction in number of knots ($p < 0.001$) on sunflower plant was observed. Results showed that use of nematicides viz, fertinemakil and vermox significantly increased the germination and growth parameters in terms of shoot length, shoot weight, root length and root weight and reduction in number of knots was observed on sunflower and okra plants. Among the different microbial antagonists used *B. thuringiensis* and *T. harzianum* showed significant results for growth of plant and reduction in number of knots on both sunflower and okra plants. Of the different coating materials used with different concentration, 2% concentration showed significant reduction in number of knots on both sunflower and okra plants (Table 3).

Present result showed that aqueous extract of microbial antagonists exhibited nematicidal activity which reduced egg hatching of *M. javanica* and increased mortality of larvae with the increase in exposure time. Presumably the production of antibiotics (Dennis & Webster, 1971) and extracellular enzymes (Elad *et al.*, 1982) are involved in antagonisms. Dawar *et al.*, (2008) observed that *Bacillus* species viz., *B. subtilis*, *B. thuringiensis* and *B. cereus* significantly reduced hatching of larvae of *M. javanica* root knot whereas mortality of larvae was significantly increased with an increase in time. Seed treatment is an attractive method for introducing biocontrol agents into a soil root environment since it protects the seed from seed-borne and soil-borne pathogens and enable the seed to germinate and become established as a healthy seedling (Chang & Kommedahl, 1968). *T. harzianum* has also been found as egg parasite of *M. javanica* race-3 killing 53% of eggs *In vitro* (Santos *et al.*, 1992). Besides parasitisms of the root knot nematode, it is also hypothesized that the production of nematicidal compounds by *Trichoderma* spp., directly affect the nematode or made root less attractive which might

Table 3. Effect of seed dressing with fungal and bacterial antagonists on growth parameters and number of knots of okra and sunflower.

Treatments	Okra					Sunflower						
	Germination %	Shoot length (cm)	Shoot weight (g)	Root length (cm)	Root weight (g)	Number of knots	Germination %	Shoot length (cm)	Shoot weight (g)	Root length (cm)	Root weight (g)	Number of knots
Fungal antagonists												
Control	20	11	0.6	6.7	0.1	45	20	8.6	1.1	4.6	0.2	52
<i>A. niger</i> 1 % sugar	33	25	8.6	8.6	0.2	38	33	15	1.4	5	0.3	39
<i>A. niger</i> 2 % sugar	87	15	5.0	5.0	0.3	32	67	16	1.6	5.6	0.3	29
<i>A. niger</i> 1 % molasses	67	14	2.3	5.6	0.2	23	73	15	1.4	8	0.4	27
<i>A. niger</i> 2 % molasses	67	16	3.5	6.0	0.2	16	67	18	1.2	6	0.4	21
<i>A. niger</i> 1 % glucose	67	15	3.2	7.6	0.2	31	53	14	1.3	7.3	0.2	31
<i>A. niger</i> 2 % glucose	60	25	3.3	14.0	0.3	15	60	15	1.2	5.3	0.3	23
<i>A. niger</i> 1 % gum arabic	73	11	2.2	8.3	0.2	13	73	12	1.3	7.3	0.2	21
<i>A. niger</i> 2 % gum arabic	33	17	3.3	7.0	0.2	7	47	12	1.4	8.6	0.4	20
LSD _{0.05} =	30.0	2.1	0.5	1.7	0.1	5.6	30.0	1.5	0.1	1.1	0.1	5.8
Bacterial antagonists												
Control	40	11	0.6	6.7	0.3	45	20	12	1.1	4.3	0.2	52
<i>T. harzianum</i> 1 % sugar	40	31	3.0	10	0.2	31	47	11	1.7	6.3	0.6	23
<i>T. harzianum</i> 2 % sugar	87	32	3.6	14	0.2	23	60	14	1.8	8.0	0.7	17
<i>T. harzianum</i> 1 % molasses	67	26	2.9	8.6	0.2	25	33	13	1.9	4.6	0.3	26
<i>T. harzianum</i> 2 % molasses	53	16	2.6	7.3	0.2	15	73	14	2.0	7.3	0.4	10
<i>T. harzianum</i> 1 % glucose	47	21	2.2	7.0	0.9	25	47	18	2.0	7.0	0.3	18
<i>T. harzianum</i> 2 % glucose	67	20	2.1	11	0.9	22	67	15	1.8	8.3	0.4	13
<i>T. harzianum</i> 1 % gum arabic	47	26	2.4	13	1.1	16	47	12	1.8	4.6	0.6	12
<i>T. harzianum</i> 2 % gum arabic	67	15	2.3	8.3	1.1	10	67	18	1.7	6.3	0.6	3
LSD _{0.05} =	32.0	1.4	0.2	1.5	0.1	6.0	36.0	1.6	0.1	1.1	0.0	4.8

Table 3. (Cont'd.).

Treatments	Okra						Sunflower					
	Germination %	Shoot length (cm)	Shoot weight (g)	Root length (cm)	Root weight (g)	Number of knots	Germination %	Shoot length (cm)	Shoot weight (g)	Root length (cm)	Root weight (g)	Number of knots
Bacterial antagonists												
Control	53	11	0.6	6.7	0.3	45	53	12	1.1	3.6	0.2	52
<i>B. thuringiensis</i> 1 % sugar	47	20	3	7.8	1.1	31	47	21	2.2	8	0.6	28
<i>B. thuringiensis</i> 2 % sugar	60	22	3.1	9	1.2	21	60	15	3.0	8.3	0.3	17
<i>B. thuringiensis</i> 1 % mollases	67	20	3	7.8	1.1	23	67	15	1.7	5.3	0.4	26
<i>B. thuringiensis</i> 2 % mollases	87	25	3	13	1.1	8	73	15	1.6	5.3	0.4	16
<i>B. thuringiensis</i> 1 % glucose	67	15	3.2	6.7	1.1	23	67	17	1.9	5	0.6	22
<i>B. thuringiensis</i> 2 % glucose	93	24	3.3	7	1.3	16	93	18	1.5	9	0.6	16
<i>B. thuringiensis</i> 1 % gum arabic	47	30	3.4	14	1.2	17	47	17	1.7	5	0.5	22
<i>B. thuringiensis</i> 2 % gum arabic	67	30	3.5	15	1.3	8	67	14	1.6	7	0.5	9
LSD _{0.05} =	45	1.1	0.3	1.7	0.1	5.7	42	1.4	0.1	1.4	0.3	5.5
Control	47	11	0.6	6.3	0.3	45	47	12	0.4	5.0	0.1	52
<i>R. meliloti</i> 1 % sugar	93	14	2.3	11	0.3	32	93	21	0.4	8.6	0.2	29
<i>R. meliloti</i> 2 % sugar	100	15	2.3	12	0.3	23	100	21	1.1	10.0	0.3	28
<i>R. meliloti</i> 1 % mollases	87	25	2.2	6.7	0.4	24	73	21	0.5	7.5	0.2	27
<i>R. meliloti</i> 2 % mollases	93	11	2.2	15	0.4	23	93	24	0.6	11	0.3	20
<i>R. meliloti</i> 1 % glucose	60	12	2.1	7.3	0.3	27	60	16	0.7	8.0	0.2	30
<i>R. meliloti</i> 2 % glucose	87	12	2.0	8.3	0.2	26	87	15	1.2	8.0	0.3	28
<i>R. meliloti</i> 1 % gum arabic	60	25	2.1	12	0.4	20	60	15	0.8	5.0	0.1	23
<i>R. meliloti</i> 2 % gum arabic	67	31	2.1	12	0.3	12	67	22	1.3	10.0	0.2	14
LSD _{0.05} =	45	1.1	0.3	1.7	0.1	5.7	49	2.1	0.1	1.1	0.1	6.0

Table 3. (Cont'd.).

Treatments	Okra						Sunflower					
	Germination %	Shoot length (cm)	Shoot weight (g)	Root length (cm)	Root weight (g)	Number of knots	Germination %	Shoot length (cm)	Shoot weight (g)	Root length (cm)	Root weight (g)	Number of knots
Nematicides												
Control	40	11	0.6	6.3	0.3	45	13	3.6	1.1	4.0	0.2	52
Fertinimakil 1 % sugar	47	18	1.7	10	0.3	25	27	8.6	2.2	5.0	1.2	16
Fertinimakil 2 % sugar	60	20	1.9	12	0.9	20	40	12	2.3	5.0	2.0	15
Fertinimakil 1 % mollasses	60	19	2.2	6.6	0.4	28	20	8.6	2.7	6.3	1.2	22
Fertinimakil 2 % mollasses	87	21	2.2	7.3	0.4	23	73	11	3.0	8.0	1.4	19
Fertinimakil 1 % glucose	60	20	2.1	7.0	0.3	28	33	9	2.9	6.3	1.3	14
Fertinimakil 2 % glucose	73	22	2.3	7.6	0.3	17	40	12	2.9	7.3	1.3	9
Fertinimakil 1 % gum arabic	60	12	2.5	6.7	0.3	22	33	11	2.1	6.0	1.1	5
Fertinimakil 2 % gum arabic	67	15	2.5	10	0.3	5	40	16	2.4	9.0	1.4	0
LSD0.05=	35	1.4	0.1	0.9	0.7	6	32	1.1	0.1	1.3	0.1	3.4
Control	53	11	0.6	6.7	0.2	45	27	9	1.1	4.0	0.2	52
Vermox 1 % sugar	60	12	2.8	8.6	0.3	29	40	9.6	2.3	4.3	0.7	22
Vermox 2 % sugar	60	18	3.0	12	0.3	19	73	12	2.4	5.0	0.7	14
Vermox 1 % mollasses	80	12	3.0	6.6	0.3	23	20	10	2.2	7.0	1.1	23
Vermox 2 % mollasses	87	16	3.3	7.3	0.4	17	47	13	2.3	8.6	1.2	16
Vermox 1 % glucose	53	19	3.0	8.6	0.4	20	33	10	2.9	5.0	1.0	19
Vermox 2 % glucose	60	21	3.3	10	1.1	13	53	19	5.6	8.0	1.2	12
Vermox 1 % gum arabic	53	17	3.1	8	1.1	18	33	14	2.4	5.6	1.2	12
Vermox 2 % gum arabic	73	21	3.2	9.6	1.2	9	33	15	2.6	7.0	1.4	7
LSD0.05=	36	1.4	0.1	1.2	0.0	6.5	37	1.5	0.1	1.2	0.1	5.3

have resulted in the reduction in nematode population. In the present study, maximum disease suppression induced by *T. harzianum* followed by *A. niger* is attributed to its parasitic nature and production of nematocidal serine proteases which degrades egg shell and check egg hatching (Bonants *et al.*, 1994). *A. niger* applied alone or in combination with the bacterial inoculants inhibited root-knot nematode galling in tomato (Siddiqui *et al.*, 2003).

In the present study, shoot length, shoot weight, root length and root weight were significantly increased in sunflower and okra when seeds were coated with *R. meliloti* and *B. thuringiensis*. Similar report was made by (Siddique *et al.*, 2000) in okra where *Rhizobia* used as seed dressing and soil drenching significantly increased growth parameters and number of nodules. In the present investigation, *Rhizobium* used either as seed dressing significantly improved plant growth and reduced disease intensity of plants due to initial colonizers of rhizospheres of test plants. It is interesting to note that *Rhizobia* not only showed significant control of root pathogens on leguminous plants like chickpea, mungbean as well as non leguminous plants like okra and sunflower but also increased plant height and fresh shoot weight (Zaki, 2000). Present results showed that seed coated with *B. thuringiensis* (Bt) strains exhibited nematocidal activity on okra and sunflower. *B. thuringiensis* toxins have also been shown to be somewhat active towards species of *Meloidogyne* (Devidas & Rehberger, 1992).

The results of the present study indicates the potentialities of seed treatment with fungal and bacterial antagonists viz., *B. thuringiensis*, *R. meliloti*, *A. niger*, *T. harzianum* and nematicides in the suppression of root knot nematode on okra and sunflower. There is therefore need to characterize nematocidal compound produced by biological antagonists resulting in control of root knot nematode instead of use of pesticides, which are costly and hazardous.

References

- Baoyn Tian, Jinkui and Ke-Qin Zhang. 2007. Bacteria used in the biological control of plant parasitic nematodes: populations, mechanism of action and future prospects. *FEMS Microbiol. Ecol.*, 61: 197-213.
- Bonants, P.J.M., F.L.F. Paul, H. Thijs, E.D. Belder, C. Waalwijk and D.M.H. Willen. 1994. A basic serine protease from *Paecilomyces lilacinus* with biological activity against *Meloidogyne hapla* eggs. *J. Microbiol.*, 141: 775-784.
- Chang, I and T. Kommedahl. 1968. Biological control of seedling blight of corn by coating kernels with antagonistic microorganisms. *Phytopathology.*, 58: 1395-1401.
- Dawar, S., M. Tariq and M.J. Zaki. 2008. Application of *Bacillus* species in control of *Meloidogyne javanica* (Treub) Chitwood on cowpea and mash bean. *Pak. J. Bot.*, 40(1): 439-444.
- Denis, C and J. Webster. 1971. Antagonistic properties of species group of *Trichoderma*. II- Production of volatile antibiotics. *Trans. Brit. Mycol. Soc.*, 57: 41-48.
- Devidas, P. and L.A. Rehberger. 1992. The effect of exotoxin (Thurigiensin) from *Bacillus thuringiensis* on *Meloidogyne incognita* and *Caenorhabditis elegans*. *Plant Soil*, 145: 115-120.
- Eisenback, J.D. and H.H. Triantaphyllous. 1991. Root knot nematode *Meloidogyne* species and races In: *Manual of Agriculture Nematology*, (Ed). W.R. Nickle. Marcel Dekker, New York. pp. 281-286.
- Elad, Y., I. Chet and Y. Hennis. 1982. Degradation of plant pathogenic fungi by *Trichoderma harzianum*. *Can. J. Microbiol.*, 28: 719-725.
- Goodey, J.B., M.T. Franklin and D.J. Hooper. 1965. *Goodey's the nematode parasite of plant Cataloged under their hosts*. Farnham Royal, Commonwealth Agriculture Bureax. 3rd edition, pp. 214.

- Hussey R.S. and K.R. Barker. 1973. A comparison of methods of collecting inocula of *Meloidogyne* spp, including a new technique *Pl. Dis. Report*, 57: 1025-1028.
- Keen, B.A. and H. Raczkowski. 1992. Clay contents and certain physical properties of soil. *J. Agric. Sci.*, 11: 441-449.
- Kerry, B.R. 1987. *Biological control*. In: *Biological control of nematodes: prospects and opportunities, Principles and practice of nematode control in crops*. (Eds.) R.H. Brown & B.R. Kerry. Sydney, Australia, Academic Press, pp. 233-263.
- Luc, M., A.R. Magneetic and R. Fortuner. 1988. A reappraisal of *Tylenchia* (Nematol) 9. The family Heteroidae Filip' ve and Schuurmans Stekhovem, 1941. *Revue Nematol.*, 11: 159-176.
- Santos, D.M.A., S. Ferraz and J.J. Muchovej. 1992. Evaluation of 20 species of fungi from Brazil for biocontrol of *Meloidogyne incognita* race-3. *Nematropica.*, 22: 183-192.
- Sasser, J.N. and D.W. Freckman. 1987. A world prospective on nematology. The role of society. In: *Vistas in Nematology*. (Ed): A. Veech and D.W. Dickerson. Hyallsville. Society of Nematologists, pp. 7-14.
- Sayre, R.M. and D.E. Walter. 1991. Factors affecting the efficacy of natural enemies of nematodes. *Ann. Rev. Phytopathol.*, 29: 149-166.
- Schneider, S.M. 1991. Root knot nematodes. In: *Compendium of Tobacco Diseases*. (Eds): H.D. Shew and G.B. Lucas. St. Paul, M.N: APS Press, pp. 37-40.
- Seuk Bae, Yeoung, OK-Hee Choi, Kyung-Soek park, Sang-Bum Lee and Choong-Hoe Kim. 2000. *A useful method for functional analysis of plant growth promoting rhizobacteria in the development of cucumber root system*. Plant pathology Division, National Institute of Agricultural Science and Technology, Suwon, 441-707, Korea.
- Siddique, I.A, S.A Qureshi, V. Sultana, S. Ehteshamul-Haque and A. Ghaffar. 2000. Biological control of root rot and root knot disease complex of tomato. *Plant Soil*, 227: 163-169.
- Siddiqui, I.A., S.S. Shaukat and A. Khan. 2003. Differential impact of some *Aspergillus* species on *Meloidogyne javanica* biocontrol by *Pseudomonas fluorescens* strain CHA0. *Letters in Applied Microbiology*, 39(1): 74-83.
- Siddiqui, Z.A. and I. Mahmood. 1995. Some observations on the management of the wilt disease complex on pigeon pea by treatment with vascular arbuscular fungus and biocontrol agent for nematodes. *Bioremediation Technology*, 54: 227-230.
- Sikora, R.A. 1992. Management of the antagonistic potential in agricultural ecosystems for the biological control of plant-parasitic nematodes. *Ann. Rev. Phytopathol.*, 30: 245-270.
- Sokal, R.R. and F.J. Rohlf. 1995. *Biometry: The Principals and practices of Statistics in Biological Research*. Freeman, New York. pp. 887.
- Stirling, G.R. 1991. *Biological control of plant-parasitic nematodes*. Wallingford, UK, CAB International. pp. 282.
- Windham, G.L., M.T. Windham and W.P. Williams. 1989. Effect of *Trichoderma* species on maize growth and *Meloidogyne arenaria* reproduction. *Pl. Dis.*, 73: 493-495.
- Zaki, M.J. 2000. *Bio-management of root- knot nematodes problem of vegetables*. DFID, UK Research Project Report. Department of Botany, University of Karachi, Karachi- 75270. pp. 131.

(Received for publication 2 August 2008)