

## RESPONSE OF TOBACCO POLYPHENOL OXIDASE GENE TO WOUNDING, ABSICISIC ACID (ABA) AND METHYL JASMONATE (MeJ)

EJAZ AZIZ<sup>1</sup>, WASIM AKHTAR<sup>2</sup>, MUHAMMAD ILYAS<sup>1</sup>, SHAZIA REHMAN<sup>1</sup>, HISASHAI KOIWA<sup>3</sup>, ZABTA KHAN SHINWARI<sup>4</sup> AND TARIQ MAHMOOD\*<sup>1</sup>

<sup>1</sup>Department of Plant Sciences, Quaid-i-Azam University, Islamabad-45320, Pakistan

<sup>2</sup>Department of Biotechnology, University of Azad Jammu and Kashmir, Muzaffarabad, Pakistan

<sup>3</sup>Department of Horticultural Sciences, MS2133 Texas A&M University 495 Horticulture Rd. College Station, TX77843-2133, USA

<sup>4</sup>Department of Biotechnology, Quaid-i-Azam University, Islamabad-45320, Pakistan

\*Correspondence: tmahmood.qau@gmail.com;tmahmood@qau.edu.pk

Tel. # +92-51-9064 3050, Fax # +92-51-9064 3138

### Abstract

Polyphenol oxidases (*PPOs*) are ubiquitous enzymes in plant kingdom and catalyze the oxidation of phenols to highly reactive quinones. *PPO* genes are induced by both biotic and abiotic stresses and implicated in several physiological processes including plant defense against pathogen and insects. Here, effect of wounding, abscisic acid (ABA) and methyl jasmonate (MeJ) applications on *PPO* induction in *Nicotiana tabacum* (*NtPPO*) was characterized by RT-PCR. In response to MeJ treatment at different concentrations, maximum expression of *NtPPO* genes was observed with six folds induction at 500  $\mu$ M concentration, while target transcript level was relatively low (0.2-0.9 fold) with different concentrations of ABA. Upon wounding, *NtPPO* gene was strongly induced (up to 14 folds) after 36 hours treatment. Thus, wound and MeJ inducibility of *NtPPO* gene is a strong indicative of its role in plant defense mechanism against biotic and abiotic stresses.

**Key words:** Polyphenol oxidase, Wounding, Tobacco, Gene expression.

### Introduction

Plants in nature are continuously exposed to environmental stresses such as drought, salinity, high temperature, which negatively affect the growth and productivity (Shinwari et al. 1998; Mahajan & Tuteja, 2005; Munns, 2005). In response, plants perceive signals that lead to the synthesis of transcription factors and subsequently regulate genes for synthesis of proteins which help to cope with stress (Shinozaki & Yamaguchi-Shinozaki, 2000). The role of polyphenol oxidases (*PPOs*) in plant defense has been implied due to its up-regulation by pathogens (Thipyapong et al., 1997; Mayer, 2006; Raj et al., 2006).

Defensive role of *PPO* has been proposed based on activities of *PPO* reaction products upon wounding, pathogen infection, insect attacks and induction of *PPO* by abiotic, biotic environmental constraints and signaling molecules (Thipyapong et al., 1995; Thipyapong et al., 1997; Maki & Morohashi, 2006; Mahmood et al., 2015; Akhtar & Mahmood, 2017). The role of *PPO* in inhibition of postharvest proteolysis further made it significant (Sullivan et al., 2004; Sullivan & Hatfield, 2006). The *PPO* activity ultimately results quinones and reactive oxygen species to cope with stresses (Thipyapong et al., 2007), and there are few studies on function of *PPO* for abiotic stress tolerance. Stresses signal transduction pathway such as abscisic acid (ABA), methyl jasmonate (MeJ), ethylene and systemin signals can trigger *PPO* expression. Chewing insects amplify wounding signals which is thought to be mediated by jasmonic acid (JA) and generation of reactive oxygen species (ROS) and is responsible for restriction of pathogen spread to the other parts of plant (Torres et al., 2002).  
C:\Users\Rehan\TOBACCO\Tobacco PPO paper.docx - ENREF 17*PPO* genes are induced by wounding and/or MeJ (Constabel et al., 2000; Zhou et al., 2003; Wang & Constabel, 2004; Raimbault et al., 2010; Shetty et al., 2011) as well as during wound sealing (Wahler et al., 2009).

In *Solanaceae* family, potato, tomato, eggplant and lulo, *PPO* genes were characterized in abiotic and biotic stresses (Li & Steffens, 2002; Thipyapong et al., 2004; Mahanil et

al., 2008; Shetty et al., 2011; Arias et al., 2012; Chi et al., 2014). To understand the role of *PPO* in plant defense and to various stresses e.g. wounding, MeJ and ABA application, a study was designed to characterize the endogenous tobacco (*Nicotiana tabacum*) *PPO* gene expression.

### Materials and Methods

**Plant growth:** Tobacco seeds (*N. tabacum* var. *Xanthi*) were grown on half strength Murashige and Skoog (MS) medium (Phyto Technology Laboratories) (Murashige & Skoog, 1962). Seedlings were grown in growth room at 25°C, 50% humidity and 16/8 light cycle. Seven days old plants were used for wounding treatment, MeJ and ABA applications.

**Wound induction:** Seedlings of *N. tabacum* were subjected to mechanical wounding by forceps. Leaves on each plant were injured carefully. Wounded plants were kept on MS media for 12, 24, 36 and 48 hours respectively along with controlled unwounded plants.

**ABA and MeJ treatments:** Solutions of ABA (Sigma-Aldrich) and MeJ (Sigma-Aldrich) were prepared from 1000 folds concentrated stock in combination with 0.01% SilvetL-77 (Merck) to facilitate infusion. Control solution (distilled water in combination with 0.01% SilvetL-77) was also used. Seven days old tobacco plants were sprayed with 100  $\mu$ M, 200  $\mu$ M, 300  $\mu$ M, 400  $\mu$ M and 500  $\mu$ M ABA and MeJ solutions and placed in growth room for 24 hours. Control plants were sprayed with control solution.

**Primer designing:** Primers were designed from coding region of *PPO* gene (A27686.1) sequence of *N. tabacum* by ApE (A plasmid Editor) software. Sequence of primers is given as:

*NtPPO*2F: AACCCGTTCCGTGTGAAAGTCC

*NtPPO*2R: CTTCGATTACGCACCGATGCCA




Fig. 1. Quantification of *NtPPO* mRNA level induced by wounding. Seven days old tobacco plants were subjected to mechanical injury growing on MS media and RT-PCR was performed after intervals of 12, 24, 36 and 48 hours to detect wound induced response of *NtPPO* gene.




Fig. 2. RT-PCR analysis of *NtPPO* gene activity by MeJ applications. Quantitative RT-PCR was carried out to detect *NtPPO* transcripts level in seven days old tobacco plants by sprays of MeJ solutions (100 µM, 200 µM, 300 µM, 400 µM and 500 µM) growing on MS media.




Fig. 3. RT-PCR analysis of *NtPPO* gene activity by ABA applications. Quantitative RT-PCR was carried out to detect *NtPPO* transcripts level in seven days old tobacco plants by sprays of ABA solutions (100 µM, 200 µM, 300 µM, 400 µM and 500 µM) growing on MS media.

**RNA isolation and cDNA synthesis:** Total RNA was extracted from the ABA, MeJ and wound stressed plants along with control plants using Spectrum™ Plant Total RNA kit (Sigma). Reverse transcription was carried out using 3 µg of total RNA to synthesize cDNA using Superscript II reverse transcriptase (Invitrogen).

***NtPPO* gene expression analysis:** Quantification of *PPO* transcripts was performed in ViiA™7 Real-Time PCR system with the Power SYBR® Green master mix (Applied Biosystems, Carlsbad, CA, USA) according to the manufacturer's instructions. *PPO* gene was used as target and 18S gene was used as internal control. After successful amplification of targets, the cycle threshold (Ct) values were exported from the ViiATM 7 software and used as raw data for the analysis of qRT-PCR data. The R software and the HTqPCR (Dvinge & Bertone, 2009) and Limma (Gentleman *et al.*, 2006) add-on packages were used for the manipulation and analysis of the Ct values.

## Results

**Induction of *NtPPO* by wounding:** Effect of wounding was monitored in seven days old tobacco plants growing on half strength MS medium. Young leaves were mechanically wounded which simulates the attack of insect pests. To analyze the expression of *NtPPO* gene in response to wounding; transcript levels of *NtPPO* endogenous gene was in increasing trend with the passage of time after wounding. Transcript level was found to increase after 12 hours (6 folds) reaching to the maximum level of 14 folds after 36 hours of wounding treatment (Fig. 1), which started to decline after 48 hours.

**Induction of *NtPPO* transcript in response to ABA and MeJ:** Plants were sprayed with various concentrations of ABA and MeJ, left for 24 hours in growth chamber. Transcript level of *NtPPO* was up-regulated from 2-6 folds in response to MeJ treatment (Fig. 2). An increase in *NtPPO* transcript level was found with increasing MeJ concentrations. It was seen that mRNA level increased up to 2 folds upon 100 µM MeJ and maximum induction of transcript level was observed at 500 µM MeJ. Unlike MeJ, lower level of *NtPPO* expression was observed with ABA application (Fig. 3).

## Discussion

*PPOs* are generally related to plant defense (Mayer, 2006; Constabel & Barbehenn, 2008). In comparisons with our findings, *PPO* mRNA accumulation increased in response to wounding and defense related hormones SA and MeJ in poplar (Mayer, 2006; Flurkey & Inlow, 2008). Disease susceptibility and herbivory were also strongly linked to *PPO* activities (Thipyapong *et al.*, 2007). Based on the previous reports, it is well known that *PPOs* are part of innate immunity in plants (Fuerst *et al.*, 2014).

**Induction of *NtPPO* gene by wounding:** Some of plant defenses are triggered through tissue damage mechanical injury (Howe & Jander, 2008). In tobacco, our data has

indicated that an endogenous induction of *PPO* gene by wounding suggests a possible role of *NtPPO* gene in plant defense. Accumulation of *NtPPO* mRNA up to 14 fold by wounding is in agreement to high induction of potato *PPO* activity (Thipyapong *et al.*, 1995). In earlier reports, localized wounding of lateral roots (Thipyapong *et al.*, 1997), wounding stress (Quarta *et al.*, 2013) and chilling injury in pineapple resulted in *PPO* gene activation (Raimbault *et al.*, 2010).

Role of *PPO* has been evaluated in anti-herbivory. *NtPPO* induction by wounding is probably the first possible indication of its defensive role which is in consistent with hybrid poplar *PPO* transcripts activated by wounding, as well as real insect herbivory (Constabel *et al.*, 2000). In aspen, forest tent caterpillar strongly induced *PPO* expression as did by mechanical wounding (Haruta *et al.*, 2001). *PPO* has a strong correlation in latex coagulation and wound sealing in dandelions (Wahler *et al.*, 2009). The pattern of *NtPPO* induction by wounding could be explained by responsiveness to insect herbivory as stimulation and *PPO* mRNA accumulated mainly in eggplant stem and fruits (Shetty *et al.*, 2011). In another report, strong up-regulation of *PPO* gene in rubber plant by wounding may correlate its probable role in plants defense (Li *et al.*, 2014).

#### Induction of *NtPPO* transcript in response to ABA and MeJ:

*PPO* has been signaled by various signaling molecules such as ethylene (Newman *et al.*, 1993), ABA (Chai *et al.*, 2013; Kaur & Zhawar, 2015; Jia *et al.*, 2016) and JA (Adeloye & Ajibade, 2011; Jia *et al.*, 2016) indicating role of *PPO* in plant defense. It is well known that MeJ induced *PPO* activity along with other defense related responses (Thaler *et al.*, 2002). Jasmonates take part in well-timed induction of plant defense responses against insects as activation of plant defense depends upon jasmonate pathway (Erb *et al.*, 2012; Bosch *et al.*, 2014). It has been revealed that treatment by JA and wounding by insects induce *PPO* activity triggering other responses which result in broad spectrum resistance against pests and microbes (Cooper *et al.*, 2004).

*PPO* up-regulation by jasmonic acid and its overexpression improved resistance against armyworm (*Spodoptera exigua*) defense in tomato (Bosch *et al.*, 2014) which can be correlated to *NtPPO* gene up-regulation by MeJ. An induction of up-regulation of *PPO* mRNA in tobacco by mechanical injury and MeJ might be a possible clue of *NtPPO* participation in biotic stresses and these findings are in consistent with banana *PPO* up-regulation by wounding and MeJ (Sreedharan *et al.*, 2012). It was reported that MeJ induced six *PPO* genes in eggplant and in transgenic tobacco (Shetty *et al.*, 2012). In another report, it was shown that red Swiss chard *PPO* promoter was regulated by MeJ in transgenic *Arabidopsis* (Yu *et al.*, 2015). Many plant's *PPOs* were regulated by MeJ treatment indicating the role of *PPO* in plant defense such as in hybrid poplar (Constabel *et al.*, 2000), trembling aspen (Haruta *et al.*, 2001), tomato (Li & Steffens, 2002), walnut (Escobar *et al.*, 2008), rubber tree (Li *et al.*, 2014) and strawberry (Jia *et al.*, 2016).

#### Conclusion

From the data of wounding, ABA and MeJ treatments, it was revealed that *NtPPO* might have role in plant protection against both biotic and abiotic stresses. These results are helpful in potential understanding of plant responses to such stresses. Conclusively, *NtPPO* is a potential candidate gene for developing transgenic crops tolerant to both biotic and abiotic stresses.

#### Acknowledgments

We are thankful to Pakistan Science Foundation, Islamabad for providing financial assistance through a research project (NSLP-77).

#### References

- Adeloye, A.O. and P.A. Ajibade. 2011. A high molar extinction coefficient mono-anthracenyl bipyridyl heteroleptic ruthenium (II) complex: Synthesis, photophysical and electrochemical properties. *Molecules*, 16(6): 4615-4631.
- Akhtar, W. and T. Mahmood. 2017. Response of rice polyphenol oxidase promoter to drought and salt stress. *Pak. J. Bot.*, 49(1): 21-23.
- Arias, D.G., C.M.M. Doria, L.R. Ramos and H.C.N. Morocho. 2012. Molecular characterization of the polyphenol oxidase gene in lulo (*Solanum quitoense* Lam.) var. Castilla. *Braz. J. Plant Physiol.*, 24(4): 261-272.
- Bosch, M., S. Berger, A. Schaller and A. Stintzi. 2014. Jasmonate-dependent induction of polyphenol oxidase activity in tomato foliage is important for defense against *Spodoptera exigua* but not against *Manduca sexta*. *BMC Plant Biol.*, 14(1): 257.
- Chai, C., Y. Lin, D. Shen, Y. Wu, H. Li and D. Dou. 2013. Identification and functional characterization of the Soybean GmaPPO12 promoter conferring *Phytophthora sojae* induced expression. *PLoS One*, 8(6): e67670.
- Chi, M., B. Bhagwat, W.D. Lane, G. Tang, Y. Su, R. Sun, B.D. Oomah, P.A. Wiersma and Y. Xiang. 2014. Reduced polyphenol oxidase gene expression and enzymatic browning in potato (*Solanum tuberosum* L.) with artificial microRNAs. *BMC Plant Biol.*, 14(1): 1.
- Constabel, C.P. and R. Barbehenn. 2008. Defensive roles of polyphenol oxidase in plants. Induced plant resistance to herbivory. Springer, pp. 253-270.
- Constabel, C.P., L. Yip, J.J. Patton and M.E. Christopher. 2000. Polyphenol oxidase from hybrid poplar. Cloning and expression in response to wounding and herbivory. *Plant Physiol.*, 124(1): 285-296.
- Cooper, W., L. Jia and F. Goggin. 2004. Acquired and R-gene-mediated resistance against the potato aphid in tomato. *J. Chem. Ecol.*, 30(12): 2527-2542.
- Dvigne, H. and P. Bertone. 2009. HTqPCR: high-throughput analysis and visualization of quantitative real-time PCR data in R. *Bioinformatics*, 25(24): 3325-3326.
- Erb, M., S. Meldau and G.A. Howe. 2012. Role of phytohormones in insect-specific plant reactions. *Trends Plant Sci.*, 17(5): 250-259.
- Escobar, M.A., A. Shilling, P. Higgins, S.L. Uratsu and A.M. Dandekar. 2008. Characterization of polyphenol oxidase from walnut. *J. Am. Soc. Hort. Sci.*, 133(6): 852-858.
- Flurkey, W.H. and J.K. Inlow. 2008. Proteolytic processing of polyphenol oxidase from plants and fungi. *J. Inorg. Biochem.*, 102(12): 2160-2170.
- Fuerst, E.P., P.A. Okubara, J.V. Anderson and C.F. Morris. 2014. Polyphenol oxidase as a biochemical seed defense mechanism. *Front. Plant. Sci.*, 5: 689.

- Gentleman, R., V. Carey, W. Huber, R. Irizarry and S. Dudoit. 2006. Bioinformatics and computational biology solutions using R and Bioconductor. Springer Science & Business Media. 397-420.
- Haruta, M., J.A. Pedersen and C.P. Constabel. 2001. Polyphenol oxidase and herbivore defense in trembling aspen (*Populus tremuloides*): cDNA cloning, expression, and potential substrates. *Physiol. Plant.*, 112(4): 552-558.
- Howe, G.A. and G. Jander. 2008. Plant immunity to insect herbivores. *Annu. Rev. Plant Biol.*, 59: 41-66.
- Jia, H.F., P.C. Zhao, B.J. Wang, P. Tariq, F.G. Zhao, M.Z. Zhao, Q.L. Wang, T.B. Yang and J.G. Fang. 2016. Overexpression of polyphenol oxidase gene in strawberry fruit delays the fungus infection process. *Plant Mol. Biol. Rep.*, 34(3): 592-606.
- Kaur, L. and V.K. Zhawar. 2015. Phenolic parameters under exogenous ABA, water stress, salt stress in two wheat cultivars varying in drought tolerance. *Indian. J. Plant Physiol.*, 20(2): 151-156.
- Li, D., Z. Deng, C. Liu, M. Zhao, H. Guo, Z. Xia and H. Liu. 2014. Molecular cloning, expression profiles, and characterization of a novel polyphenol oxidase (PPO) gene in *Hevea brasiliensis*. *Biosci. Biotechnol. Biochem.*, 78(10): 1648-1655.
- Li, L. and J.C. Steffens. 2002. Overexpression of polyphenol oxidase in transgenic tomato plants results in enhanced bacterial disease resistance. *Planta*, 215(2): 239-247.
- Mahajan, S. and N. Tuteja. 2005. Cold, salinity and drought stresses: an overview. *Arch. Biochem. Biophys.*, 444(2): 139-158.
- Mahanil, S., J. Attajarusit, M.J. Stout and P. Thipyapong. 2008. Overexpression of tomato polyphenol oxidase increases resistance to common cutworm. *Plant Sci.*, 174(4): 456-466.
- Mahmood, T., M.U. Rehman, E. Aziz, I. Ali and Z.K. Shinwari. 2015. *In silico* analysis, mapping of regulatory elements and corresponding DNA-protein interaction in polyphenol oxidase gene promoter from different rice varieties. *Pak. J. Bot.*, 47(6): 2321-2327.
- Maki, H. and Y. Morohashi. 2006. Development of polyphenol oxidase activity in the micropylar endosperm of tomato seeds. *J. Plant Physiol.*, 163(1): 1-10.
- Mayer, A.M. 2006. Polyphenol oxidases in plants and fungi: going places? A review. *Phytochemistry*, 67(21): 2318-2331.
- Munns, R. 2005. Genes and salt tolerance: bringing them together. *New Phytol.*, 167(3): 645-663.
- Murashige, T. and F. Skoog. 1962. A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiol. Plant.*, 15(3): 473-497.
- Newman, S.M., N.T. Eannetta, H. Yu, J.P. Prince, M.C. de Vicente, S.D. Tanksley and J.C. Steffens. 1993. Organisation of the tomato polyphenol oxidase gene family. *Plant Mol. Biol.*, 21(6): 1035-1051.
- Quarta, A., G. Mita, M. Durante, M. Arlorio and A. De Paolis. 2013. Isolation of a polyphenol oxidase (PPO) cDNA from artichoke and expression analysis in wounded artichoke heads. *Plant Physiol. Biochem.*, 68: 52-60.
- Raimbault, A.-K., P.-A. Marie-Alphonsine, J.-P. Horry, M. Francois-Haugrin, K. Romuald and A. Soler. 2010. Polyphenol oxidase and peroxidase expression in four pineapple varieties (*Ananas comosus* L.) after a chilling injury. *J. Agric. Food Chem.*, 59(1): 342-348.
- Raj, S.N., B. Sarosh and H.S. Shetty. 2006. Induction and accumulation of polyphenol oxidase activities as implicated in development of resistance against pearl millet downy mildew disease. *Funct. Plant Biol.*, 33(6): 563-571.
- Shetty, S.M., A. Chandrashekar and Y.P. Venkatesh. 2011. Eggplant polyphenol oxidase multigene family: cloning, phylogeny, expression analyses and immunolocalization in response to wounding. *Phytochemistry*, 72(18): 2275-2287.
- Shetty, S.M., A. Chandrashekar and Y.P. Venkatesh. 2012. Promoter analyses and transcriptional profiling of eggplant polyphenol oxidase 1 gene (SmePPO1) reveal differential response to exogenous methyl jasmonate and salicylic acid. *J. Plant Physiol.*, 169(7): 718-730.
- Shinozaki, K. and K. Yamaguchi-Shinozaki. 2000. Molecular responses to dehydration and low temperature: differences and cross-talk between two stress signaling pathways. *Curr. Opin. Plant Biol.*, 3(3): 217-223.
- Shinwari, Z.K., Nakashima K, Miura S., Kasuga M., Seki M., Yamaguchi-Shinozaki K, and Shinozaki K 1998. An Arabidopsis gene family encoding DRE binding protein involved in low temperature - responsive gene expression. *Biochem. Biophys. Res. Commun.*, 250: 161-170.
- Sreedharan, S., U.K.S. Shekhawat and T.R. Ganapathi. 2012. MusaSAP1, a A20/AN1 zinc finger gene from banana functions as a positive regulator in different stress responses. *Plant Mol. Biol.*, 80(4-5): 503-517.
- Sullivan, M.L. and R.D. Hatfield. 2006. Polyphenol oxidase and-diphenols inhibit postharvest proteolysis in red clover and alfalfa. *Crop Sci.*, 46(2): 662-670.
- Sullivan, M.L., R.D. Hatfield, S.L. Thoma and D.A. Samac. 2004. Cloning and characterization of red clover polyphenol oxidase cDNAs and expression of active protein in *Escherichia coli* and transgenic alfalfa. *Plant Physiol.*, 136(2): 3234-3244.
- Thaler, J.S., R. Karban, D.E. Ullman, K. Boege and R.M. Bostock. 2002. Cross-talk between jasmonate and salicylate plant defense pathways: effects on several plant parasites. *Oecologia*, 131(2): 227-235.
- Thipyapong, P., M.D. Hunt and J.C. Steffens. 1995. Systemic wound induction of potato (*Solanum tuberosum*) polyphenol oxidase. *Phytochemistry*, 40(3): 673-676.
- Thipyapong, P., D.M. Joel and J.C. Steffens. 1997. Differential expression and turnover of the tomato polyphenol oxidase gene family during vegetative and reproductive development. *Plant Physiol.*, 113(3): 707-718.
- Thipyapong, P., S. Mahanil, A. Bhonwong, J. Attajarusit, M. Stout and J. Steffens. 2004. Increasing resistance of tomato to lepidopteran insects by overexpression of polyphenol oxidase, IX International Symposium on the Processing Tomato 724, pp. 29-38.
- Thipyapong, P., M.J. Stout and J. Attajarusit. 2007. Functional analysis of polyphenol oxidases by antisense/sense technology. *Molecules*, 12(8): 1569-1595.
- Torres, M.A., J.L. Dangel and J.D. Jones. 2002. Arabidopsis gp91phox homologues AtrbohD and AtrbohF are required for accumulation of reactive oxygen intermediates in the plant defense response. *Proceedings of the National Academy of Sciences*, 99(1): 517-522.
- Wahler, D., C.S. Gronover, C. Richter, F. Foucu, R.M. Twyman, B.M. Moerschbacher, R. Fischer, J. Muth and D. Prüfer. 2009. Polyphenoloxidase silencing affects latex coagulation in *Taraxacum* species. *Plant Physiol.*, 151(1): 334-346.
- Wang, J. and C.P. Constabel. 2004. Polyphenol oxidase overexpression in transgenic *Populus* enhances resistance to herbivory by forest tent caterpillar (*Malacosoma disstria*). *Planta*, 220(1): 87-96.
- Yu, Z.-H., Y.-N. Han and X.-G. Xiao. 2015. A PPO Promoter from betalain-producing red swiss chard, directs petiole- and root-preferential expression of foreign gene in anthocyanins-producing plants. *Int.J.Mol. Sci.*, 16(11): 27032-27043.
- Zhou, Y., T.J. O'Hare, M. Jobin-Decor, S.J. Underhill, R.B. Wills and M.W. Graham. 2003. Transcriptional regulation of a pineapple polyphenol oxidase gene and its relationship to blackheart. *Plant Biotechnol. J.*, 1(6): 463-478.