

SYNCHRONIZATION OF ZINC RATES THROUGH SOIL AND FOLIAR APPLICATION WITH SYNTHETIC FERTILIZER TO MAXIMIZE GROWTH, YIELD AND QUALITY ATTRIBUTES OF MANGO

MUHAMMAD MEHBOOB HASSAN KHAN¹ AND NIAZ AHMED^{1*}

¹Department of Soil Science, Faculty of Agricultural Sciences and Technology, Bahauddin Zakariya University, Multan, Punjab, 60800, Pakistan

*Corresponding author's email: niaz.ahmad@bzu.edu.pk

Abstract

Zinc (Zn) deficiency is among the major constraints for reduced fruit production and quality in mango orchards soils. Therefore, the judicious use of Zn in mango orchards is gaining attention. Although, the farming communities are using Zn as a micronutrient fertilizer; however, the synchronization of inorganic fertilizers and Zn application rate is yet to be performed under changing climatic conditions. Therefore, the current experiment was performed to assess better application methods (soil and foliar application) and judicious rates of Zn with recommended doses of nitrogen (N), phosphorus (P) and potassium (K) for mango trees. Zinc was applied as ZnSO₄ at the rate of 0, 150, and 300 g plant⁻¹ as basal application and 0, 0.5, and 1.0% as a foliar application. Results confirmed with an increase in soil and foliar-applied Zn solely, growth, quality and yield of mango fruit improved. Combined use of soil and foliar Zn gave more positive effects over sole application. Treatment T₇ (recommended NPK + ZnSO₄ (150 g plant⁻¹ as basal and 1.0% as foliar spray)) followed by T₈ (recommended NPK + ZnSO₄ (300 g plant⁻¹ as basal and 1.0% as foliar spray)) were statistically alike for the improvement in yield per tree, photosynthetic rate, transpiration rate and sugar contents. In conclusion, recommended NPK + ZnSO₄ (150 g plant⁻¹ as basal and 1.0% as foliar spray) is a better approach for maximizing the yield, growth attributes and quality of mango fruits.

Key words: Growth attributes, Gas exchange attributes, *Mangifera indica* L., Zinc

Introduction

Mango (*Mangifera indica* L.) is cultivated in nearly 63 countries worldwide. It has been documented that 69 species of genus *Mangifera* are found worldwide and there are 11,595 varieties available on the globe (Bose, 1999). Single mango fruit can deliver up to 40% of everyday dietary fiber requirements (Sing *et al.*, 2005). It is a tropical, climacteric fruit liked by all due to its taste, flavor and excellent nutritional properties (Baloch *et al.*, 2017). The mango is the most important fruit of Asia. Pakistan ranks fourth in the total production among major fruit crops worldwide after bananas, citrus, grapes and apples. Pakistani mango is recognized as one of the best of its kind in the world market. Pakistan is among the leading mango producers and is considered the original and natural habitat of mangoes (Madonsela, 2019). Pakistan was the 2nd biggest producer of mango in the world during 1970's (Usman *et al.*, 2003); however, its position is dropped to 4th (Anon., 2010; Memon, 2016) with 1,732 thousand tones annual production under 167.5 thousand hectares of area (Ahmed *et al.*, 2020). The reduction in mango yield and quality is a matter of great concern. Multiple factors or their combination might have contributed to yield and quality reduction. The major factors that can decrease the mango yield are poor fruit setting, malnutrition, plant and irrigation protection measures, poor management of plant, for instance, no pruning, intercropping, dieback, mango malformation, mango sudden death, alternate bearing and poor post-harvest handling (Usman *et al.*, 2003; Kazmi *et al.*, 2005; Sharma & Singh, 2006). Moreover, mango undergoes numerous disorders like blacktip, internal necrosis, fruit cracking, jelly seed, spongy tissue, leaf scorching, and fruit pitting which further contributed to the loss in yield and quality (Sharma, 2006; Saran & Kumar, 2011).

The use of micronutrients is very important for the achievement of optimum crops yield. These are involved in enzymatic activities, hormones biosynthesis and assimilation. Deficiency of micronutrients occurs in various areas and the majority of the mango growers do not follow the application of micronutrients and eventually, orchards show the hunger signs (Fageria, 2009). The micronutrients are required only in traces from soil or by using chemical sources. The primary reasons for micronutrient deficiencies are intensive agriculture, unbalanced use of inorganic fertilizers, exhaustion of nutrients, and their inadequate supply (Rajendran & Ramamurthy, 2009). The micronutrient deficiency in the mango crop caused poor fruit quality. Low fruit sets and unnecessary fruits drop at numerous stages of fruit development (Siddiq *et al.*, 2017). Soils of Pakistan are mostly alkaline, and excessive phosphatic fertilizers can cause an increased level of phosphorous concentration in the soils. Such increase disturbs the solubility and mobility of zinc (Zn) in soil and soils become Zn deficient (Akhtar *et al.*, 2019).

When Zn deficiency occurs, plants undergo physiological stresses, which caused abnormalities in crops and then deficiency symptoms. In case of acute deficiency of Zn, plants showed evident symptoms comprising stunted growth, small leaves, spikelet sterility and chlorosis of leaves. Acute deficiency of Zn in field crops has been reported in the literature since 1937 (Brown *et al.*, 1993). However, in such condition, the quality of produce, i.e., protein contents, size, and appearance of fruit, may undergo harmful effects and susceptibility increased by injury to light intensity, temperature, and infection of certain fungal diseases (Cakmak, 2000). In hidden deficient conditions of Zn, reduction in crop yields occurred and quality also suffered without the presence of obvious symptoms

Madonsela (2019). Zn regulates growth of plant by reducing the water stresses and tackling the conditions of drought faced by plants (Jan *et al.*, 2019). The current study was designed to explore the optimum application (soil basal dose and foliar) rate of ZnSO₄ with inorganic fertilizer. It is hypothesized that combined application of Zn i.e., as basal soil dose and foliar is a better approach than sole application for improving the quality and yield of mango.

Materials and Methods

Experimental site: The experiment was done in a mango orchard (30°09' N, 71°26' E) at Multan. Three-years mean annual maximum and minimum temperatures were 35.8 and 10.3 °C with precipitation 82 mm.

Soil characteristics: Experimental soil texture was silt loam having calcareous parent material with pH 8.1. Zinc (Zn) in soil was 0.49 ug g⁻¹ with 6–10% CaCO₃ equivalent.

Treatments plan: The experiment comprises of 9 treatments and three replications per treatment, where a tree was experimental unit with RCBD arrangement. The following treatments were applied for Zn nutrient management; T₀: recommended dose (RD) of NPK + ZnSO₄ (0 g plant⁻¹ as basal and 0% as foliar spray), T₁: RD + ZnSO₄ (150 g plant⁻¹ as basal and 0% as foliar spray), T₂: RD + ZnSO₄ (300 g plant⁻¹ as basal and 0% as foliar spray), T₃: RD + ZnSO₄ (0 g plant⁻¹ as basal and 0.5% as foliar spray), T₄: RD + ZnSO₄ (150 g plant⁻¹ as basal and 0.5% as foliar spray), T₅: RD + ZnSO₄ (300 g plant⁻¹ as basal and 0.5% as foliar spray), T₆: RD + ZnSO₄ (0 g plant⁻¹ as basal and 1.0% as foliar spray), T₇: RD + ZnSO₄ (150 g plant⁻¹ as basal and 1.0% as foliar spray), and T₈: RD + ZnSO₄ (300 g plant⁻¹ as basal and 1.0% as foliar spray).

Zinc application methods and time: Zn fertilizers were applied into the soil prior to the flowering stage in mango trees. While, the foliar application of Zn was performed twice a year. The first foliar application was made at the time of the inflorescence stage. However, a second foliar spray was applied when fruit size becomes like a pea.

Data collection: After harvesting, soil and leaf samples were collected, stored and processed for further analyses. Attributes such as average fruit weight, fruit size, number of flowers per panicle, number of fruits matured per panicle, fresh pulp weight, stone weight, shelf life and yield per tree were noted as per standard procedures after harvesting.

Sugar contents: Total sugars and Non-reducing sugars were determined according to the method of Ranganna (1986). Determination of reducing sugars was according to the method described by Hortwitz (1960).

Statistical analysis: Collected data were examined statistically by applying analysis of variance (ANOVA) and Tukey's test at $p \leq 0.05$. Statistical computer

software package (Statistics 8.1) was used for the statistical analysis.

Results

Average fruit weight, fruit length and fruit width: The effect of treatments was significant ordinal for average fruit weight (AFW), fruit length (FL) and fruit width (FW). The effect of treatments was significant for the improvement in AFW, FL and FW. It was observed that all the treatments were significant from control for the improvement in AFW and FW. Treatments T₃, T₄, T₅, T₆, T₇ and T₈ differed significantly over control for the improvement in FL (Table 1). However, T₁ and T₂ did not differ over control for FL. Chord diagram showed that the percentage share of treatment T₇ was highest than T₀ for average fruit weight, fruit width and fruit length (Fig. 1). Pearson correlation showed a significant positive correlation between treatments, AFW, FL, and FW (Fig. 6). The maximum increase in AFW, FL and FW were noted where T₇ was applied.

Shelf life, number of flowers and fruits matured panicle⁻¹: Application of treatments significantly affected the shelf life (SL), number of flowers per panicle (NFP) and number of fruits matured per panicle (NFMP). Effect of treatments was significant for the improvement in SL, NFP and NFMP. It was observed that all the treatments were significant from control for the improvement in SL and NFMP. Treatments T₄, T₅, T₆, T₇ and T₈ differed significantly over control for the improvement in NFP (Table 2). However, T₁, T₂ and T₃ did not differ over control for NFP. Chord diagram showed that the percentage share of treatment T₇ was highest than T₀ for shelf life, number of flowers per panicle and number of fruits matured per panicle of mango (Fig. 2). Pearson correlation showed a significant positive correlation between treatments, SL, NFP and NFMP (Fig. 6). Maximum increase in SL, NFP and NFMP were noted where T₇ was applied.

Pulp fresh weight, stone weight and yield tree⁻¹ of mango: Treatments differed significantly for pulp fresh weight (PFW) stone weight (SW) and yield per tree of mango (YT). Impact of treatments was significant for the improvement in PFW, SW and YT. It was observed that all the treatments were significant from control for the improvement in PFW and SW. Treatments T₂, T₃, T₄, T₅, T₆, T₇ and T₈ differed significantly over control for the improvement in YT (Table 3). However, T₁ did not differ over control for YT. Chord diagram showed that the percentage share of treatment T₇ was highest than T₀ for shelf life, number of flowers per panicle and number of fruits matured per panicle of mango. Chord diagram showed that the percentage share of treatment T₇ was highest than T₀ for pulp fresh weight, stone weight and yield per tree of mango (Fig. 3). Pearson correlation showed a significant positive correlation between treatments, PFW, SW, and YT (Fig. 6). The maximum increase in PFW, SW and YT were noted where T₇ was applied.

Table 1. Effects of Zn as foliar and soil application on average fruit weight, fruit length and fruit width of mango.

Treatment	Average fruit weight	Fruit length	Fruit width
	(g)	(cm)	(cm)
T ₀	450 g	11.25 d	5.63 f
T ₁	480 f	12.00 cd	6.00 e
T ₂	499 e	12.475 b-d	6.24 de
T ₃	512 de	12.80 bc	6.40 cd
T ₄	525 cd	13.125 a-c	6.56 b-d
T ₅	531 c	13.275a-c	6.64 bc
T ₆	550 b	13.75 ab	6.88 ab
T ₇	574 a	14.35 a	7.18 a
T ₈	570 a	14.25a	7.13 a

Table 2. Effects of Zn as foliar and soil application on Shelf life, Number of flowers per panicle and Number of fruits matured per panicle of mango.

Treatment	Shelf life	Number of flowers per panicle	Number of fruits matured per panicle
	(Days)	-	-
T ₀	8 c	630.00 e	1.80 g
T ₁	10 b	672.00 de	1.92 f
T ₂	10 b	698.60 cd	2.00 e
T ₃	11 ab	716.80 cd	2.05 de
T ₄	11 ab	735.00 bc	2.10 cd
T ₅	11 ab	743.40 bc	2.12 c
T ₆	11 ab	770.00 ab	2.20 b
T ₇	12 a	803.60 a	2.30 a
T ₈	12 a	798.00 a	2.28 a

Table 3. Effects of Zn as foliar and soil application on Pulp fresh weight, Stone weight and Yield per tree of mango.

Treatment	Pulp fresh weight	Stone weight	Yield per tree
	(g)	(g)	(kg)
T ₀	214.29 g	150.00 f	150.00 e
T ₁	228.57 f	160.00 e	160.00 de
T ₂	237.62 e	166.33 de	166.00 cd
T ₃	243.81 de	170.67 cd	170.00 cd
T ₄	250.00 cd	175.00 b-d	175.00 bc
T ₅	252.86 c	177.00 bc	177.00 bc
T ₆	261.90 b	183.33 ab	183.00 ab
T ₇	273.33 a	191.33 a	191.00 a
T ₈	271.43 a	190.00 a	190.00 a

Total sugars (TS), reducing sugars (RS) and non-reducing sugars (NRS) of mango: Treatments effect was significantly different for TS, NRS and RS of mango. The effect of treatments was significant for the improvement in TS, RS and NRS. It was observed that all the treatments were significant from control for the improvement in NRS. Treatments T₃, T₄, T₅, T₆, T₇ and T₈ differed significantly over control for the improvement in TS. However, T₁ and T₂ did not differ over control for TS. In the case of RS, treatments T₂, T₃, T₄, T₅, T₆, T₇ and T₈ differed significantly over control for the improvement in TS (Table 4). While only T₁ did not differ over control for RS. Chord diagram showed that the percentage share of treatment T₇ was highest than T₀ for total sugars, reducing sugars and non-reducing sugars of mango (Fig. 4). Pearson correlation showed that a significant positive correlation existed between treatments, TS, RS and NRS. The maximum increase in TS, RS and NRS were noted where T₇ was applied.




Fig. 1. Chord diagram showing the percentage share of each treatment for average fruit weight, fruit width and fruit length.




Fig. 2. Chord diagram showing the percentage share of each treatment for shelf life, number of flowers per panicle and number of fruits matured per panicle of mango.




Fig. 3. Chord diagram showing the percentage share of each treatment for pulp fresh weight, stone weight and yield per tree of mango.

Table 4. Effects of Zn as foliar and soil application on Total sugars, Reducing sugars and Non-Reducing sugars of mango.

Treatment	Total sugars	Reducing sugars	Non-reducing sugars
	(%)		
T ₀	18.00 d	14.85 e	7.43 g
T ₁	19.20 cd	15.84 de	7.92 f
T ₂	19.96 b-d	16.47 cd	8.23 e
T ₃	20.48 bc	16.90 cd	8.45 de
T ₄	21.00 a-c	17.33 bc	8.66 cd
T ₅	21.24 a-c	17.52 bc	8.76 c
T ₆	22.00 ab	18.15 ab	9.08 b
T ₇	22.96 a	18.94 a	9.47 a
T ₈	22.80 a	18.81 a	9.41 a

Table 5. Effects of Zn as foliar and soil application on Photosynthetic rate, Transpiration rate and Stomatal conductance of mango.

Treatment	Photosynthetic rate	Transpiration rate	Stomatal conductance
	($\mu\text{mol}/\text{m}^2/\text{s}$)	($\text{mmol}/\text{m}^2/\text{s}$)	($\text{mol}/\text{m}^2/\text{s}$)
T ₀	14.85 g	4.50 f	0.23 e
T ₁	15.84 f	4.80 e	0.24 d
T ₂	16.47 e	4.99 de	0.25 d
T ₃	16.90 de	5.12 cd	0.26 cd
T ₄	17.33 cd	5.25 b-d	0.26 bc
T ₅	17.52 c	5.31 bc	0.27 bc
T ₆	18.15 b	5.50 ab	0.28 ab
T ₇	18.94 a	5.74 a	0.29 a
T ₈	18.81 a	5.70 a	0.29 a

Photosynthetic rate, Transpiration rate and Stomatal conductance of mango: Effect of treatments was significant for photosynthetic rate (Pn), transpiration rate (E) and stomatal conductance (SC) of mango. The influence of treatments was significant for the improvement in TS, RS and NRS. It was observed that all the treatments were significant from control for the improvement in Pn, E and SC (Table 5). Chord diagram showed that the percentage share of treatment T7 was highest than T0 for photosynthetic rate, transpiration rate and stomatal conductance of mango (Fig. 5). Pearson correlation showed a significant positive correlation between treatments, Pn, E and SC (Fig. 6). The maximum increase in Pn, E and SC were noted where T7 was applied.

Discussion

In the current experiment, the application of T7 and T8 maximized the growth and yield attributes of mango. The improvement in fruit width and length, average fruit weight, stone weight, TSS, and yield per tree might be due to better regulation of gas exchange attributes and nutrient uptake. These results can be correlated with the findings of Ahmad *et al.*, (2018). Zn-finger transcription factors regulate the physiological processes that developed the floral parts, i.e., anthers, pollen, pistil and secretory tissues in many plant species (Kobayashi *et al.*, 1998). It has been observed that deficiency of Zn disrupts the development of flowers and fruits (Epstein & Bloom, 2005). Rodríguez *et al.*, (2005) added Zn + KCl in 'Valencia' orange trees and observed that K and Zn foliar concentrations positively affect the production of medium and large-sized fruits. These results are in




Fig. 4. Chord diagram showing the percentage share of each treatment for total sugars, reducing sugars and non-reducing sugars of mango.




Fig. 5. Chord diagram showing the percentage share of each treatment for photosynthetic rate, transpiration rate and stomatal conductance of mango.

agreement with the work of Bariya *et al.*, (2014), and Ahmad *et al.*, (2018). They found that Zn improved the growth of mango by activating the enzymatic reactions. Zinc is a co-factor of more than 300 enzymes and proteins and has a quick and precise effect on cell division, protein synthesis and nucleic acid metabolism (Marschner, 1986). Zinc is an important micronutrient for plants as it is involved in numerous enzymatic reactions and is necessary for good growth and development. Zinc is also involved in regulating carbohydrate and protein metabolism (Swietlik, 1999). Furthermore, Zn uptake in mango plants will be more if ZnSO₄ is applied as foliar application than soil application (Bahadur, 1998). The positive correlation of its foliar application in enhancing the mango productivity was cited by Singh & Maurya (2004); Ranjit *et al.*, (2008) and elevating the mango fruit quality in terms of total sugar and total soluble solids

(TSS) (Rashmi & Singh, 2007). However, the balanced use of Zn and the best application method needs more attention to synchronize the application rate of Zn with inorganic fertilizers. Hasani *et al.*, (2016) also found similar results of enhancement in yield via Zn over control. The enhancement of mango fruit yield, vitamins, sugar concentrations, and some physiological features and improvement in quality was due to the use of Zn. In the current experiment, improvement in total sugar, reducing sugar and non-reducing sugar also validated the efficacious functioning of Zn. The application of Zn regulated the rate of photosynthesis in plants. Improvement in photosynthesis increase the synthesis of carbohydrates, i.e., sugar contents. According to Hegde & Venkatesh, (2007) Zn positively affects the biochemical activities of plants by establishing or stimulating enzymes, cells osmoregulation and altering membrane permeability. Zinc also played an imperative role in better mineral

nutrition uptake by catalyzing the macronutrient uptake (Phillips, 2004). A rise in the TSS and SC of fruits via foliar Zn has been documented in pomegranate and mango cv. Dusehri (Khorsandi *et al.*, 2009; Anees *et al.*, 2011). The improvement in fruit quality might be via catalytic activation caused by micronutrients mainly at high concentrations. Hence, the foliar application of micronutrients quickly increased the uptake of macronutrients in the tissues and organs of the mango plants, decreased the nutritional deficiencies and improved the fruit quality. The increase of mango fruit quality and yield in response to the combined application of Zn may be due to improvements in sugar concentrations, vitamins and some physiological features Hegde & Venkatesh, (2007). Anees *et al.*, (2011) detected improved fruit ascorbic acid and sugar via ZnSO₄ addition. They also documented enhancement in total sugars of fruit by increasing zinc rates.




Fig. 6. Pearson correlation for treatments and different attributes. * = 0.05, ** = 0.01, *** = 0.001

Conclusion

It is concluded that applying both soil and foliar application of ZnSO₄ has the potential to improve the growth and yield attributes of mango. Treatments T₇ (RD + ZnSO₄ + 150 g plant⁻¹ as basal and 1.0% as foliar spray) followed by T₈ (RD + ZnSO₄ + 300 g plant⁻¹ as basal and 1.0% as foliar spray) should be applied to maximize the fruit length, fruit width, average fruit weight, stone weight, total soluble solid and yield per tree of mango. In addition, T₇ and T₈ were also capable of modifying the gas exchange attributes, i.e., photosynthetic rate, stomatal conductance and transpiration rate of mango. It is recommended to use T₇ for maximizing the growth, yield and quality of mango, keeping in mind the efficacious use of fertilizer.

References

Ahmad, I., F. Bibi, H. Ullah and T. Munir. 2018. Mango fruit yield and critical quality parameters respond to foliar and soil applications of zinc and boron. *Plants*, 7(4): 97.

Ahmed, N., S. Khalid, A.G. Grewal, M.A. Ali, M.A. Anjum, A.A. Rahi and S. Danish. 2020. Performance of mango scion cultivars under various levels of artificially induced salinity stress. *Pak. J. Bot.*, 52. 10.30848/PJB2020-4(11).

Akhtar, M., S. Yousaf, N. Sarwar and S. Hussain. 2019. Zinc biofortification of cereals-role of phosphorus and other impediments in alkaline calcareous soils. *Environ. Geochem. Hlth.*, 41: 2365-2379.

Anees, M., F.M. Tahir, J. Shahzad and N. Mahmood. 2011. Effect of foliar application of micronutrients on the quality of mango (*Mangifera indica* L.) cv. Dusehri'fruit. *Mycopathologia*, 9: 25-28.

- Anonymous. 2010. Production Year Book. Food and Agriculture Organization of the United Nation.
- Bahadur, L., C.S. Malhi and Z. Singh. 1998. Effect of foliar and soil applications of zinc sulphate on zinc uptake, tree size, yield, and fruit quality of mango. *J. Plant Nutr.*, 21: 589-600.
- Baloch, F.S., S.S. Tahir, S.T.H. Sherazi, N.S. Jilani, A.L. Khokhar and M.T. Rajput. 2017. Evaluation of essential oil components from the fruit peelings of sindhri and langra varieties of mango (*Mangifera indica* L.). *Pak. J. Bot.*, 49(4): 1479-1484.
- Bariya, H. S. Bagtharia and A. Patel. 2014. Boron: A promising nutrient for increasing growth and yield of plants. In: *Nutrient Use Efficiency in Plants*; Kopriva, S., Ed.; Springer: Basel, Switzerland.
- Bose, T.K. 1999. Fruits, history and products: Tropical Horticulture Dept. of Horti. Bidhan Chandra *Krishi Viswavidyalaya*, Naya Udyog Publishers, Calcutta 32: 179-198.
- Brown, P.H., I. Cakmak and Q. Zhang. 1993. Form and function of zinc in plants. In: (Ed.): Robson, A.D. Zinc in soils and plants (pp. 90-106). Dordrecht: Kluwer Academic Publishers. Evaluate antioxidant activity. *LWT – Food Sci. Technol.*, 28: 25-30.
- Cakmak, I. 2000. Role of zinc in protecting plant cells from reactive oxygen species. *New Phytol.*, 146: 185-205.
- Epstein, E. and A.J. Bloom. 2005. Mineral nutrition of plants: principles and perspectives. Sinauer Associates, Sunderland.
- Fageria, N.K. 2009. The use of nutrients in crop plants. Boca Raton, FL: CRC Press.
- Hasani, M., Z. Zamani, G. Savaghebi and R. Fatahi. 2016. Effects of zinc and manganese as a foliar spray on pomegranate yield, fruit quality and leaf minerals. *J. Soil Sci. Plant Nutr.*, 12: 471-480.
- Hegde, V.L. and Y.P. Venkatesh. 2017. Anaphylaxis following ingestion of mango fruit. *J. Invest. Allergy*, 17: 341-344.
- Hortwitz, W. 1960. Official and tentative methods of analysis. Ed. 9. Assoc. *Off. Anal. Chem.*, Washington, D.C., USA.
- Jan, A.U., A. Shah and F. Hadi. 2019. Role of potassium, zinc and gibberellic acid in increasing drought stress tolerance in sunflower (*Helianthus annuus* L.). *Pak. J. Bot.*, 51, 809-815. DOI: 10.30848/PJB2019-3(4).
- Kazmi, M.R., F.S. Fateh, K. Majeed, A.M. Kashkhely, I. Hussain, I. Ahmad and A. Jabeen. 2005. Incidence and etiology of mango sudden death phenomenon in Pakistan. *Pak. J. Phytopathol.*, 17: 154-158.
- Khorsandi, F., F.A. Yazdi and M.R. Vazifehshenas. 2009. Foliar zinc fertilization improves marketable fruit yield and quality attributes of pomegranate. *Int. J. Agri. Biol.*, 11: 766-770.
- Kobayashi, A., A. Sakamoto, K. Kubo, Z. Rybka, Y. Kanno and H. Takatsuji. 1998. Seven zinc-finger transcription factors are expressed sequentially during the development of anthers in petunia. *Plant J.*, 13: 571-576.
- Madonsela, E.N. 2019. Defining and describing fruit quality of mango, its development and protection in fruit production. *Acta Sci. Agric.*, 3(9): 38-41, DOI: 10.31080/ASAG.2019.03.0603.
- Marschner, H. 1986. The Mineral Nutrition of Higher Plants. 1st Ed., Academic Press, New York, USA.
- Memon, N.A. 2016. "Mango: Pakistan 4th Largest Producer in the World." <http://foodjournal.pk/2016/March-April-2016/PDF-March-April-2016/Exclusive-article-Mango.pdf>.
- Phillips, M. 2004. Economic benefits from using micronutrients for the farmer and the fertilizer producer. IFA, International symposium on micronutrients. ND, India.
- Rajendran, C. and K. Ramamurthy. 2009. Nutritional and physiological disorders in crop plants, scientific publishers, Jodhpur; pp. 6-10.
- Ranganna, S. 1986. Handbook of analysis and quality control for fruit and vegetable products. Tata McGraw-Hill Publishing Company, New Delhi, India
- Ranjit, K., K. Pawan and U.P. Singh. 2008. Effect of foliar application of nitrogen, Zinc, Boron on flowering and fruiting of mango (*Mangifera indica* L.) c.v amepali. *Environ. Ecol.*, 26: 1965-1967.
- Rashmi, P. and C.P. Singh. 2007. Effect of pre-harvest foliar spray of micronutrients on chemical properties of mango fruit cv. Langra. *Pant. J. Res.*, 5: 56-61.
- Rodriguez, V.A., S.M. Mazza, G.C., Martanez and A.R. Ferrero. 2005. Zn and K influence in fruit sizes of Valencia orange. *Rev. Brasil. de Fruticul.*, 27: 132-135.
- Saran, L.S. and R. Kumar. 2011. Boron deficiency disorders in mango (*Mangifera indica* L.): field screening, nutrient composition and amelioration by boron application. *Int. J. Agric. Sci.*, 81: 143-149.
- Sharma, R.R. 2006. Fruit production: Problems and Solutions. International Book Distributing Co, Lucknow, India. 342-345.
- Sharma, R.R. and R. Singh. 2006. Pruning intensity modifies canopy microclimate, and influences sex ratio, malformation incidence and development of fruited panicles in 'Amrapali' mango (*Mangifera indica* L.). *Sci. Hort.*, 109: 118-122.
- Siddiq, M., J.K. Brecht and J.S. Sidhu. 2017. Handbook of mango fruit: production, postharvest science, processing technology and nutrition. Oxford, UK: Wiley.
- Sing, Z., A.U. Malik and T. Davenport. 2005. Fruit drop in mango. *Hort. Rev.*, 31: 111-140.
- Singh, G. and A.N. Maurya. 2004. Effect of micronutrients on bearing of mango cv. Malika. *Prog. Agric.*, 4: 47-50.
- Swietlik, D. 1999. Zinc nutrition in horticultural crops. Horticultural Reviews. *John Wiley & Sons, Inc.* New York. pp. 109-180.
- Usman, M., B. Fatima, M.M. Khan and M.I. Chaudhry. 2003. Mango in Pakistan: a chronological review. *Pak. J. Agri. Sci.*, 40: 151-154.

(Received for publication 21 September 2020)