

SPATIO-TEMPORAL ANALYSIS OF CUCUMBER MOSAIC VIRUS DISEASE AND ITS SUSTAINABLE MANAGEMENT

MUHAMMAD AHMAD ZESHAN¹, YASIR IFTIKHAR^{1*}, ASHARA SAJID¹, MUHAMMAD USMAN GHANI², MUHAMMAD ZAFAR-UL-HYE³ AND MAZHAR ABBAS⁴

¹Department of Plant Pathology, College of Agriculture, University of Sargodha, Pakistan.40100.

²Institute of Soil and Environmental Sciences, University of Agriculture Faisalabad, Pakistan. 38000.

³Department of Soil Science, Bahauddin Zakariya University Multan 60800, Pakistan

⁴Department of Agriculture and Agribusiness Management, Faculty of Science, University of Karachi, Pakistan

*Corresponding author's email: yasir.iftikhar@uos.edu.pk

Abstract

Cucumber fields were surveyed in the vicinity of Sargodha to map the spatial and temporal CMV disease distribution. Infected samples were collected for pathogenicity tests, biophysical properties and serological assays. After confirmation of virus; data of disease prevalence, incidence and severity was recorded which showed strong positive linear relationship. There was gradual increase in disease after each temporal assessment. Bhatti Town (BT) showed more than 90% disease prevalence followed by College of Agriculture (COA) and Adaptive Research Farm (ARF). This experiment also demonstrated the significant contribution of NPK solution, Naphthalene acetic acid (NAA), benzothiadiazole (BTH) and elephant ear plant (EEP) extract in improving the cucumber performance against CMV disease. There was positive linear relationship between all disease parameters. All the treatments gave significant contribution for sustainable disease management. NPK was the efficient among rest of the treatments with more than 75% disease inhibition. The second most effective treatment was BTH followed by NAA and EEP with 63%, 55% and 39% disease control, respectively. The study provided useful information for estimation disease dynamics in varied spatio-temporal scales. It could be concluded that NPK and BTH are the suitable tools for sustainable management of CMV disease.

Key words: Naphthalene acetic acid, Micronutrients, Epidemiology, Reducing agents, Inoculums.

Introduction

Cucumber mosaic disease (CMD) is caused by cucumber mosaic virus (CMV) that was first reported in 1916 in New York (Yoon *et al.*, 2019). CMV belongs to familyBromoviridae and genusCucumovirus that has ssRNA genome (Nalam *et al.*, 2019).The virusinterrupts with somatic and reproductive physiology of plants and cause extensive yield losses (Donnelly *et al.*, 2019).The highest percentage of CMD severity may result in 100% yield losses (Groen *et al.*, 2017).CMV has wider host range and can attack more than 1000 plant species (Wu *et al.*, 2017). It passes through the cell wall of host and finds the replication sites, removes coat protein, becomes a part of host and replicates through replicases enzyme (Mauck, 2016). It damages the biosynthetic machinery of host plants as it replicates in that system (Casteel *et al.*, 2015).

The salient symptoms of the disease are stunting, dark and light green patterns, malformation and yellow streaking of leaves (Tungadi *et al.*, 2017). It is transmitted through sap inoculation (Gao *et al.*, 2016), seeds (Arogundade *et al.*, 2019), aphids (Tungadi *et al.*, 2020) and 10 species of dodder (Ntui *et al.*, 2013). Aphids transmit CMV in a stylet borne manner with acquisition feeding period of 10 seconds (Jo *et al.*, 2016).

The activation of induced systemic resistance triggers the expression of plant defense genes (Kumar *et al.*, 2017). It can be strengthened by using different nutritional amendments (Meena *et al.*, 2016) and by the application of salicylic acid, jasmonates and ethylene (Qin *et al.*, 2013). Jasmonic acid and ethylene induce the systemic resistance in plants against the signaling mechanisms of pathogens (Aznar *et al.*, 2015).

Similarly, spraying of insecticides is also an option available to the farmers for managing CMV through vectors control. However, as CMV is an aphid-transmitted non-persistent virus, only insecticidal spray may not control the disease effectively. Moreover, dependence on a single method is highly vulnerable to failure (Borer *et al.*, 2016). However, when integrated with more than one management strategies may repress disease significantly more than any single tactic alone (Anitha, 2016). Therefore, if available an integrated approach is preferred. Many reports are available on the successful application of integrated management tactics for CMV (Al-Zahrani *et al.*, 2018).The aqueous extract of *Tanacetumvulgare* was applied for the management of CMV and many other mechanically transmitted plant viruses (Petrov *et al.*, 2016).There are essentially two approaches to manage virus diseases. The first approach is to decrease the sources of infection and secondly to minimize the rate of spread by vector control (Rahman *et al.*, 2020).

The potential of nutrients and plant extracts for improving the tolerance of cucumber plants against cucumber mosaic virus disease has not been documented so far. Therefore, this study, evaluated the potential of NPK solution, naphthalene acetic acid and elephant plant extract for improving salt tolerance in wheat.

Materials and Methods

Plant material and experimental layout: The experiment was conducted in research area Department of Plant Pathology, College of Agriculture, University of Sargodha, Pakistan 32.0754° N and 72.41168° E using completely randomized design (CRD). The prevailing environmental conditions during the experiment were; having average

temperature of 29°C with relative humidity 57.43%, 10.4 h sunshine, 15.2 mm rainfall, 14 km/h wind speed and 1.8 mm potential evapo-transpiration. Cucumber seeds of Poinsettee variety were obtained from local market, Sargodha. Clay pots of 40 cm diameter were filled with 25% sand, 75% clay and appropriate amount of farmyard manure. The 30 pots were seeded with cucumber seeds (1 cm deep) at the rate of 10 seeds per pot and kept under natural daylight in glasshouse. The pots were irrigated with tap water regularly till the end of experiment. The fertilizers i.e. potassium sulphate, urea, and diammonium phosphate were applied as sources of potassium, nitrogen and phosphorous respectively.

Sources of viruses and inoculums preparation:

Cucumber crop was surveyed at Adaptive Research Farm (ARF), Sargodha (32.1228° N, 72.6786° E) for the collection of diseased samples. The samples were collected in zip lock polythene bags, placed in ice-bucket and stored in the refrigerator at 4°C in Plant Pathology laboratory. Virus inoculums were prepared from infected fresh leaves of cucumber having alternate light green patches intermingled with normal green color. Infected cucumber leaves (5 g), were macerated in 5 ml of 0.01 M phosphate buffer in pre-chilled pestle and mortar, filtered through muslin cloth and stored in a falcon tube (Gao *et al.*, 2016).

Virus inoculation on cucumber and indicator plants:

The primary leaves of cucumber and indicator plants (chili, tomato and chinopodium) were dusted with carborandum

powder. A cotton swab was dipped in falcon tube containing inoculums and injured leaf surfaces were rubbed. The inoculated leaves were sprinkled with water to remove extraneous inoculums. The inoculated plants were covered with polythene sheet to avoid from the attack of insects and other pathogens (Crespo *et al.*, 2018).

Serological assay (DAS-ELISA) for detection of CMV:

Infected samples were collected and double antibody sandwich (DAS-ELISA) was performed in Laboratory of Plant Virology section Ayub Agricultural Research Institute Faisalabad-Pakistan. Two plants from each location were selected for serological assay and sap was extracted from leaf lamina and leaf mid rib. The standard procedure of ELISA as described by Clark & Adams (1977) was followed and after stopping the reaction, micro-titre plate was kept in ELISA reader for recording the absorbance readings.

Spatio-temporal assessments of CMV disease severity:

After confirmation of CMV through mechanical inoculation; the cucumber fields at ARF, COA and Bhatti Town were visited weekly for noting disease observations. The D_{50} and D_{95} were calculated for each location by determining the nearest neighbor distance from each CMV block at a specific time. The nearest neighbor distance was calculated by using ArcMap software (Gougherty & Nutter, 2012). The data of disease incidence and severity were recorded four times from three places at weekly intervals by using following formulae.

$$\text{Disease incidence (\%)} = \text{Number of diseased } \frac{\text{plants}}{\text{Total}} \text{ number of plants} \times 100$$

$$\text{Disease severity (\%)} = \text{Number of diseased } \frac{\text{leaves}}{\text{Total}} \text{ number of leaves} \times 100$$

$$\text{Disease prevalence (\%)} = \text{Number of symptomatic } \frac{\text{plots}}{\text{Total}} \text{ number of plots} \times 100$$

Area under disease progressive curve (AUDPC) was calculated by the trapezoidal integration $n - 1$
 $AUDPC = \sum [(x_i + x_{i+1})/2](t_{i+1} - t_i)$
 $i = 1$

where n is the number of assessment; x, disease incidence (%); and $(t_{i+1} - t_i)$, duration between two consecutive assessments (Shaner and Finney, 1977).

$$\text{Disease severity index (\%)} = \sum P \times Q / (M \times N) \times 100$$

where P = severity score, Q = number of infected plants having the same score; M = Total number of plants observed, N = Maximum rating scale number

Disease was categorized according to following disease rating scale as mentioned in table 1 (Bashir *et al.*, 2002).

Effect of sap storage on virus infectivity: Sap was prepared with standard method and stored under optimum conditions. Then sap was inoculated with the standard method as described above. Four groups of plants containing 5 pots each were inoculated with varied time interval i.e. same day the inoculums were prepared, 2 days after preparation, 4 days after preparation and 6 days after preparation (Mujtaba *et al.*, 2019). The symptoms produced by each set of inoculums were assessed on the inoculated plants.

Table 1. Disease rating scale for CMV

Disease rating	Disease severity (%)	Response
0	All plants free of virus symptoms	HR (Highly Resistant)
1	1-10% infection	R(Resistant)
2	11-20% infection	MR (Moderately Resistant)
3	21-30% infection	MS(Moderately Susceptible)
4	30-50% infection	S (Susceptible)
5	More than 50% infection	HS (Highly Susceptible)

Effect of reducing agents on crude (CMV) sap: To evaluate the efficacy of reducing agents on crude CMV sap, 5ml of 2.5% carbon tetrachloride (CCl₄) and 5ml of 5% chloroform (CHCl₃) were added in the crude CMV sap (Ali *et al.*, 2020). The sap containing reducing agents was applied on two sets of plants (5 pots each) including one control. After inoculation, plants were covered and symptoms were observed to assess the efficacy of reducing agents on CMV virulence.

Demonstration of dilution end point (DEP): Sap was prepared with the standard method. 1ml of sap was taken and added into 9ml of distilled water in the test tube (10⁻¹ dilution). 1ml of sap was taken from 10⁻¹ dilution and added into 9ml of distilled water in the test tube (10⁻² dilution). Similarly, 1ml was taken from 10⁻² dilution and added into 9ml of distilled water in the test tube (10⁻³ dilution). These dilutions were inoculated with the different dates (Sharma *et al.*, 2020). All sets of diluted sap were inoculated on three sets of plants including positive and negative control. In positive control, undiluted sap was applied while negative control remained untreated.

Effect of different treatments against CMV disease under field conditions: Elephant ear plant (*Clocasia esculenta* L.) extract was used against CMV disease at three different concentrations. The leaves of test plants were collected from premises of COA, UOS and extract was prepared by grinding the leaves with manual plant extraction machine. Equal v/v of sample and water used to make 100% solution. The relative concentrations of 5%, 7.5% and 10% were made by adding 5 ml, 7.5 ml and 10 ml of extract in 95 ml, 92.5 ml and 90 ml of distilled water, respectively. The extract obtained was considered as standard and was stored at -20°C until use. Three foliar sprays of plant extracts at 5%, 7.5% and 10% concentrations were applied at an interval of 7 days. After each spray disease incidence data was recorded two times at five days' interval.

Ali Akbar product named "Fashion" was taken as NPK and was used at three concentrations i.e. 2.5%, 5% and 7%. This was prepared by adding 25ml, 50ml and 75ml NPK solution into 1 liter of water, then apply as a foliar spray. STEDEC product name Phytifix was taken as a naphthalene acetic acid (NAA). The concentrations (1.5%, 2.5% and 3.5%) were prepared by adding 15ml, 25ml and 35ml into 1 liter of water. Benzothiadiazole (BTH) was applied at three concentrations (0.2%, 0.3% and 0.5%) prepared by adding 2g, 3g and 5g in one liter of distilled water.

The data of disease severity was recorded before and after each spray by following above mentioned formula. The efficacy of each treatment was calculated by using the formula given below described by (Helal, 2019).

$$I\% = C - T / T * 100$$

where I% = Inhibition or efficacy; C = Control and T = Treated

Comparison of diseased and healthy leaf to check leaf area: Ten leaves each from diseased and healthy plants were taken to check the leaf area cm² with the leaf area meter (Plate 1).

Dry preservation of inoculums: In order to perform the virus indexing procedure conveniently, diseased samples were collected, cut the veins, chopped into small pieces and put into desiccators for 4-6 days. Took the dried inoculums into a bottle having four layers (Silica, muslin cloth, chopped leaves, and muslin cloth) for preservation (Plate 1).

Comparison of chlorophyll contents from diseased and healthy plants: Ten leaves from diseased and healthy plants were taken to check the chlorophyll contents with the chlorophyll content meter (Plate 1).

Statistical analysis

Data were analyzed statistically and means were compared by using Least Significant Difference (LSD) test and standard error was calculated. ANOVA was used to determine the effect of treatments on disease severity (Steel *et al.*, 1977).

Results and Discussions

Spatial distribution of CMV disease in different locations: Significant differences were observed for prevalence, incidence and severity of CMV at three locations of ARF, COA and BT in Sargodha district. The parameters of disease estimation showed higher values in all locations of BT and COA, respectively while minimum observed in ARF. Maximum disease prevalence (100%) was recorded in BT at location 1, while location 1 and 2 of ARF gave minimum disease prevalence (60%). Disease prevalence ranged from 60-70%, incidence ranged from 51%-59% and severity from 32-39% in ARF. The overall disease prevalence (60-100%), disease incidence (51-75%), and disease severity (32-55%) was recorded in all locations of three places (Table 2). The locations with more CMV disease prevalence gave more disease incidence and severity while low values of disease incidence and severity were recorded in less prevalent locations respectively. In all three locations, mean disease prevalence depicted strong linear relationship with disease incidence and severity (Figs. 1 & 2). For each per unit increase in disease prevalence there was relative increase in disease incidence (0.21-0.56) and severity (0.34-0.52). The values of D₉₅ and D₅₀ were calculated for all locations collectively which conformed the disease distribution pattern to real model. The values of (D₉₅ and D₅₀) for prevalence are higher than incidence and severity indicating the preciseness of disease distribution. These values indicate the conservative measurement of pathogen spread. Gougherty and Nutter proposed that these values provide a guide for the establishment of a quarantine area for careful future survey. Furthermore, these are also helpful in devising disease eradication programs for an area. Pearson's correlation indicated more than 95% strong relation among disease prevalence and incidence and severity.

Cucumber mosaic virus ranked among the most damaging disease causing agents worldwide. The remarkable yield losses incited by CMV disease may be eliminated by sowing resistant germplasm against the virus. There are very limited studies to explore the natural genetic potential of cucumber cultivars against CMV (Iqbal *et al.*, 2012). The present research was undertaken to unveil the hot spots for CMV disease in three different locations of Sargodha that would be helpful in deciding the sustainable management options against quality and yield destruction. All the designated locations were

regarded as susceptible and highly susceptible when rated according to a devised scale. The non-availability of resistant sources is a key factor in minimizing the cost-benefit ratio. CMV is the most damaging viral pathogen with wide host range which infects many crops from various families (Ashfaq *et al.*, 2014). Due to multiple hosts and high mutation rate, provision of genetically resistant germplasm against CMV is limited. That's why conventional methods including cultural control, physical eradication, chemical and bio-control of vectors are mostly relied upon for CMV management (Hull, 2014)

Table 2. Spatial distribution of CMV disease in various cucumber fields of district Sargodha.

Spatial distribution with GPS location	Locations	DP (%)	DI (%)	DS (%)
ARF 32.1228°N 72.6786°E	1	60 ± 3.16 b	51.57 ± 1.32 c	32.16 ± 1.24 c
	2	60 ± 1.32 b	57.08 ± 1.41 b	35.53 ± 1.73 b
	3	70 ± 2.41 a	59.06 ± 1.73 a	39.09 ± 1.31 a
COA 32.0754°N 72.41168°E	1	80 ± 1.41 a	67.02 ± 1.73 b	44.18 ± 1.59 b
	2	70 ± 2.24 b	62.51 ± 2.24 c	41.97 ± 1.73 c
	3	80 ± 2.83 a	69.15 ± 2.65 a	47.42 ± 1.41 a
BT 32.1087°N, 72.6543°E	1	100 ± 0.0 a	75.17 ± 2.24 a	55.14 ± 1.89 a
	2	90 ± 1.73 b	73.38 ± 1.89 b	52.87 ± 1.63 b
	3	90 ± 2.47 b	72.78 ± 1.63 ab	50.62 ± 1.62 c
	D ₉₅	96.01	74.45	54.23
	D ₅₀	80.24	67.02	44.18

Means with similar letters in columns against each location are not significantly different 5% probability level

To assess the temporal distribution of CMV disease, the cucumber crop was observed. CMV disease was evaluated by recording prevalence, incidence and severity from all locations at fortnight interval for three times. The overall minimum values of disease parameters were found after first assessment in all locations while all parameters increased after second and third assessments. Significant differences were recorded for disease prevalence, incidence and severity in all assessments (Table 3). During first assessment disease prevalence was minimum at all locations which ranged from 63.55%-93.53% that gradually increased after second assessment (66.31-95.67%) and third assessment (69%-97%). It was assessed that disease incidence and severity also increased with each consecutive assessment. The range of temporal increase in disease incidence and disease severity was 55.91%-78.02% and 35.59%-58.19%, respectively.

The evaluation of cucumber crop at all locations indicated that there was no resistance against CMV disease (Table 3). The disease severity index and AUDPC value showed increasing trend at each assessment. The crop surveyed at ARF and COA was regarded as susceptible when compared with disease rating scale while BT gave highly susceptible response. The disease severity index ranged from 0.71-1.39% with disease rating of 4 and 5. The value of AUDPC was lower in temporal assessment at ARF which ranged from 157.98 to 1309.28 which increased in case of COA (1491.08-1572.07) and BT (1660.05-1755.45).

There was highly significant difference in each temporal assessment of CMV disease distribution which showed increasing trend. These results are in line with the findings of a cucumber screening experiment which was conducted for evaluation of tomato germplasm against

CMV (Akhtar *et al.*, 2010). The spatial and temporal disease dynamics would provide a strong base for eradication plans in the future (Gougherty *et al.*, 2015). Spatio-temporal disease distribution maps are very intuitive in plant disease management strategies (Byamukama *et al.*, 2010). The spatial analysis for disease measurement was strengthened by devising a new method to differentiate by two consecutive spaces through the calculation of D₉₅ and D₅₀ values (Gougherty and Nutter, 2012). A more advance method in spatial analysis was to determine the hot spot among locations by applying Ripley's K function which specifies the site needs more attention (Nutter *et al.*, 2011). While studying spatial disease distribution, each unit should be sampled systemically due to clustered pattern (Byamukama *et al.*, 2011).

Effect of sap storage on virus infectivity (Longivity in vitro-LIV): The studies on LIV done at room temperature which showed 100% transmission and maximum lesions when inoculated with crude sap immediately after extraction from infected leaves. Gradual decrease in transmission was observed when the crude sap was kept at room temperature for varied duration. The results showed that the virus could retain infectivity up to 6days and got inactivated from 6th day when kept at room temperature (Fig. 3).

The effect of aging LIV has been studied by many researchers worldwide. The attenuation of CMV sap infectivity with increased storage was recorded by Chandankaret *et al.*, (2013). In another study, maximum transmission in main host and indicator plants were observed by inoculation with fresh sap that decreased drastically when inoculation was done 6 days old sap (Mujtaba *et al.*, 2019).




Fig. 1. Relationship between disease prevalence and incidence at three locations.




Fig. 2. Relationship between disease prevalence and severity at three locations temporal distribution of CMV disease.

Table 3. Temporal distribution of CMV disease at fortnight intervals.

Locations	Data assessments	DP (%) ^	DI (%)	DS (%)	DSI (%) ^	Disease rating	AUDPC^	Response
ARF	First	63.55 ± 1.44i	55.91 ± 1.41i	35.59 ± 2.42i	0.71	4	1257.98	S^
	Second	66.31 ± 1.23 h	58.03 ± 2.65h	37.13 ± 2.13 h	0.74	4	1305.68	S
	Third	69.53 ± 2.82 g	58.19 ± 1.82 g	38.98 ± 1.93 g	0.77	4	1309.28	S
COA	First	73.53 ± 1.73 f	66.27 ± 1.57 f	42.51 ± 2.01 f	0.85	4	1491.08	S
	Second	75.14 ± 1.37 e	68.05 ± 1.26 e	44.73 ± 1.06 e	0.89	4	1531.13	S
	Third	79.29 ± 1.31 d	69.87 ± 1.21 d	48.22 ± 2.11 d	0.96	4	1572.07	S
BT	First	93.53 ± 1.01 c	73.78 ± 1.18 c	52.88 ± 1.78 c	1.27	5	1660.05	HS
	Second	95.67 ± 1.22 b	75.87 ± 1.34 b	55.06 ± 1.89 b	1.32	5	1707.08	HS
	Third	97.19 ± 1.38 a	78.02 ± 1.63 a	58.19 ± 1.71 a	1.39	5	1755.45	HS

^value given against each assessment in all columns is the mean of three readings

DP= Disease prevalence; DS= Disease severity; DI= Disease incidence

^DSI= Disease severity index; AUDPC= Area under disease progress curve; S= Susceptible; HS= Highly susceptible

Effect of reducing agents on crude (CMV) sap: The virulence of CMV was increased when its sap was mixed with 5% carbon tetrachloride (CCl₄) and 2.5% chloroform (CHCl₃). These reducing agents showed significant higher number of lesions and percent transmission of CMV as compared with control. In consecutive inoculations, maximum infection (65%) was recorded in CCl₄ treated sap followed by CHCl₃ (53%) as compared to control (Fig. 4).

Plants have constitutive antiviral substances which may interrupt with virus replication, in order to overcome these substances reducing agents were mixed with crude virus sap. Chloroform is a strong reducing agent which gave significant higher transmission as compared to crude virus sap (Sapno *et al.*, 2017). Ali *et al.*, (2020) also used chloroform and carbon tetra chloride as reducing in cucumber mosaic virus sap.

Demonstration of dilution end point (DEP): The undiluted crude sap produced maximum number of lesions with 100% transmission. However, the sap retained its infectivity up to the dilution of 10⁻³. From this experiment, it was concluded that the dilution end point of the virus is 10⁻³ (Fig. 5). Assay plants inoculated with the crude sap diluted (10⁻¹-10⁻³) produced mosaic, leaf distortion and leaf puckering symptoms. However, sap failed to produce symptoms with extract diluted to 10⁻⁴ and above. These results corroborate with those of Parvin *et al.*, (2007) who described decreasing trend of virus transmission with every dilution. The infectivity loss of virus was noted with diluted sap inoculation on indicator plants (Mochizuki & Ohki, 2012).

Evaluation of different treatments against CMV disease: The mean values of disease severity showed that all the treatments were highly significant as compared to control with 71.95% disease severity (Table 4). Among all the treatments, the minimum mean disease severity (17.59%) was recorded in NPK treated plants followed by benzothiadiazole (BTH) 63.68%, NAA (55.06%) and elephant plant extract (39.44%). There was a significant difference in disease severity values after each spray. The comparison disease severity after three sprays showed that the minimum disease severity (22.53%) was recorded after 3rd spray and the maximum disease severity (41.25%) was recorded after 1st spray. The interaction of all the treatments was also significant with three sprays. The efficacy of all

the treatments was also calculated which indicated that NPK was the most efficient treatment with 75.55% reduction in CMV disease severity. The second most efficient treatment was BTH followed by NAA and elephant ear plant extract which showed 63.68%, 55.06% and 39.44% efficacy respectively.

The CMV disease severity was recorded for all the treatments which were applied at different concentrations (Table 5). The analysis showed that there was significant disease severity reduction after each concentration. In C₁, all treatments were applied at lower concentration which gave minimum disease reduction that was gradually increased in case of C₂ and C₃ where higher disease reduction values were noted. Disease severity of CMV under all the treatments was significantly lower than control. The interaction of concentrations with treatments was highly significant which gave maximum disease reduction in C₃ and minimum in C₁. The values of z-score inferred that each treatment was random in its respective concentration and none was clustered at ($p \leq 0.05$). The z-score values indicated that disease severity was random in concentration 1, 2 and 3 in all the treatments.

It's very interesting as none of the above mentioned treatments have insecticidal and viricidal properties. Then how these managed to combat the CMV disease because CMV impairs the plant physiology and these treatments have their independent pathways to repair the damages and triggering the plant defense. The infection caused by CMV disturbs the level of carbon assimilation and metabolic activities in infected plants. Previously in a study, biochemical characterization of infected leaf samples determined reduced carbohydrate concentration and photosynthesis with raised respiration rate (Shalitin and Wolf, 2020). There was significant reduction in biochemical reactions, photosynthetic pigments, protein contents, total phenols and alkaloids of CMV infected plants (Al-Zahrani *et al.*, 2018). The functioning of 2b counter defense proteins is impaired due to CMV attack which causes host volatile emissions (Tungadi *et al.*, 2017). Effect of CMV infection on host physiology was studied to investigate the alteration of cellular components like chlorophyll, stomatal conductance and β -carotene (Rahman *et al.*, 2019). There was considerable destruction in plant physiology by disruption in signals of hormones, cell deformities and carbohydrate movement (Mauck *et al.*, 2019).




Fig. 3. Effect of storage time on transmission of CMV.




Fig. 4. Graphical representation regarding effect of reducing agents on CMV transmission.




Fig. 5. Effect of series sap dilutions on transmission of CMV (crude sap= no dilution; 1= 10⁻¹; 2= 10⁻²; 3= 10⁻³; 4= 10⁻⁴). Values with different letters are significantly different from each other at 5% probability level.

The most efficient treatment in managing CMV disease was the foliar application of NPK followed by BTH, NAA and elephant ear plant extract. The plant defense mechanism is activated through the induction of pathogenesis-related (PR) proteins, phytoalexins accumulation and callose depositions (Saldajeno and Hyakumachi, 2011). Nutrients enhance growth, development and initiate defense against pathogenic and physiological stresses (Meddad-Hamza *et al.*, 2010). Nitrogen (N) acts as driving force in growth, development and yield of plants by nitrogen assimilation and amino acid transmission (Mur *et al.*, 2016). It provides nitric oxide defense signal activation against pathogen invasion (Liu *et al.*, 2015). Previous studies focused on plant defense as a function of nitrogen absorption through foliar application (Hout *et al.*, 2014). Nitrogen is a constituent of polyamines that follows systemic pathway against biotic stresses through oxidative phosphorylation (Tiburcio *et al.*, 2014). Nitric oxide signals stimulated by nitrogen metabolism initiate the salicylic acid dependent gene expression (Yu *et al.*, 2014). Nutrient metabolism enhances sugar contents, amino acids and GABA (γ -aminobutyric acid) in the challenged cells (Gupta *et al.*, 2013). Nitrogen yields metabolism yields products that take product in detoxification of reactive oxygen species that is linked with disease resistance (Chern *et al.*, 2013). NO works as a driving force in generating the hypersensitive response against biotic stresses (Vitor *et al.*, 2013). Potassium (K) regulates synthesis of protein, enzyme activation, carbohydrate metabolism, stomatal conductance and photosynthetic processes (Hasanuzzaman *et al.*, 2018). K interacts with plant signals which regulates biochemical processes and metabolism that affects plant growth, development and enhances stress tolerance (Shani *et al.*, 2017). It has vital role in photosynthesis, carbohydrate translocation and metabolism which ultimately enhance crop yield and quality (Lu *et al.*, 2016). Enhanced photosynthetic activity was recorded in phosphorus treated plants (Naem *et al.*, 2010). The biochemical analysis of phosphorus treated plants indicated enhanced growth, development, metabolic reactions and defense signals (Hakeem *et al.*, 2012). Similar results regarding increased metabolic activities were also observed in another study (Lambers & Shane, 2007).

The second most significant reduction was recorded in BTH applied plants. BTH is an appropriate treatment due to minute toxic hazards, quick degradation eco-friendly nature (Cao *et al.*, 2011). BTH triggers SAR pathway against viruses in various plant species in different plant species (Nischwitz *et al.*, 2008). It suppresses viral RNA accumulation in tobacco plants (Mandal *et al.*, 2008). BTH interfered with RNA replication resulting in reduced number of local lesion in BTH treated plants (Namitharan *et al.*, 2014). The antiviral activity of BTH derived chiral α -amino phosphonate studied by Zhang (Zhang *et al.*, 2017). In virus infected plants peroxidases (POD), polyphenol oxidases (PPO) and phenylalanine ammonia lyase (PAL) enzymes were significantly reduced (Helal, 2019). The efficiency of naphthalene acetic acid (NAA) in reducing CMV disease severity could be explained by the fact that increases protein, carbohydrates and sugar contents of the plants (Tümová *et al.*, 2018). The previous research resulted in increased plant growth and development by the application of NAA (Kareem *et al.*, 2017). As virus induces

decrease in protein, minerals and vitamin contents of the infected plants; the efficacy of elephant ear plant extract (*Clocasiaesculenta*) was attributed to its abundance of proteins, ascorbic acid, minerals and vitamins in its constituents (Chakraborty *et al.*, 2015). Ascorbic acid provides the basic defense against destructive effects of reactive oxygen species (ROS) (Gallie, 2013).

However, despite the clear importance of virus symptom severity as a driver of both yield losses and virus spread, strategies for mitigating symptoms and enhancing plant tolerance to virus infection are rarely considered as components of integrated disease management. It can be concluded that cultivars selection, proper dose of nutrients and plant extract application could be an effective integrated approach to manage CMV disease. In present study, spatio-temporal disease increment was recorded with each assessment. All the applied treatments induced strong defense responses in plants which resulted in significant disease severity reduction.

Comparison of chlorophyll contents and leaf area in diseased and healthy plants: Chlorophyll content values noted from the chlorophyll meter for each leaf sample and then average was taken which was plotted for five plants. There was significant decrease in the chlorophyll contents of diseased and healthy leaves in all plants. Maximum chlorophyll contents (2.7 mg g^{-1}) were recorded in case of healthy plants which were reduced upto (0.9 mg g^{-1}) in diseased plants (Fig. 6). There was significant reduction in chlorophyll contents of susceptible plants infected with CMV (Patel *et al.*, 2013).




Fig. 6. Comparison of chlorophyll contents in diseased and healthy plants.

There was clear difference in leaf area of diseased and healthy plants with maximum (70 cm^2) in healthy plants and minimum (27 cm^2) in diseased plants (Fig. 7). Leaf area reduction was recorded after mechanical inoculations in many cucumber advance lines along with other growth parameters (Azizi & Shams-Bakhsh, 2014).

ELISA for CMV detection: The samples collected from all locations were positive for the presence of CMV both in leaf lamina and midrib (Fig. 8; Table 6). The virus titer was high in leaf midrib than leaf lamina in all the samples. Optical density (OD) values were higher for the samples of BT than COA and ARF respectively and very strong reaction was observed in those samples. There was positive relationship between symptom severity and absorbance values of serological assays in different infected samples (Ashfaq *et al.*, 2012). These results in line with previous serological studies where moderately resistant samples gave low absorbance values and high values were observed when ELISA performed on susceptible and highly susceptible samples (Iqbal *et al.*, 2012). Arogundade *et al.*, (2019) detected CMV from infected samples by using a modified method of ELISA that was antigen coated plate (ACP). Kumar *et al.*, (2016) observed high virus titer in leaf mid rib than in lamina which strengthened the results of current experiment. Seed samples of cucumber were used for CMV detection by using anti rabbit polyclonal antibodies (Yoon *et al.*, 2019).




Fig. 7. Comparison of leaf area in healthy CMV infected cucumber plants.

Table 4. Disease severity as influenced by the treatments at various intervals

Treatments	Sprays			Mean	Efficacy (%)
	S ₁	S ₂	S ₃		
NPK solution	25.36±1.12	16.82±1.64	10.59±1.26	17.59 e	75.55 a
NAA*	40.18±1.24	32.75±1.55	24.04±1.34	32.33 c	55.06 c
Elephant plant extract	50.14±1.09	44.32±2.24	36.26±1.26	43.57 b	39.44 d
BTH	33.44±1.26	25.71±2.01	19.24±1.29	26.13 d	63.68 b
Control	70.07±1.22	70.88±1.32	74.91±1.63	71.95 a	---
Mean**	41.25 a	29.9 b	22.53 c		---

Means sharing similar letter in a row or in a column are statistically non-significant ($p > 0.05$)

*NAA = Naphthalene acetic acid; **Means don't include control values



Fig. 8. Microtitre plate depicting ELISA results.

Table 5. Disease severity as influenced by the treatments at different concentrations.

Treatments	Concentrations				Pattern
	C ₁	C ₂	C ₃	z-score	
NPK solution	22.08 ± 1.41	12.91±1.73	8.66±1.53	-0.94	Random
NAA	38.34 ± 1.27	29.09±1.36	22.64±1.02	-0.26	Random
Elephant plant extract	48.97 ± 1.13	41.04±1.25	33.58±1.34	0.24	Random
BTH	30.52±1.05	19.13±1.16	14.83±1.82	-0.64	Random
Control	69.05±1.25	71.54±1.56	75.86±1.47	1.61	Random
z-score for conc.	1.07	-0.16	-0.91		
Pattern	Random	Random	Random		

Means sharing similar letter in a row or in a column are statistically non-significant ($p>0.05$)

*NAA = Naphthalene acetic acid

Conclusions

The spatio-temporal disease assessments provide a road map to devise sustainable disease management strategies. The virus disrupts the plant physiology and its biochemical contents. All the treatments used, worked to repair the physiological and biochemical damages incited by CMV. The extraordinary results were noted with the plant's ability to defend against viruses with application of NPK. This contribution of NPK is an essential landmark for sustainable agriculture. Many environmental concerns convince the growers to use nutrients like NPK for disease management. It would be helpful to get rid from hazardous synthetic chemicals.

References

- Ahmed, A., Z. Ahmad, F. Begum, Ubairah and N. Raees. 2015. Varietal reaction of cucumber against *Cucumber mosaic virus*. *Ame. J. Plant Sci.*, 6: 833-838. <http://dx.doi.org/10.4236/ajps.2015.67090>.
- Akhtar, K.P., M.Y. Saleem, M. Asghar, M. Ahmad and N. Sarwar. 2010. Resistance of Solanum species to cucumber mosaic virus subgroup IA and its vector *Myzus persicae*. *Eur. J. Plant Pathol.*, 128: 435-450.
- Ali, A., M.A. Zeshan, Y. Iftikhar, M. Abid, S.F. Ehsan, M.U. Ghani and A.A. Khan. 2020. Role of plant extracts and salicylic acid for the management of *chili vein mottle virus* disease. *Pak. J. Phytopath.*, 32(2): 147-157.
- Al-Zahrani, H.S.M., E.K.F. Elbeshy, A.Y. Aldhebiani and T. Elbeaino. 2018. Effect of cucumber mosaic virus (CMV) infection on antineoplastic alkaloids from periwinkle (*Catharanthus roseus* L.) culture in Mecca region and resistance induction by plant growth promoting rhizobacteria (PGPR). *Biotech. Biotechnol. Equip.*, 32(1): 49-57.
- Anitha, S.S. 2016. Pharmacological activity of vinca alkaloids. *Res. Rev. J. Pharm. Phytochem.*, 4(3): 27-34.
- Arogrundade, O., O.S. Balogun and P.L. Kumar. 2019. Seed transmissibility of Cucumber mosaic virus in Capsicum species, *Int. J. Veg. Sci.*, 25(2): 146-153. DOI: 10.1080/19315260.2018.1487498.
- Arogrundade, O., A. Oresanya, J.O. Matthew, K.T. Kareem and H. Onyeanus. 2019. Natural occurrence of cucumber mosaic virus in Ogeira (*Eleutheranthera ruderalis*) in

- Nigeria, *Arch. Phytopath. Plant Protect.*, 52; 1-2, 108-113, DOI: 10.1080/03235408.2019.1585407.
- Azizi, A. and M. Shams-Bakhsh. 2014. Impact of cucumber mosaic virus infection on the varietal traits of common bean cultivars in Iran. *Virus Dis.*, 25(4): 447-454.
- Aznar, A., N.W.G. Chen, S. Thomine and A. Dellagi. 2015. Immunity to plant pathogens and iron homeostasis. *Plant Sci.*, 240: 90-97.
- Bashir, M., Z. Ahmad and A. Ghafoor. 2002. Cowpea aphid-borne mosaic potyvirus: A review. *Int. J. Pest Manag.*, 48: 155-168.
- Borer, E.T., A.L. Laine and E.W. Seabloom. 2016. A multi-scale approach to plant disease using the meta community concept. *Annu. Rev. Phytopathol.*, 54: 397-418. doi: 10.1146/annurev-phyto-080615-095959.
- Byamukama E., A.E. Robertson and F.W. Nutter. 2011. Quantifying the within-field temporal and spatial dynamics of *Bean pod mottle virus* in soybean. *Plant Dis.*, 95: 126-136.
- Byamukama, E., A.R. Robertson and F.W. Nutter. 2010. Quantification of temporal and spatial dynamics of *Bean pod mottle virus* at different spatial scales. Online. *Plant Health Prog.*, doi:10.1094/PHP-2010-05XX-01-SY.
- Cao, S., Z. Hu, Y. Zheng, Z. Yang and B. Lu. 2011. Effect of BTH on antioxidant enzymes, radical-scavenging activity and decay in strawberry fruit. *Food Chem.*, 125: 145-149.
- Casteel, C.L., M. De Alwis, A. Bak, H. Dong, S.A. Whitham and G. Jander. 2015. Disruption of ethylene responses by turnip mosaic virus mediates suppression of plant defense against the green peach aphid vector. *Plant Physiol.*, 169: 209-218.
- Chakraborty, P., P. Deb, S. Chakraborty, B. Chatterjee and J. Abraham. 2015. Cytotoxicity and antimicrobial activity of *Colocasia esculenta*. *J. Chem. Pharm. Res.*, 7(12): 627-635.
- Chandankar, V.D., M.K. Mondhe, P.R. Bhojar, B.N. Ninawe and G. Jadesha. 2013. Biophysical characterization, host range and transmission studies of Cucumber mosaic virus. *The Bioscan.*, 8(2):437-441.
- Chem, M., W. Bai, X. Chen, P.E. Canlas and P.C. Ronald. 2013. Reduced expression of glycolate oxidase leads to enhanced disease resistance in rice. *Peer J.*, 1: e28. doi:10.7717/peerj.28.
- Clark, M.F. and A.N. Adams. 1977. Characteristics of the microplate method of enzyme linked immune-sorbent assay for the detection of plant viruses. *J. Gen. Virol.*, 34: 475-483.
- Crespo, O., D. Janssen, C. Robles and L. Ruiz. 2018. Resistance to cucumber green mottle mosaic virus in *Cucumis sativus*. *Euphyt.*, 214: 201. <https://doi.org/10.1007/s10681-018-2286>.
- Donnelly, R., N.J. Cunniffe, J.P. Carr and C.A. Gilligan. 2019. Pathogenic modification of plants enhances long-distance dispersal of non-persistently transmitted viruses to new hosts. *Ecol.*, 100e02725.
- Gallie, D.R. 2013. The role of l-ascorbic acid recycling in responding to environmental stress and in promoting plant growth. *J. Exp. Bot.*, 64: 433-443.
- Gao, W., Y. Wang, C. Zhang, A. Zhang and X. Zhu. 2016. Investigation and pathogen preliminary identification of pepper virus disease in Tianjin. *Shandong Agric. Sci.*, 48: 91-94. doi: 10.14083/j.issn.1001-4942.2016.03.023.
- Gougherty, A.V. and F.W. Nutter. 2012. An epidemiological comparison of the US and Canadian Plum pox virus eradication programs. Online. *Plant Health Prog.*, doi:10.1094/PHP-2012-0723-03-RS.
- Gougherty, A.V. and F.W. Nutter. 2015. Impact of eradication programs on the temporal and spatial dynamics of plum pox virus on *Prunus* spp. in Pennsylvania and Ontario, Canada. *Plant Dis.*, 99: 593-603.
- Groen, S.C., F.O. Wamonje, A.M. Murphy and J.P. Carr. 2017. Engineering resistance to virus transmission. *Curr. Opin. Virol.*, 26: 20-27.
- Guo, G., S. Wang, J. Liu, B. Pan, W. Diao and W. Ge. 2017. Rapid identification of QTLs underlying resistance to cucumber mosaic virus in pepper (*Capsicum frutescens*). *Theor. Appl. Genet.*, 130: 41-52. doi: 10.1007/s00122-016-2790-3.
- Gupta, K.J., Y. Brotman and S. Segu. 2013. The form of nitrogen nutrition affects resistance against *Pseudomonas syringae* pv. *phaseolicola* in tobacco. *J. Exp. Bot.*, 64: 553-568.
- Hakeem, K.R., F. Khan, R. Chandna, T.O. Siddiqui and M. Iqbal. 2012. Proteomic analysis for low and high nitrogen responsive proteins in the leaves of rice genotypes grown at three nitrogen levels. *Appl. Biochem. Biotechnol.*, 16: 834-850.
- Hasanuzzaman, M., M.H.M.B. Bhuyan, K. Nahar, M.S. Hossain, J.A. Mahmud, M.S. Hossen, A.A.C. Masud, Moumita and M. Fujita. 2018. Potassium: A Vital Regulator of Plant Responses and Tolerance to Abiotic Stresses. *Agron.*, 8, 31; doi:10.3390/agronomy8030031.
- Helal, I.M. 2019. Use of biocides for controlling viral diseases that attack common bean and cucumber plants. *Folia Hort.*, 31(1): 159-170.
- Hull, R. 2014. *Plant Virology*. 5th ed. Academic Press, London, UK. pp. 854.
- Hout, B., J. Yao, B.L. Montgomery and S.Y. He. 2014. Growth-defense tradeoffs in plants: a balancing act to optimize fitness. *Mol. Plant.*, 7: 1267-1287.
- Iqbal, S., M. Ashfaq, H. Shah, M.I. Haq and A. Din. 2012. Prevalence and distribution of cucumber mosaic Cucumovirus (CMV) in major chili growing areas of Pakistan. *Pak. J. Bot.*, 44: 1749-1754.
- Jo, Y.D., Y. Ha, J.H. Lee, M. Park, A.C. Bergsma and H.I. Choi. 2016. Fine mapping of *Restorer-of-fertility* in pepper (*Capsicum annuum* L.) identified a candidate gene encoding a pentatricopeptide repeat (PPR)-containing protein. *Theor. Appl. Genet.*, 129: 2003-2017. doi: 10.1007/s00122-016-2755-6.
- Kareem, F., H. Rihan and M. Fuller. 2017. The effect of exogenous applications of salicylic acid and molybdenum on the tolerance of drought in wheat. *Agric. Res. Tech.*, 9(4): 555768.
- Kumar, A., B.R. Maurya, R. Raghuvanshi, V.S. Meena and M.T. Islam. 2017. Co-inoculation with *Enterobacter* and *Rhizobacteria* on yield and nutrient uptake by wheat (*Triticum aestivum* L.) in the alluvial soil under indo-gangetic plain of India. *J. Plant Growth Regul.*, doi:10.1007/s00344-016-9663-5.
- Kumar, S., P.S. Chauhan, L. Agrawal, R. Raj, A. Srivastava and S. Gupta. 2016. *Paenibacillus lentimorbus* inoculation enhances tobacco growth and attenuates the virulence of *Cucumber mosaic virus*. *PLoS ONE.*, 11(3): e0149980. <https://doi.org/10.1371/journal.pone.0149980>.
- Lambers, H. and M. Shane. 2007. Role of root clusters in phosphorus acquisition and increasing biological diversity in agriculture. *Frontis.*, 21: 235-248.
- Liu, Q., X. Chen, K. Wu and X. Fu. 2015. Nitrogen signaling and use efficiency in plants: what's new? *Curr. Opin. Plant Biol.*, 27: 192-198.
- Lu, Z., J. Lu, Y. Pan, P. Lu, X. Li, R. Cong and T. Ren. 2016. Anatomical variation of mesophyll conductance under potassium deficiency has a vital role in determining leaf photosynthesis. *Plant Cell Environ.*, 39: 2428-2439.
- Mandal, B., S. Mandal, A. Csinos, N. Martinez, A. Cullbreath and H. Pappu. 2008. Biological and molecular analyses of the Acibenzolar-S-Methyl- induced systemic acquired resistance in flue-cured tobacco against tomato spotted wilt virus. *Phytopath.*, 98: 196-204.
- Mauck, K.E. 2016. Variation in virus effects on host plant phenotypes and insect vector behavior: What can it teach about virus evolution? *Curr. Opin. Virol.*, 21: 114-123.

- Mauck, K.E., J. Kenney and Q. Chesnais. 2019. Progress and challenges in identifying molecular mechanisms underlying host and vector manipulation by plant viruses. *Curr. Opin. Insect Sci.*, 33: 7-18.
- Meddad-Hamza, A., A. Beddiar, A. Gollotte, M.C. Lemoine, C. Kuszala and S. Gianinazzi. 2010. Arbuscularmycorrhizal fungi improve the growth of olive trees and their resistance to transplantation stress. *Afric. J. Biotechnol.*, 9: 1159-1167.
- Meena, V.S., J.P. Verma and S.K. Meena. 2016. Towards the current scenario of nutrient use efficiency in crop species. *J. Clean Prod.*, 102: 556-557. doi:10.1016/j.jclepro.2015.04.030.
- Mochizuki, T. and S. Ohki. 2012. Cucumber mosaic virus: viral genes as virulence determinants. *Mol. Plant Pathol.*, 13(3): 217-225.
- Mujtaba, A.V., A.K. Cherian, P.M. Namitha, V.I.M.I. Louis and S. Beena. 2019. Detection and biophysical characterization studies of cucumber mosaic virus causing infectious chlorosis disease of banana. *J. Pharmac. Phytochem.*, 8(1): 2606-2611.
- Mur, L.A., C. Simpson, A. Kumari, A.K. Gupta and K.J. Gupta. 2016. Moving nitrogen to the centre of plant defense against pathogens. *Ann. Bot.*, 119: 703-709.
- Naeem, M.S., Z.L. Jin, G.L. Wan, D. Liu, H.B. Liu, K. Yoneyama and W.J. Zhou. 2010. Aminolevulinic acid improves photosynthetic gas exchange capacity and ion uptake under salinity stress in oilseed rape (*Brassica napus* L.). *Plant Soil.*, 332: 405-415.
- Nalam, V., J. Louis and J. Shah. 2019. Plant defense against aphids, the pest extraordinaire. *Plant Sci.*, 279: 96-107.
- Namitharan, K., T.S. Zhu, J.J. Cheng, P.C. Zheng, X.Y. Li, S. Yang, B.A. Song and Y.R. Chi. 2014. Metal and carbene organocatalytic relay activation of alkynes for stereo selective reactions. *Nat. Commun.*, 5: 3982.
- Nischwitz, C., A.S. Csinos, S.W. Mullis, L.L. Hickman, K.L. Stevenson and R.D. Gitaitis. 2008. Effect of transplant age, tobacco cultivar Acibenzolar-S-methyl, and imidacloprid on *Tomato spotted wilt* infection in flue-cured tobacco. *Plant Dis.*, 92: 1524-1528.
- Ntui, V.O., K. Kong, P. Azadi, R.S. Khan, D.P. Chin, T. Igawa, M. Mii and I. Nakamura. 2013. RNAi-mediated resistance to cucumber mosaic virus (CMV) in genetically engineered tomato. *Amer. J. Plant Sci.*, 5: 554-572. <http://dx.doi.org/10.4236/ajps.2014.55071>.
- Nutter, F.W., E. Byamukama, R.A. Coelho-Netto, S.K. Eggenberger, M.L. Gleason, A. Gougherty, A.E. Robertson and N. Van Rij. 2011. Integrating GPS, GIS, and remote sensing technologies with disease management principles to improve plant health. In: *GIS Applications in Agriculture-Invasive Species*, 2nd ed.; Clay, S., ed. Taylor & Francis Group LLC: Boca Raton, FL. Pp. 59-90.
- Parvin, M.S., A.M. Akanda and A.H.M.A. Rahman. 2007. Summer Cosmos - A Host of Cucumber mosaic virus. *J. Agric. Rural Dev.*, 5(1-2): 84-93.
- Patel, H., R. Kalaria, M. Mahatma, D.A. Chauhan and L. Mahatma. 2013. Physiological and biochemical changes induced by mungbean yellow mosaic virus (MYMV) in mungbean [*Vigna radiata* (L.) Wilczek]. *J. Cell Tissue Res.*, 13: 3927-3930.
- Petrov, N., M. Stoyanova and M. Valkova. 2016. Antiviral activity of plant extract from *Tanacetum vulgare* against Cucumber mosaic virus and Potato virus Y. *J. Bio-Sci. Biotechnol.*, 5(2): 189-194.
- Qin, J., H. He, S. Luo and H. Li. 2013. Effects of rice-water chestnut intercropping on rice sheath blight and rice blast diseases. *Crop Prot.*, 43: 89-93.
- Rahman, M.S., A.U. Ahmed, K. Jahan and F. Khatun. 2020. Management of cucumber mosaic virus (CMV) infecting cucumber in Bangladesh. *Bangla. J. Agric. Res.*, 45(1): 65-76.
- Rahman, M.S., K. Jahan, A.A. Sabuz and A.M. Akanda. 2019. Effects of cucumber mosaic virus on cellular components, host physiology and yield of chili. *Plant Tissue Culture Biotechnol.*, 29(2): 219-230.
- Saldajeno, M.G.B. and M. Hyakumachi. 2011. The plant growth promoting fungus *Fusarium equiseti* and the arbuscularmycorrhizal fungus *Glomus mosseae* stimulate plant growth and reduce severity of anthracnose and damping off diseases in cucumber (*Cucumis sativus*) seedlings. *Ann. Appl. Biol.*, 159: 28-40.
- Sapno, R., D. Gallitelli and T. Mascia. 2017. Grafting to manage infections of top stunting and necrogenic strains of cucumber mosaic virus in tomato. *Ann. Appl. Biol.*, 171: 393-404.
- Shalitin, D. and S. Wolf. 2020. Cucumber mosaic virus infection affects sugar transport in melon plants. *Plant Physiol.*, 123: 597-604.
- Shaner, G. and R.E. Finney. 1977. The effect of nitrogen fertilization on the expression of slow-mildewing resistance in Knox wheat. *Phytopath.*, 67: 1051-1056.
- Shani, E., M. Salehin, Y. Zhang, S.E. Sanchez, C. Doherty, R. Wang, C.C. Mangado, L. Song, I. Tal and O. Pisanty. 2017. Plant stress tolerance requires auxin-sensitive Aux/IAA transcriptional repressors. *Curr. Biol.*, 27: 437-444.
- Sharma, S., A. Sharma and S. S. 2020. Kang. Insight in biophysical properties of *Pepper mottle virus* prevalent in Indian Punjab. *Plant Dis. Res.*, 33(2): 217-221.
- Steel, R.G.D., J.H. Torri and D.A. Dickey. 1997. Principles and procedures of statistics: A biometrics approach, 2nd ed. McGraw-Hill, New York, USA.
- Tiburcio, A.F., T. Altabella, M. Bitrian and R. Alcazar. 2014. The roles of polyamines during the lifespan of plants: from development to stress. *Planta.*, 240: 1-18.
- Tümová, L., D. Tarkowská, K. Řehořová, H. Marková, M. Kočová, O. Rothová, P. Čečetka and D. Holá. 2018. Drought-tolerant and drought-sensitive genotypes of maize (*Zea mays* L.) differ in contents of endogenous brassinosteroids and their drought-induced changes. *PloS One.*, 13(5): 0197870.
- Tungadi, T., R. Donnelly, L. Qing, J. Iqbal, A.M. Murphy, A.E. Pate, N.J. Cunniffe and J.P. Carr. 2020. Cucumber mosaic virus 2b proteins inhibit virus induced aphid resistance in tobacco. *Mol. Plant Pathol.*, 21(2): 250-257.
- Tungadi, T., S.C. Groen, A.M. Murphy, A.E. Pate, J. Iqbal, T.J.A. Bruce, N.J. Cunniffe and J.P. Carr. 2017. Cucumber mosaic virus and its 2b protein alter emission of host volatile organic compounds but not aphid vector settling in tobacco. *Viol. J.*, 14:91.
- Vitor, S.C., G.T. Duarte, E.E. Saviani, M.G.A. Vincentz, H.C. Oliveira and I. Salgado. 2013. Nitrate reductase is required for the transcriptional modulation and bactericidal activity of nitric oxide during the defense response of *Arabidopsis thaliana* against *Pseudomonas syringae*. *Planta*, 238: 475-486.
- Wu, D., T. Qi, W.X. Li, H. Tian, H. Gao, J. Wang, J. Ge, R. Yao, C. Ren, X.B. Wang, Y. Liu, L. Kang, S.W. Ding and D. Xie. 2017. Viral effectors protein manipulates host hormone signaling to attract insect vectors. *Cell Res.*, 27: 402-415.
- Yoon, J.Y., P. Palukaitis and S.K. Choi. 2019. Host range. In: (Eds.): Palukaitis, P., F. Garcia-Arenal. *Cucumber mosaic virus*, American Phytopathological Society, St. Paul, MN, USA, pp. 15-18.
- Yu, M., L. Lamattina, S.H. Spoel and G.J. Loake. 2014. Nitric oxide function in plant biology: a redox cue in deconvolution. *New Phytologist.*, 202: 1142-1156.
- Zhang, W.Y., X.Y. Li, G.P. Zhang, Y. Ding, L.L. Ran, L.Z. Luo, J. Wu, D.Y. Hu and B.A. Song. 2017. Binding interactions between enantiomeric α -aminophosphonate derivatives and tobacco mosaic virus coat protein. *Int. J. Biol. Macromol.*, 94: 603-610.