

CLONING OF PGPP2C2 GENE AND ANALYSIS OF ITS EXPRESSION IN RESPONSE TO NITROGEN AND PHOSPHORUS DEFICIENCY, AND EXOGENOUS GR24 TREATMENT IN PANAX GINSENG

HAO LIANG¹, HAI SUN¹, CAI SHAO¹, JIAQI QIAN¹, YAYU ZHANG^{1,2*}

¹*Institute of Special Economic Animal and Plant Sciences, Chinese Academy of Agricultural Sciences, Changchun 130112, China*

²*College of Pharmacy and Biological Engineer, Chengdu University, Chengdu 610106, China*

*Corresponding author's email: caassh@126.com

Abstract

Type 2C protein phosphatases (PP2Cs) are the most abundant members of the plant phosphatase family and perform vital functions in response to various environmental stressors. To explore the role of *Panax ginseng* C.A. Meyer (*P. ginseng*) *PgPP2C*, a gene identified from a previously assembled transcriptome database, under adverse conditions, we created a clone of the gene and named it *PgPP2C2*. The length of its open reading frame was 1269 base pairs, and it coded for a 422-amino-acid protein that contained a 2C family serine/threonine phosphatase catalytic domain characteristic of the phosphatase 2C family proteins. Phylogenetic analysis of *PgPP2C2* protein sequences from multiple species revealed that the *P. ginseng* sequence was similar to homologous sequences from *Macleaya cordata*, *Malus domestica*, *Nicotiana tomentosiformis*, *Solanum lycopersicum*, and *Handroanthus impetiginosus*. Transient expression analysis identified a *PgPP2C2*–GLOsfp fusion protein in the nucleus. The expression of *PgPP2C2* was detected by transcriptional analysis in all investigated tissues, but at a higher level in the roots. Moreover, nitrogen (N) and phosphorus (P) deficiency treatments were observed to trigger *PgPP2C2* expression. Specifically, the downregulation of *PgPP2C2* and the accumulation of abscisic acid–glucose ester (ABA–GE) were induced under N deficiency. The synthetic strigolactone analog GR24 restored the expression of *PgPP2C2* under N deficiency, but had no obvious effect on the ABA–GE content. The findings enhance our understanding of the genetic regulatory mechanism of *P. ginseng* in response to N and P deficiencies, as well as provide a potential strategy for improving the quality of cultivation of *P. ginseng* through hormone-mediated nutritional regulation.

Key words: *Panax ginseng*, *PgPP2C2*, Molecular cloning, Expression, Strigolactone, Nitrogen (N) and phosphorus (P).

Abbreviations: PP2Cs, Type 2C protein phosphatases; ABA, abscisic acid; SnRK2, SNF1-related kinase 2; ORF, open reading frame; MAPK, mitogen-activated protein kinases; HOG1, high osmolarity glycerol 1; NBP2, NAPI-binding protein 2.

Introduction

The perennial herb ginseng (*Panax ginseng* C.A. Meyer) is a member of the Araliaceae family. It is known as the “king of herbs” and is the most prized Chinese medicinal material in East Asia and North America (Wen & Zimmer, 1996). Ginsenosides are the primary medicinal ingredients of ginseng and are used to treat cardiovascular diseases, age-related diseases, obesity, and stress (Chu & Zhang, 2009; Yang *et al.*, 2012; Kim *et al.*, 2014). However, in recent years, the cultivation of ginseng has encountered adverse conditions, such as climate warming, drought, and nutrient shortage, which have limited its growth. Consequently, it is imperative to identify stress resistance genes and breed ginseng for higher stress tolerance to increase overall yield and meet the ever-increasing market demand.

Protein kinases phosphorylate proteins, whereas protein phosphatases dephosphorylate them. During the regulation of plant metabolism, the two protein types work together to regulate the phosphorylation of substrate proteins. Type 2C protein phosphatases (PP2Cs) comprise a multi-member protein family with distinctive N-terminal extensions and are an important subtype of plant phosphatases. PP2Cs are serine/threonine protein phosphatases that exist in the form of monomeric enzymes. Their dephosphorylation activity depends on Mn^{2+} and Mg^{2+} , and they are widespread in animals, plants, bacteria, and fungi (Hu *et al.*, 2020); in particular, higher plants produce a large number of diverse PP2C enzymes. These proteins are involved in a variety of stress signaling pathways, such as those that respond to drought, salinity, and nutrient deficiency, and perform critical functions in plants under stress. Moreover, as regulators of the signal transduction pathway, PP2Cs are involved in abscisic acid (ABA) signal transduction. The interaction between the receptor and PP2C releases downstream SNF1-related kinase 2 (SnRK2) that further phosphorylates downstream proteins (Fujii *et al.*,

2009; Umezawa *et al.*, 2009), resulting in ABA transcription and ion channel activation (Hubbard *et al.*, 2009; PriTal *et al.*, 2009; Weiner *et al.*, 2010). After the ABA receptor receives a molecular signal, it allosterically inhibits PP2C activity, thereby reducing or eliminating the inhibition of PP2C on the SnRK2 kinase and enhancing the phosphorylation of the substrate protein by SnRK2 kinase to regulate the overall response of the plant to ABA (Rigoulot *et al.*, 2019). PP2Cs have been categorized into ten major groups (A–J) with six members that fall outside these main groups and are encoded in the *Arabidopsis* genome (Chu *et al.*, 2021). There are 82 species? of PP2C in *Arabidopsis thaliana*, 90 in rice (*Oryza sativa* L.), 88 in pepper (*Capsicum annuum* L.), and 91 in tomato (*Solanum lycopersicum* Mill.) (Jiang, 2020). PP2C proteins regulate seed germination (Rovira *et al.*, 2021; Yu *et al.*, 2020), root growth (Miao *et al.*, 2020), stress resistance (Zhang *et al.*, 2021), hormone responses (Hu *et al.*, 2020; Wang *et al.*, 2021), and other processes. The expression profiles of PP2Cs have been shown to fluctuate significantly as a result of biotic and abiotic factors, such as drought, cold, light, salt, and plant hormones (Chu *et al.*, 2021; Yu *et al.*, 2020; Miao *et al.*, 2020; Zhang *et al.*, 2021; Wang *et al.*, 2021; Wang *et al.*, 2020a).

Currently, the *PP2C* gene sequences and expression profiles in *P. ginseng* are unclear. In our previous study, the *PgPP2C* gene was identified by high-throughput sequencing (Lei *et al.*, 2023). To the best of our knowledge, the present study is the first to demonstrate the first isolation of a cDNA clone for the *PP2C* gene (*PgPP2C2*). Bioinformatic analyses were performed on the identified polypeptide product to gain insight into its gene homology, phylogeny, amino acid sequence, structure, and physiochemical characteristics. This study also offers the first detailed examination of *PgPP2C2* subcellular localization and distribution analysis of *PgPP2C2* expression in various tissues. Moreover, it demonstrates the activity of *PgPP2C2* under N and P deficiency in *P. ginseng*. Currently, *P. ginseng* is being

promoted for ecological planting, however, its yield and quality under this system are limited by the unavailability of nutrients in the soil (Ma *et al.*, 2021). The findings of our study provide insight into the regulation of *PP2C* by exogenous hormones in response to N and P nutrient deficiency during the growth of *P. ginseng*, with implications for addressing the issue of soil nutrient limitation in the ecological planting of *P. ginseng* through nutritional management.

Material and Methods

Plant materials, cultivation, and stress treatments: *P. ginseng* seeds (*P. ginseng* collected and preserved in the *P. ginseng* planting base of Baishan City, Jilin Province, China) were germinated on wet Whatman filter paper in a rectangular culture tray (34 × 25 cm) under darkness for 3 days at 25°C. Germinated seeds were transferred to a hydroponic tank supplemented with pure water for 3 days at 70% relative humidity and 22–25°C ambient temperature. The seedlings were then cultured for 5 days in Hoagland's solution (Hoagland & Arnon, 1950). Subsequently, 40 uniform seedlings were selected and transplanted into each hydroponic tank, and the nutrient solution in the hydroponic tank was composed of 5 mmol L⁻¹ Ca(NO₃)₂·4H₂O, 5 mmol L⁻¹ KNO₃, 1 mmol L⁻¹ NH₄NO₃, 2.5 mmol L⁻¹ MgSO₄·7H₂O, 2 mmol L⁻¹ KH₂PO₄, 0.4 mmol L⁻¹ Fe-Na-EDTA, and 5 mL of trace elements (Hoagland & Arnon, 1950). The pH of the solution was adjusted to 5.8. For N and P treatments, the nutrient solution was also supplemented with 0 mmol L⁻¹ (–N) or 5 mmol L⁻¹ (+N) Ca(NO₃)₂·4H₂O and KNO₃, and 0 mol L⁻¹ (–P) or 2 mmol L⁻¹ (+P) KH₂PO₄. CaCl₂ and KCl were separately supplied to complement the Ca²⁺ and K⁺ concentrations under N- and P-deficient conditions. The nutrient solution was replaced every 2 days. After 72 h, 20 seedlings were randomly selected from each hydroponic tank, and their root, stem, and leaf organs were collected, rapidly frozen in liquid nitrogen, and preserved at –80°C for subsequent analyses. For the remaining 20 seedlings in each hydroponic tank, 10 mM synthetic strigolactone analog GR24 (Yuanye Bio-Technology Co., Ltd, Shanghai, China) was applied to the plants in the –N, –P, and –N–P groups every 8 h for 24 h. The experimental design included four groups, namely –N+GR24, –P+GR24, –N–P+GR24, and CK (without GR24). After treatment, roots were collected, rapidly frozen using liquid nitrogen, and preserved at –80°C for further use. Each treatment had three biological replicates.

RNA isolation and cDNA cloning: We used the RNAsimple Total RNA Kit (Tiangen, Beijing, China) following the manufacturer's instructions for RNA isolation. The RNA quality was assessed using the NanoPhotometer® spectrophotometer (IMPLEN, CA, USA) and 1.0% (w/v) agarose gel electrophoresis. Following the protocol provided by the manufacturer of RNase H-Reverse Transcriptase FastKing gDNA Dispelling RT SuperMix (Tiangen, Beijing, China), 9 µL of total RNA (1 µg/mL) was utilized to synthesize first-strand cDNA. Primers for *PgPP2C2* were designed based on a transcriptome database produced by our laboratory in a previous study (Lei *et al.*, 2023): forward primer *PgPP2C2*, 5'-ATGATGATCAA TACCAACCTAAAA-3' and reverse primer *PgPP2C2*, 5'-CGCCTAGTATTCTCTCTGATG-3'. The thermal cycling program for amplification was 95°C for 4 min; 35 cycles of 95°C for 30 s, 51°C for 35 s, and 72°C for 90 s; and a final extension at 72°C for 7 min. After PCR amplification, the target fragment was purified and subcloned using the TIANgel Midi Purification Kit (Tiangen, Beijing, China) and pEASY-T5 Zero Cloning Kit (TransGen Biotech, Beijing, China), respectively, following the manufacturer's instructions. Sequencing of the resulting product was conducted at Sangon Biotech (Shanghai, China), and the *PgPP2C2* cDNA sequence was obtained using the DNAMAN 6.0 Software based on fragment assembly.

Bioinformatic analysis of *PgPP2C2*: The open reading frame (ORF) of the *PgPP2C2* coding sequence, as well as the identity and similarity of *PgPP2C2* to other sequences, was determined using tools available on the NCBI website (<https://www.ncbi.nlm.nih.gov/>). Its protein structure and conserved domains were predicted using Prosite tools, including ProtParam (physicochemical properties), ProtScale (<http://web.expasy.org/protscale>), SOPMA (<https://npsa-prabi.ibcp.fr/>), TMHMM (<http://www.cbs.dtu.dk/services/TMHMM/>), and SignalP-5.0 (<http://www.cbs.dtu.dk/services/SignalP/>). Interaction proteins were predicted by STRING (<https://string-db.org/cgi/input>), and protein structures were analyzed using the Swiss Model (<https://www.swissmodel.expasy.org/>). The NCBI database was searched for protein homologs, and an online alignment tool (<http://www.uniprot.org/align>) was used to construct multiple sequence alignments of protein sequences from diverse species. To determine the relationships among *PgPP2C2* from *P. ginseng* and *PP2C* genes from other species, we performed a phylogenetic analysis with MEGA6.0 software using the JTT substitution model and the G + I model for rates among sites (Tamura *et al.*, 2013). In total, 1000 bootstrap replicates were used to determine the reliability of the internal branches. The neighbor-joining approach was used to generate a phylogenetic tree of *PP2C* homologs from various species.

Subcellular localization of *PgPP2C2*: To analyze the subcellular localization of *PgPP2C2*, the *PgPP2C2* ORF (minus the termination codon) was amplified from *PgPP2C2* cDNA with specific primers (SI-F: cagtGGTCTCacaacatgatgatcaataccaact and SI-R: cagtGGTCTCatacagtattctctctgatgatact). Subsequently, the *PgPP2C2* sequence was separately cloned into the pBWA(V)HS-ccdb-GLosgfp vector fused to the 5' terminus of GFP and driven by the 35S promoter. The pBWA(V)HS-*PgPP2C2*-Glosgfp fusion vectors and GFP empty constructs were introduced into the *Agrobacterium tumefaciens* strain GV3101. Transformed GV3101 cells were grown for 2 days at 30°C and inoculated into 10 mL of YEB liquid medium cultured at 170 rpm/min for 1 h. It was suspended with 10 mM MgCl₂ (including 120 µMAS) when the absorbance at 600 nm was 0.6. One-month-old *Nicotiana benthamiana* plant with good growth was selected and the leaf lower epidermis was injected using a 1 mL syringe without the gun head. The plants were cultured for 2 days under low light. The leaves injected with *Agrobacterium tumefaciens* were placed on slides, and confocal microscopy (Nikon C2-ER; Nikon, Tokyo, Japan) was performed to determine the subcellular localization of *PgPP2C2* based on the associated GFP fluorescence intensity. For co-localization, the marker plasmid was transformed into *Agrobacterium tumefaciens*, suspended with the constructed vector plasmid *Agrobacterium tumefaciens*, mixed in a 1:1 ratio before injection, and then injected into *Nicotiana benthamiana* leaves.

Expression analysis of *PgPP2C2* by quantitative real-time PCR (qPCR) in *P. ginseng*: The template for qPCR was the cDNA obtained by reverse transcription with the primers *PgPP2C2*-F (5'-TCAGTTAATCTCGCGGAAAGC-3') and *PgPP2C2*-R (5'-GTCTCGTCTTCGTTCTCAAGTT-3'). *GAPDH* served as the reference gene for normalization with the primers *GAPDH*-F (5'-AATGACCTTGCCGACAGCCTTG-3') and *GAPDH*-R (5'-TACACCGCCACCCAGAAGACC-3') (Zhang *et al.*, 2022). The 20 µL qPCR amplification system contained 0.5 µL of template cDNA, 0.5 µL of Primer F (10 µM, Sangon), 0.5 µL of Primer R (10 µM, Sangon), and 8.5 µL of ddH₂O mixed with 10 µL of 2× qPCR Mix (Takara, Japan). All reactions were performed with three biological replicates. The

Fig. 2. Deduced amino acid sequence of *PgPP2C2* and overall length of the cDNA sequence. The red boxes represent the initiation and termination codons.




Fig. 3. Domain prediction of the PgPP2C2 protein.




Fig. 4. Interaction prediction of the PgPP2C2 protein.




Fig. 5. Tertiary structure prediction of the PgPP2C2 protein.

Phylogenetic analysis of protein sequences homologous to PgPP2C2: To elucidate the structure of the PgPP2C2-encoded protein sequence, we constructed multiple sequence alignments of PgPP2C2 and other plant homologs obtained from the NCBI database. The primary structure of PgPP2C2 was quite similar to that of PgPP2C2-like proteins from five other species, all of which exhibited a substantial degree of sequence conservation (Fig. 6). According to previous studies, PP2C protein phosphatase is a Mg^{2+}/Mn^{2+} -dependent monomeric enzyme that is highly conserved in numerous diverse plant species (Yu *et al.*, 2008; Cao *et al.*, 2013). Thereafter, we used the neighbor-joining method in MEGA 6.0 to generate a phylogenetic tree of the PgPP2C2 and homologous protein sequences from 20 related plant species obtained from the NCBI database. PgPP2C2 was the most closely related to sequences from *Macleaya cordata* (OVA01953.1), *Malus domestica* (XP_028962043.1), *Nicotiana tomentosiformis* (XP_009592346.1), *Solanum lycopersicum* (NP_001333898.1), and *Handroanthus impetiginosus* (PIN20518.1), and was more distantly related to sequences from other species (Fig. 7).

Currently, the PP2C genes of *Malus domestica* and *Solanum lycopersicum* have been sequenced (Yang *et al.*, 2022; Chen *et al.*, 2016). The *PYL* gene in *Malus domestica* and *Solanum lycopersicum* interacts with PP2CAs in ABA-dependent and -independent manners and is affected by characteristics of the PP2C protein (Yang *et al.*, 2022; Chen *et al.*, 2016). Our analysis suggests that PgPP2C2 may have functions similar to that of PP2C proteins in *Malus domestica*, *Solanum lycopersicum*, and *Nicotiana tabacum*. These results laid a foundation for the analysis of the relationship between PgPP2C2 expression and the ABA content in subsequent experiments.

PgPP2C2-related subcellular localization: We developed a pBWA(V)HS-PgPP2C2-GLOsgfp vector and transiently expressed it in tobacco leaves via *Agrobacterium tumefaciens*-mediated transformation to validate the specific subcellular location of PgPP2C2. *Nicotiana benthamiana* cells transformed with the empty pBWA(V)HS-ccdb-GLOsgfp vector exhibited a uniform distribution of GLOsgfp fluorescence (Fig. 8A-B). In contrast, PgPP2C2-GLOsgfp fluorescence was localized to the nucleus. The PP2C protein is expressed differently in different plants. For example, AcPP2C3 and PbrPP2C are expressed in the nucleus (Wang *et al.*, 2021; Zhai *et al.*, 2023), BpPP2C1 is expressed in the nucleus and cell membrane (Xing *et al.*, 2021), and AtPP2C49 is expressed in the cytoplasm and nucleus (Chu *et al.*, 2021). These discrepancies in expression localization reflect differences in the functions of PP2C proteins and are also likely related to the diversity of PP2C proteins, which represent the largest protein phosphatase family.

Analysis of PgPP2C2 expression in various tissues and under different treatments: We extracted RNA from the roots, stems, and leaves of *P. ginseng* cultured for 72 h, reverse transcribed it into cDNA, and detected the tissue-specific expression of PgPP2C2 via qPCR. We discovered that PgPP2C2 expression varied among different organs of *P. ginseng* under normal conditions, with higher expression in the roots than in the aboveground tissues (Fig. 9A). The tissue specificities of PgPP2C2 gene expression were similar to those of AtPP2CG1 and AtPP2C49 in *Arabidopsis thaliana* (Chu *et al.*, 2021; Liu *et al.*, 2021). The root is the main medicinal part of *P. ginseng*. Our result suggests that PgPP2C2 may have an important function in the highly prized *P. ginseng*. The gene expression of PP2Cs is widespread in different plants and their tissues. For example, the cotton PP2C gene, *GhDRP1*, has high expression in the cotyledons and anthers (Chen *et al.*, 2021), whereas the oilseed rape PP2C gene *BcAB11* is expressed at high levels in the pods, flowers, and leaves (Kong *et al.*, 2018). In the present study, the broad expression profile of the PgPP2C2 gene across multiple organs suggests that it may also have regulatory functions in a variety of tissues throughout *P. ginseng* development.

To analyze the effect of N and P deficiencies on PgPP2C2 expression, we used cDNA isolated from ginseng roots in the -N, -P, -N-P, and CK groups and performed PCR analysis. The expression of PgPP2C2 in the -N, -P, and -N-P groups was lower than that in the control group, with the lowest expression observed in the -N group (Fig. 9B).

PgPP2C	RVTSSSR	SDTTQTAR	RAHSLA	DA	C	GS	Y	G	K	I	S	V	I	G	R	130																								
McPP2C	GGGPPTV	GIIIVAGG	SEICGEEELRS	LQ	CV	SH	GS	V	SV	CG	R						159																								
MdPP2C51	SGSVAGS	SDKGGGKRG	SEGL	GARG	G	Y	GV	V	SL	K	G	M				119																								
NtPP2C51	TTSS	EVKT	VRES	LQR	VT	CG	SH	G	L	I	SL	I	G	R	120																								
SlPP2C	ALSS	EVKKV	RES	L	VT	CC	SH	GS	I	SL	I	G	R		114																								
HiPP2C	VQLP	ENDV	VGS	RE	VT	CL	SH	GT	I	SV	I	G	R		118																								
Consensus											g		s		g																										
PgPP2C	RREMEDAV	TVEL	GFV	SAS	SRKY	DDFFG	VYDGHG	HGS	SRV							166																								
McPP2C	RREMEDAV	TVVP	GLVS	DHKY	DDFFA	VYDGHG	HGS	SGV								193																								
MdPP2C51	RKEMEDAV	RVEV	GF	..	G	KF	DFY	G	VYDGHG	HG	PD	V				150																								
NtPP2C51	RREMEDAV	TVKP	G	LLSKG	NRKY	DDFFG	VYDGHG	HGS	SRV							156																								
SlPP2C	RREMEDAV	A	IYP	CFFSE	G	GGGGGS	RRY	DYF	G	VYDGHG	HGS	SRV					154																								
HiPP2C	RREMEDAV	A	VEL	G	FLKR	G	E	KRY	DDFFG	VYDGHG	HG	WRV				154																								
Consensus	r	emedav						d			v y d g h g g																														
PgPP2C	AHAC	R	D	R	L	H	H	L	V	V	K	E	V	E	E	E	189																								
McPP2C	ANVCR	D	R	L	H	R	V	L	V	K	E	I	E	E	R	E	T	233																							
MdPP2C51	AGAC	R	E	R	M	H	E	V	V	A	E	A	V	E	M	E	G	I	183																						
NtPP2C51	AHAC	R	E	W	L	H	R	L	V	I	E	E	I	V	E	E	E	E	184																						
SlPP2C	ANVCR	D	F	L	H	R	L	V	I	Q	V	S	E	G	E	D	Y	182																						
HiPP2C	AQS	C	G	E	M	L	H	K	M	L	R	N	V	L	E	E	S	179																						
Consensus	a	c																h																							
PgPP2C	WE	K	V	M	V	A	S	F	G	K	M	D	E	E	V	I	A	S	225																					
McPP2C	WEN	V	M	V	S	S	F	L	K	M	D	E	E	V	N	C	R	G	I	E	L	273																			
MdPP2C51	WET	V	M	E	E	C	F	G	K	I	D	G	E	V	S	D	I	A	A	R	M	I	G	A	T	A	V	A	V	V	T	218								
NtPP2C51	WE	K	V	M	R	E	S	F	V	K	M	D	E	E	V	E	K	K	G	A	E	M	A	T	M	G	S	T	A	V	V	A	M	V	S	220				
SlPP2C	WE	K	V	M	T	E	S	F	R	K	M	D	E	K	V	N	K	E	G	A	E	M	A	T	I	G	S	T	A	V	V	A	V	G	218					
HiPP2C	WE	R	V	M	A	A	G	F	K	R	M	D	E	E	V	K	K	S	G	A	A	V	A	T	T	G	S	T	A	T	V	A	V	I	G	215				
Consensus	w	e		v	m			f			d		v												g	t	a		v	a											
PgPP2C	EE	V	V	V	A	N	C	G	D	S	R	A	V	L	S	R	A	R	V	P	M	P	L	S	N	D	H	K	P	D	R	P	D	E	L	E	R	I	265		
McPP2C	RN	K	I	I	V	A	N	C	G	D	S	R	A	V	L	S	R	G	G	A	A	V	P	L	S	R	D	H	R	P	D	R	T	D	E	M	E	R	V	313	
MdPP2C51	EE	Q	V	V	A	N	C	G	D	S	R	A	V	L	S	R	G	G	V	A	L	A	L	S	N	D	H	K	P	D	R	V	D	E	L	K	R	I	258		
NtPP2C51	EE	E	V	I	V	A	N	C	G	D	S	R	A	V	L	S	R	G	G	V	T	V	P	L	S	I	D	H	K	P	N	R	P	D	E	L	D	R	I	260	
SlPP2C	V	E	E	F	V	A	N	C	G	D	S	R	A	V	L	S	R	A	G	V	A	V	P	L	S	I	D	H	K	P	D	R	P	D	E	L	D	R	I	258	
HiPP2C	EE	E	V	V	A	N	C	G	D	S	R	A	V	L	S	R	G	S	G	A	V	Q	L	S	E	D	H	K	P	D	R	P	D	E	L	E	R	I	255		
Consensus																																									
PgPP2C	EV	S	G	G	R	V	I	N	W	N	G	Q	R	V	L	G	V	L	A	T	S	R	S	I	G	D	Q	Y	L	K	P	Y	V	I	A	Q	P	E	V	305	
McPP2C	EA	A	G	G	R	V	I	N	W	N	G	Y	R	V	L	G	V	L	A	T	S	R	S	I	G	D	H	Y	L	K	P	Y	V	I	S	V	P	E	V	353	
MdPP2C51	EA	A	G	G	R	V	I	T	W	N	G	N	R	V	L	G	V	L	A	T	S	R	S	I	G	D	H	Y	L	R	P	Y	V	I	S	K	P	E	V	298	
NtPP2C51	EK	S	G	G	K	V	I	N	W	N	G	H	R	V	L	G	V	L	A	T	S	R	S	I	G	D	T	Y	L	K	P	Y	V	I	P	D	P	E	V	300	
SlPP2C	EN	S	G	G	K	V	I	N	W	N	G	Q	R	V	L	G	V	L	A	T	S	R	S	I	G	D	M	Y	L	K	P	Y	V	I	P	D	P	E	V	298	
HiPP2C	EQ	I	G	G	K	V	I	N	W	N	G	Q	R	V	L	G	V	L	A	T	S	R	S	I	G	D	E	Y	L	K	P	F	V	I	S	E	P	E	V	295	
Consensus	e																																								
PgPP2C	TV	S	K	R	N	D	R	D	E	F	L	I	L	A	S	D	G	L	W	D	V	I	T	N	E	L	A	C	R	V	V	R	K	C	L	D	G	R	L	345	
McPP2C	T	I	T	R	R	N	E	S	D	E	F	M	I	L	A	S	D	G	L	W	D	V	L	S	N	E	V	A	C	D	V	A	R	R	C	L	E	G	R	I	393
MdPP2C51	TV	T	K	R	T	D	Q	D	E	F	L	I	L	A	S	D	G	L	W	D	V	M	S	N	D	V	A	C	Q	V	V	K	K	C	L	G	G	R	.	337	
NtPP2C51	K	V	S	K	R	T	D	A	D	E	F	L	I	L	A	S	D	G	L	W	D	V	I	P	N	D	V	A	C	D	I	T	R	R	C	L	N	G	Q	339	
SlPP2C	I	V	S	K	R	S	D	E	D	E	F	L	I	L	A	S	D	G	L	W	D	V	I	P	N	D	V	A	C	D	V	T	R	R	C	L	N	G	Q	T	338
HiPP2C	K	V	I	K	R	T	D	A	D	E	F	L	I	L	A	S	D	G	L	W	D	V	V	S	N	N	V	A	C	Q	V	V	R	R	C	L	D	G	K	.	334
Consensus																																									
PgPP2C	RR	R	F	S	Q	G	I	V	N	E	E	T	S	G	G	H	D	D	D	D	D	D	D	D	D	D	D	E	I	N	I	V	N	D	E	S	R	A	I	385	
McPP2C	I	R	R	F	S	S	D	T	T	T	T	I	T	T	S	G	S	G	S	G	S	G	S	G	V	A	419				
MdPP2C51	M	R	R	V	S	I	H	D	E	H	Q	V	L	L	N	358			
NtPP2C51	M	K	R	C	S	R	Q	S	K	S	H	S	K	T	D	E	K	D	E	Q	T	S	E	V	K	K	E	S	L	A	A	370
SlPP2C	F	R	R	C	D	Q	Q	T	K	S	Y	K	R	D	361		
HiPP2C	I	K	R	N	S	Q	L	P	I	F	Q	D	N	I	P	G	364		
Consensus																																									
PgPP2C	EAA	A	V	L	A	E	L	A	V	A	R	G	S	E	D	N	I	S	V	V	V	V	E	L	N	K	P	G	Q	Y	H	Q	E	E	Y	422				
McPP2C	DAA	A	V	L	A	E	L	A	M	A	R	G	S	K	D	N	I	S	V	V	I	I	E	L	K	K	L	N	R	Q	L	A	S	454					
MdPP2C51	EAA	A	V	L	A	D	L	A	M	A	R	G	S	R	D	N	I	S	V	I	V	V	D	L	T	K	T	S	388				
NtPP2C51	QA	A	S	F	L	A	E	L	A	I	A	R	G	S	E	D	N	I	S	V	I	V	I	E	L	N	R	S	V	C	S	T	S	K	405				
SlPP2C	RA	A	S	F	L	A	E	L	A	I	A	R	G	S	R	D	N	I	S	V	I	V	N	L	N	R	S	V	R	S	S	I	D	S	397				
HiPP2C	EAA	A	V	L	A	E	L	A	I	A	R	G	S	R	D	N	I	S	V	I	V	V	L	K	S	F	G	L	G	Q	I	G	K	F	T	402				
Consensus	a	a																																							

Fig. 6. Multiple alignments of PP2C2 protein sequences from *Panax ginseng* and other plant species according to DNAMAN 6.0 analysis. The aligned proteins were *Macleanya cordata* (OVA 01953.1), *Malus domestica* (XP 028962043.1), *Nicotiana tomentosiformis* (XP 009592346.1), *Solanum lycopersicum* (NP 001333898.1), and *Handroanthus impetiginosus* (PIN 20518.1). The dark blue color denotes identical amino acids, whereas teal and pink colors indicate conserved amino acids.




Fig. 7. Phylogenetic analysis of PP2C2 proteins derived from several species generated in MEGA 6.0 using the neighbor-joining technique with 1000 bootstrap replicates. The bootstrap values are provided next to each node.

Notably, the expression of *PgPP2C2* in the $-N$ group was substantially lower than that in the control group ($p < 0.05$), indicating that N deficiency had the greatest impact on *PgPP2C2* expression. N-limited conditions could enhance the expressional

up-regulation of the *PP2C* genes (Effah *et al.*, 2018; Wen *et al.*, 2018), and a protein phosphatase gene that is rapidly induced by $-N$ is Os09g0325700, which encodes protein phosphatase 2 C 68 (PP2C68) in rice (Hsieh *et al.*, 2018). In contrast, our research showed that $-N$ induced the downregulation of *PgPP2C2*, which may be affected by abundant medicinal substances in *P. ginseng* root. However, *PP2C* overexpression in the medicinal plant *Ammopiptanthus mongolicus* facilitates N acquisition in *Arabidopsis thaliana* (Han *et al.*, 2018). Therefore, further functional studies of *PgPP2C2* are still needed. PP2C is crucial for plants to adapt to stress, and most researchers have focused on how it responds to salt (Chu *et al.*, 2021; Wang *et al.*, 2021), light (Rovira *et al.*, 2021), drought (Yu *et al.*, 2020) and hormones (Miao *et al.*, 2020). However, few studies have focused on its roles in nutrient stress responses, especially multi-element synergistic stress. *AtPP2C49* is expressed at high levels in root vascular tissues when exposed to NaCl stress, and the salt-induced sensitivity of *AtPP2C49* is mediated by the *Arabidopsis thaliana* high-affinity potassium transporter (*AtHKT1;1*). By negatively regulating *AtHKT1;1* activity, *AtPP2C49* mediates the systemic Na^+ distribution under salt stress (Chu *et al.*, 2021). Moreover, the overexpression of *AmPP2C* (*Ammopiptanthus mongolicus* PP2C) facilitates K and N acquisition driven by transpiration (Han *et al.*, 2018). The *PP2C* synergistic response to several nutrient elements is an important feature. In our results, *PgPP2C2* expression in the $-N-P$ group was higher than that in the $-N$ and $-P$ groups. Therefore, we speculate that, under $-N$ and $-P$ stress, N and P transport-related proteins promote or inhibit the expression of *PgPP2C2*, but further experiments are needed to confirm this. Our previous studies revealed the role of the *P. ginseng* PP2C protein in response to drought and water stress (Lei *et al.*, 2023), and it was shown that N deficiency could also induce *PgPP2C2* expression in this study, indicating that PP2Cs can participate in a variety of stress signal regulation. Although the role of *PgPP2C2* in response to $-N$ and their relationship have not been determined, our findings contribute to the study of these potential N regulatory genes and may provide a novel approach to enhance N use efficiency in *P. ginseng*.




Fig. 8. Subcellular localization of *PgPP2C2*. (A) From left to right: target protein fluorescence signal, chlorophyll fluorescence signal, bright-field image, and merged image. Scale bar, 20 μ m. (B) Empty vector (control). Panels are as described in (A). Scale bar, 20 μ m.




Fig. 9. *PgPP2C2* expression in various tissues and under different treatments. (A) Expression of *PgPP2C2* in distinct tissues of *P. ginseng*. (B) *PgPP2C2* expression in *P. ginseng* roots in response to N and/or P deficiency. Each bar represents the mean of six replicates with standard error. Different letters indicate significant differences ($p < 0.05$, Duncan's test).




Fig. 10. Effects of different N and P deficiencies on ABA-glucose ester (ABA-GE) content in *P. ginseng* roots. Each bar represents the mean of six replicates with standard error. Different letters indicate significant differences ($P < 0.05$; Duncan's test).

Effect of N and P deficiency on ABA content: The total ABA content in the roots of *P. ginseng* was measured in response to N and P deficiencies. The levels of ABA-glucose ester (ABA-GE), the main form of ABA, were significantly higher ($p < 0.05$) in the -N group than in the control group, and were higher in the -N group than in the -P and -N-P groups, indicating that -N treatment had the greatest effect on the ABA-GE content (Fig. 10). Plants accumulate ABA upon stress exposure and activate either ABA-independent or -dependent defensive stress response pathways (Yoshida *et al.*, 2006). Drought (Xie *et al.*, 2006; Zhang *et al.*, 2006), salt (Song *et al.*, 2023), and heat (Itai & Ben-Zioni, 1974) stress can induce ABA accumulation in plants. Notably, the ABA content also increases in plants that lack minerals, especially N (Xiong & Zhu, 2003). Similarly, we found that N deficiency induced a significant increase in the ABA-GE content of *P. ginseng* roots. N plays a role in regulating the production and transport of ABA, particularly during root development. Elevated concentrations of NO_3^- in roots are positively related to ABA biosynthesis (Signora *et al.*, 2001). In Wheat, NO_3^- may upregulate β -glucosidase 1 (BG1), which, in turn, de-conjugates the inactive ABA-GE to release bioactive ABA (Wang *et al.*, 2020b; Ondzighi *et al.*, 2016). Arabidopsis PP2Cs have been shown to act as negative modulators of ABA (Gosti *et al.*, 1999; Merlot *et al.*, 2001; Schweighofer *et al.*, 2004; Wasilewska *et al.*, 2008). When ABA binds to the pyrabactin resistance1/PYR1-like/regulatory components of ABA receptors (PYR/PYL/RCARs), it attenuates

the suppression of PP2Cs on protein kinase SnRK2s by inhibiting PP2Cs, phosphorylates transcription factors, such as ABI5 and RAV1, and activates the downstream ABA response gene (Li & Li, 2019). Through PCR analysis, we discovered that the expression level of the *PgPP2C2* gene under -N treatment was considerably reduced compared with that in the control group. Therefore, we speculate that, when plants are under N deficiency stress, they may downregulate *PgPP2C2* expression by accumulating ABA-GE. Research has shown that P deprivation induces the expression of the ABA biosynthesis genes, promoting ABA accumulation and activating the ABA signal pathway (Castro *et al.*, 2023; Ribot *et al.*, 2008). -N-induced genes, *PSY3* (phytoene synthase 3) and *BCH1* (beta-carotene hydroxylase 1) are associated with ABA biosynthesis (Hao *et al.*, 2000; Li *et al.*, 2008; Vallabhaneni & Wurtzel, 2009; Welsch *et al.*, 2008). -N can rapidly and strongly induce the expression of *PSY3* and *BCH1* in rice roots (Hsieh *et al.*, 2018). Similarly, our results showed that N deficiency triggered both the downregulation of *PgPP2C2* and the accumulation of ABA-GE. Therefore, *PgPP2C2* may act as a mediator of ABA modulation under N deficiency.

Exogenous GR24 restored *PgPP2C2* expression and ABA-GE levels: We conducted PCR analysis with cDNA derived from the roots of ginseng plants grown under various conditions to examine the influence of exogenous GR24 on *PgPP2C2* expression under N and P deficiency: -N+GR24, -P+GR24, -N-P+GR24, and CK. The results showed that exogenous GR24 restored *PgPP2C2* expression under N and P deficiency, to some extent (Fig. 11A-C). It has been reported that a variety of exogenous hormones regulate plant nutrition. Strigolactones play a fundamental role in plant responses to N and P nutrient (Marro *et al.*, 2022). Regarding the effect of exogenous GR24 on PP2C gene expression, a study on alfalfa showed that *MsPP2C50* expression was down regulated by GR24 treatment under normal conditions, whereas *MsPP2C1* expression was upregulated after GR24 treatment under drought stress (Yang *et al.*, 2022). In this study, plants treated with exogenous GR24 under N and P deficiency did not show ABA-GE recovery to CK levels (Fig. 11D-F). Therefore, GR24 may not have affected the synthesis of ABA-GE. However, -N+GR24 treatment resulted in attenuation of the *PgPP2C2* expression under -N treatment alone, with levels similar to those in the CK group (Fig. 11A). In contrast, the expression of *PgPP2C2* was lower after -P+GR24 and -N-P+GR24 treatments than that under -P and -N-P treatments (Fig. 11B-C), respectively. It has been reported that strigolactones regulate stomatal opening and closing by affecting the activity of the S-type anion channel SLAC1, which is independent of the ABA pathway (Lv *et al.*, 2018). Our results were consistent with

this in that, under N deficiency, GR24 appeared to alleviate stress by modulating the activity of *PgPP2C2* in an ABA-independent manner, but further research is still needed. Interestingly, the administration of GR24 in the -P and -N-P treatment groups did not restore the expression of *PgPP2C2*, but rather further downregulated it. N and P may, therefore, have a synergistic effect in this process that, in turn, affects the expression of *PgPP2C2*. Similar studies have illustrated that soybean P

homeostasis may be regulated in response to N deficiency by GmNLA1-1 and GmNLA1-3 (Zhou *et al.*, 2022); unlike our study, these did not mention whether the genes were related to GR24. Therefore, *PgPP2C2* may also exhibit redundant and diversified functions in the coordinated response to N and P. Although further experiments are needed to confirm this, our study shows that *PgPP2C2* is involved in the response to N and P deficiency, and its expression is affected by exogenous GR24.




Fig. 11. Effects of exogenous GR24 treatment on *PgPP2C2* expression (A-C) and ABA-glucose ester (ABA-GE) content (D-F) under N and/or P deficiency. Different letters indicate significant differences ($P < 0.05$; Duncan's test).

Conclusions

To elucidate the tolerance mechanisms of *P. ginseng* under diverse stressors, it would be useful to examine the characteristics and expressional profiles of resistance genes, as these play an active role in various biotic and abiotic stress responses. In this study, the cDNA of the *PP2C* gene in *P. ginseng*, *PgPP2C2*, was effectively cloned and characterized for the first time. Our research showed that the *PgPP2C2* protein exhibited nuclear localization, and *PgPP2C2* was spatially modulated at the transcription level and expressed primarily in the roots. This gene was also found to be stimulated by N and P deficiency, as was the accumulation of endogenous ABA. These findings suggest that *PgPP2C2* may represent an N and P nutrient stress gene that participates in ABA-associated pathways. Furthermore, exogenous GR24 restored or enhanced

the expression of *PgPP2C2* under N and P deficiency treatment. Therefore, *PgPP2C2* may represent a potential target gene to regulate the growth of *P. ginseng* under N and P deficiency through the application of GR24. This is different from the traditional regulation of ginseng growth through soil nutrients, but similar to plant internal genes and exogenous hormones in regulating ginseng growth, which may help farmers to optimize *P. ginseng* cultivation; however, this method still requires validation through actual application in a production setting. Briefly, the cloning and characterization of *PgPP2C2* shown in this study have remarkable implications. It establishes a foundation for further investigations of the mechanisms by which *P. ginseng* adapts to environmental stress, and cultivate "excellent shaped and high-quality" ginseng, as well as represents a valuable reference for future molecular biology research of medicinal plant cultivation.

Acknowledgments

This study was supported by the China Agriculture Research System of MOF [MARACARS-21], the Agricultural Science and Technology Innovation Program of CAAS (CAAS-XTX20190025-6), the National Key R&D Program of China (2021YFD1600902), the Non-Woodland Old Ginseng Land Remediation Technology and Cultivation Demonstration (LNFYZBDL-JL18-246C), and the Science and Technology Development Program of Jilin Province (20200708024YY and 20210401106YY).

References

- Cao, J., M. Jiang, P. Li and Z. Chu. 2013. Genome-wide identification and evolutionary analyses of the PP2C gene family with their expression profiling in response to multiple stresses in *Brachypodium distachyon*. *BMC Genom.*, 17: 175.
- Castro, V.P., P.A. Martínez-Melgarejo, F. Pérez-Alfocea, J. Tian and I.C. Dodd. 2023. Stem girdling enhances ABA-induced stomatal closure of phosphorus-deprived soybean plants. *Environ. Exp. Bot.*, 208: 105266.
- Chen, P., Y.F. Sun, W.B. Kai, B. Liang, Y.S. Zhang, X.W. Zhai, L. Jiang, Y.W. Du and P. Leng. 2016. Interactions of ABA signaling core components (SIPYLs, SIPP2Cs, and SlSnRK2s) in tomato (*Solanum lycopersicon*). *J. Plant Physiol.*, 205:67-74.
- Chen, Y., J.B. Zhang, N. Wei, Z.H. Liu, Y. Li, Y. Zheng and X.B. Li. 2021. A type-2C protein phosphatase (GhDRP1) participates in cotton (*Gossypium hirsutum*) response to drought stress. *Plant Mol. Biol.*, 107: 499-517.
- Chu, M., P. Chen, S. Meng, P. Xu and W. Lan. 2021. The Arabidopsis phosphatase PP2C49 negatively regulates salt tolerance through inhibition of AtHKT1;1. *J. Integr. Plant Biol.*, 63: 528-542.
- Chu, S.F. and J.T. Zhang. 2009. New achievements in ginseng research and its future prospects. *Chin. J. Integr. Med.*, 15: 403-408.
- Effah, Z., L. Li, J. Xie, B. Karikari, C. Liu, A. Xu and M. Zeng. 2018. Transcriptome profiling reveals major structural genes, transcription factors and biosynthetic pathways involved in leaf senescence and nitrogen remobilization in rainfed spring wheat under different nitrogen fertilization rates. *Genomics*, 114: 110271.
- Floková, K., D. Tarkowská, O. Miersch, M. Strnad, C. Wasternack and O. Novák. 2014. UHPLC-MS/MS based target profiling of stress-induced phytohormones. *Phytochemistry*, 105: 147-157.
- Fujii, H., V. Chinnusamy, A. Rodrigues, S. Rubio, R. Antoni, S.Y. Park, S.R. Cutler, J. Sheen, P.L. Rodriguez and J.K. Zhu. 2009. *In vitro* reconstitution of an abscisic acid signaling pathway. *Nature*, 462: 660-664.
- Gosti, F., N. Beaudoin, C. Serizet, A.A. Webb, N. Vartanian and J. Giraudat. 1999. ABI1 protein phosphatase 2C is a negative regulator of abscisic acid signaling. *Plant Cell*, 11: 1897-1910.
- Han, L., J. Li, M. Jin and Y. Su. 2018. Functional analysis of a type 2C protein phosphatase gene from *Ammopiptanthus mongolicus*. *Gene*, 653: 29-42.
- Hao, D., W. Nili, C. Fei, L. Xiang, X. Jing and X. Li. 2000. Characterization of the β -carotene hydroxylase gene DSM2 conferring drought and oxidative stress resistance by increasing xanthophylls and abscisic acid synthesis in rice. *Plant Physiol.*, 154: 1304-1318.
- Hoagland, D.R. and D.I. Arnon. 1950. The water culture method for growing plants without soil. *California Agricultural Experiment Station. Circular*, pp. 1-32.
- Hsieh, P.H., C.C. Kan, H.Y. Wu, H.C. Yang and M.H. Hsieh. 2018. Early molecular events associated with nitrogen deficiency in rice seedling roots. *Sci. Rep.*, 8: 12207.
- Hu, Q.T., D. Hou, Z.Y. Zhao, H.T. Wei and X.C. Lin. 2020. Identification and expression analysis of PP2C gene family in *Phyllostachys edulis*. *J. Agri. Bioch.*, 28: 1776-1787.
- Hubbard, K.E., N. Nishimura, K. Hitomi, E.D. Getzoff and J.I. Schroeder. 2009. Early abscisic acid signal transduction mechanisms: newly discovered components and newly emerging questions. *Gene. Dev.*, 24: 1695-1708.
- Itai, C. and A. Ben-Zioni. 1974. Regulation of plant response to high temperature, In: (Eds.): Bielecki, R.L., A.R. Ferguson and M.M. Cresswell. *Mechanisms of Regulation of Plant Growth. The Royal Society of New Zealand, Wellington*, pp. 477-482.
- Jiang, Y. 2020. Research on the mechanism of SIPP2C in regulating tomato senescence. *PhD dissertation. Shenyang Agricultural University, Shenyang, China*.
- Kim, K.H., I. Choi, Y.W. Lee, C.K. Cho, H.S. Yoo, S.B. Lee, S.H. Choi, K.R. Kwon and J.H. Jang. 2014. Target genes involved in antiproliferative effect of modified ginseng extracts in lung cancer A549 cells. *Acta Bioch. Biophys. Sin.*, 46: 441-449.
- Kong, L., H. Deng, S. Hu, F. Wang, L. Miao, C. Chen, K. Zhao and X. Yu. 2018. Isolation, expression, and evolution analysis of the type 2C protein phosphatase gene BcABI1 involved in abiotic and biotic stress in *Brassica campestris* ssp., chinensis. *Plant Growth Regul.*, 85: 317-327.
- Lei, H., H. Zhang, Z. Zhang, H. Sun, M. Li, C. Shao, H. Liang, H. Wu and Y. Zhang. 2023. Physiological and transcriptomic analyses of roots from *Panax ginseng* C.A. Meyer under drought stress. *Ind. Crop. Prod.*, 191: 115858.
- Li, F., R. Vallabhaneni and E.T. Wurtzel. 2008. PSY3, a new member of the phytoene synthase gene family conserved in the Poaceae and regulator of abiotic stress-induced root carotenogenesis. *Plant Physiol.*, 146: 1333-1345.
- Li, J. and C.Y. Li. 2019. Seventy-year major research progress in plant hormones by Chinese scholars (in Chinese). *Sci. Sin. Vita.*, 49: 1227-1281.
- Li, Y., C. Zhou, X. Yan, J. Zhang and J. Xu. 2016. Simultaneous analysis of ten phytohormones in *Sargassum horneri* by high-performance liquid chromatography with electrospray ionization tandem mass spectrometry. *J. Sep. Sci.*, 39: 1804-1813.
- Liu, X., Y. Zhu, H. Zhai, H. Cai, W. Ji, X. Luo, J. Li and X. Bai. 2021. AtPP2CG1, a protein phosphatase 2C, positively regulates salt tolerance of Arabidopsis in abscisic acid-dependent manner. *Biochem. Biophys. Res. Co.*, 422: 710-715.
- Lv, S., Y. Zhang, C. Li, Z. Liu, N. Yang, L. Pan, J. Wu, J. Wang, J. Yang, Y. Lv, Y. Zhang, W. Jiang, X. She and G. Wang. 2018. Strigolactone-triggered stomatal closure requires hydrogen peroxide synthesis and nitric oxide production in an abscisic acid-independent manner. *New Phytol.*, 217: 290-304.
- Ma, W.Q., H.Y. Wang, W.J. Zhang, S. Wang, X.F. Wan, C.Z. Kang and L.P. Guo. 2021. Effects of ecological factors on shape and ginsenoside of *Panax ginseng*. *Chin. Mat. Med.*, 46(8): 1920-1926.
- Marro, N., J. Lidoy, M.Á. Chico, C. Rial, J. García, R.M. Varela, F.A. Macías, M.J. Pozo, M. Janoušková and J.A. López-Ráez. 2022. Strigolactones: New players in the nitrogen-phosphorus signaling interplay. *Plant Cell Environ.*, 45: 512-527.
- Merlot, S., F. Gosti, D. Guerrier, A. Vavasseur and J. Giraudat. 2001. The ABI1 and ABI2 protein phosphatases 2C act in a negative feedback regulatory loop of the abscisic acid signaling pathway. *Plant J.*, 25: 295-303.
- Miao, J. X.F. Li, X.B. Li, W.C. Tan, A.Q. You, S.J. Wu, Y.J. Tao, C. Chen, J. Wang, D.P. Zhang, Z.Y. Gong, C.D. Yi, Z.F. Yang, M.H. Gu, G.H. Liang and Y. Zhou. 2020. OsPP2C09, a negative regulatory factor in abscisic acid

- signaling, plays an essential role in balancing plant growth and drought tolerance in rice. *New Phytol.*, 227: 1417-1433.
- Ondzighi, C.A., S. Chakraborty and J.M. Harris. 2016. Environmental nitrate stimulates abscisic acid accumulation in *Arabidopsis* root tips by releasing it from inactive stores. *Plant Cell*, 28: 729-745.
- PriTal, O., L. Shaar-Moshe, G. Wiseglass, Z. Peleg and A. Mosquna. 2009. Non-redundant functions of the dimeric ABA receptor BdPYL1 in the grass *Brachypodium*. *Plant J.*, 92: 774-786.
- Ribot, C., Y. Wang and Y. Poirier. 2008. Expression analyses of three members of the AtPHO1 family reveal differential interactions between signaling pathways involved in phosphate deficiency and the responses to auxin, cytokinin, and abscisic acid. *Planta*, 227: 1025-1036.
- Rigoulot, S.B., H.E. Petzold, S.P. Williams, A.M. Brunner and E.P. Beers. 2019. *Populus trichocarpa* clade A PP2C protein phosphatases: their stress-induced expression patterns, interactions in core abscisic acid signaling, and potential for regulation of growth and development. *Plant Mol. Biol.*, 100: 303-317.
- Rovira, A., M. Sentandreu, A. Nagatani, P. Leivar and E. Monte. 2021. The sequential action of MIDA9/PP2C.D1, PP2C.D2, and PP2C.D5 is necessary to form and maintain the hook after germination in the dark. *Front. Plant Sci.*, 12: 636098.
- Schmittgen, T.D. and K.J. Livak. 2008. Analyzing real-time PCR data by the comparative CT method. *Nat. Protoc.*, 3: 1101-1108.
- Schweighofer, A., H. Hirt and I. Meskiene. 2004. Plant PP2C phosphatases: emerging functions in stress signaling. *Trends Plant Sci.*, 9: 236-243.
- Shi, Y. 2009. Serine/threonine phosphatases: mechanism through structure. *Cell*, 139: 468-484.
- Signora, L., S.I. De, C.H. Foyer and H. Zhang. 2001. ABA plays a central role in mediating the regulatory effects of nitrate on root branching in *Arabidopsis*. *Plant J.*, 28: 655-662.
- Smékalová, V., A. Doskočilová, G. Komis and J. Samaj. 2013. Crosstalk between secondary messengers, hormones and MAPK modules during abiotic stress signaling in plants. *Biotechnol. Adv.*, 32: 2-11.
- Song, L., R. Ding, T. Du, S. Kang, L. Tong, F. Xue and Z. Wei. 2023. Stomatal conductance parameters of tomatoes are regulated by reducing osmotic potential and pre-dawn leaf water potential via increasing ABA under salt stress. *Environ. Exp. Bot.*, 206: 105176.
- Tamura, K., G. Stecher, D. Peterson, A. Filipski and S. Kumar. 2013. MEGA6: Molecular evolutionary genetics analysis Version 6.0. *Mol. Biol. Evol.*, 30: 2725-2729.
- Uhrig, R.G., A.M. Labandera and G.B. Moorhead. 2013. *Arabidopsis* PPP family of serine/threonine protein phosphatases: Many targets but few engines. *Trends Plant Sci.*, 18:505-513.
- Umezawa, T., N. Sugiyama, M. Mizoguchi, S. Hayashi, F. Myouga, S.K. Yamaguchi, Y. Ishihama, T. Hirayama and K. Shinozaki. 2009. Type 2C protein phosphatases directly regulate abscisic acid-activated protein kinases in *Arabidopsis*. *P. Natl. Acad. Sci. USA*, 106: 17588-17593.
- Vallabhaneni, R. and E.T. Wurtzel. 2009. Timing and biosynthetic potential for carotenoid accumulation in genetically diverse germplasm of maize. *Plant Physiol.*, 150: 562-572.
- Wang, G., X. Sun, Z. Guo, D. Joldersma, L. Guo, X. Qiao, K. Qi, C. Gu and S. Zhang. 2021. Genome-wide identification and evolution of the PP2C gene family in eight rosaceae species and expression analysis under stress in *Pyrus bretschneideri*. *Front. Genet.*, 12: 770014.
- Wang, M., P. Zhang, Q. Liu, G. Li, D. Di, G. Xia, H.J. Kronzucker, S. Fang, J. Chu and W. Shi. 2020b. TaANR1-TaBG1 and TaWabi5-TaNRT2s/NARs link ABA metabolism and nitrate acquisition in wheat roots. *Plant Physiol.*, 182: 1440-1453.
- Wang, Z.J., Z.Y. Ren, C.H. Cheng, T. Wang, H.T. Ji, Y. Zhao, Z.P. Deng, L.Y. Zhi, J.J. Lu, X.Y. Wu, S.M. Xu, M.M. Cao, H.T. Zhao, L. Liu, J.K. Zhu and X. Li. 2020a. Counteraction of ABA-mediated inhibition of seed germination and seedling establishment by ABA signaling terminator in *Arabidopsis*. *Mol. Plant*, 13: 1284-1297.
- Wasilewska, A., F. Vlad, C. Sirichandra, Y. Redko, F. Jammes, C. Valon, N.F. Frey and J. Leung. 2008. An update on abscisic acid signaling in plants and more. *Mol. Plant*, 1: 198-217.
- Weiner, J.J., F.C. Peterson, B.F. Volkman and S.R. Cutler. 2010. Structural and functional insights into core ABA signaling. *Curr. Opin. Plant Biol.*, 13: 495-502.
- Welsch, R., F. Wüst, C. Bär, S. AlBabili and P. Beyer. 2008. A third phytoene synthase is devoted to abiotic stress-induced abscisic acid formation in rice and defines functional diversification of phytoene synthase genes. *Plant Physiol.*, 147: 367-380.
- Wen, B., X. Gong, X. Chen, Q. Tan, L. Li and H. Wu. 2018. Transcriptome analysis reveals candidate genes involved in nitrogen deficiency stress in Apples. *J. Plant Physiol.*, 279: 153822.
- Wen, J. and E.A. Zimmer. 1996. Phylogeny and biogeography of *Panax* L. (the ginseng genus, Araliaceae): inferences from ITS sequences of nuclear ribosomal DNA. *Mol. Phylogenet. Evol.*, 6: 167-177.
- Xie, Y., Y. Mao, W. Zhang, D. Lai, Q. Wang and W. Shen. 2006. Reactive oxygen species-dependent nitric oxide production contributes to hydrogen-promoted stomatal closure in *Arabidopsis*. *Plant Physiol.*, 165: 759-773.
- Xing, B., C. Gu, T. Zhang, Q. Zhang, Q. Yu, J. Jiang and G. Liu. 2021. Functional study of BpPP2C1 revealed its role in salt stress in *Betula platyphylla*. *Front. Plant Sci.*, 14: 617635.
- Xiong, L. and J.K. Zhu. 2003. Regulation of abscisic acid biosynthesis. *Plant Physiol.*, 133: 29-36.
- Xue, T., D. Wang, S. Zhang, J. Ehrling, F. Ni, S. Jakab, C. Zheng and Y. Zhong. 2008. Genome-wide and expression analysis of protein phosphatase 2C in rice and *Arabidopsis*. *BMC Genom.*, 9: 550.
- Yang, N.Q., P.S. Chen, Z.W. Tao, N.T. Zhou, X.X. Gong, Z.H. Xu, M. Zhang, D.G. Zhang, B. Chen, Z.X. Tao and Z.J. Yang. 2012. Beneficial effects of ginsenoside-Rg1 on ischemia-induced angiogenesis in diabetic mice. *Acta Bioch. Bioph. Sin.*, 44: 999-1005.
- Yang, Y., M. Gu, J. Chen, R. Zhang, Z. Liu, Y. Shi, D. Liu and L. Wang. 2022. Comparative transcriptomes reveal the mitigation effect of GR24 in alfalfa under drought stress. *J. Plant Growth Regul.*, 42: 3150-3161.
- Yoshida, R., T. Umezawa, T. Mizoguchi, S. Takahashi, F. Takahashi and K. Shinozaki. 2006. The regulatory domain of SRK2E/OST1/SnRK2.6 interacts with ABI1 and integrates abscisic acid (ABA) and osmotic stress signals controlling stomatal closure in *Arabidopsis*. *J. Biol. Chem.*, 281: 5310-5318.
- Yu, X.F., J.P. Han, E. Wang, J. Xiao, R. Hu, G.X. Yang and G.Y. He. 2008. Genome-wide identification and homoeologous expression analysis of PP2C genes in wheat (*Triticum aestivum* L.). *Front. Genet.*, 10: 561.
- Yu, X.F., J.P. Han, L. Li, Q. Zhang, G.X. Yang and G.Y. He. 2020. Wheat PP2C-a10 regulates seed germination and drought tolerance in transgenic *Arabidopsis*. *Plant Cell Rep.*, 39: 635-651.
- Zhai, X., D. Lan, M. Jia, Z. Gan, C. Chen, X. Zeng, J. Chen and W. Kai. 2023. Characterization and expression analysis of key abscisic acid signal transduction genes during kiwifruit development. *Sci. Hort.*, 309: 111672.

- Zhang, B.H., N.Z. Chen, X.J. Peng and S.H. Shen. 2021. Identification of the PP2C gene family in paper mulberry (*Broussonetia papyrifera*) and its roles in the regulation mechanism of the response to cold stress. *Biotechnol. Lett.*, 43: 1089-1102.
- Zhang, Y., W. Xu, Z. Li, X.W. Deng, W. Wu and Y. Xue. 2006. F-box protein DOR functions as a novel inhibitory factor for abscisic acid-induced stomatal closure under drought stress in *Arabidopsis*. *Plant Physiol.*, 148: 2121-2133.
- Zhang, Y., Z. Yakun, C. Jiang, D. Yan, J. Gao, Y. Jiang and C. Chen. 2022. Screening and verification of internal reference genes in *Panax ginseng* roots infected by *Ilyonectria robusta* causing ginseng rusty rot by real-time quantitative PCR. *J. Jilin Agri. Uni.*, 3: 1-10.
- Zhou, M., Y. Li, X. Lu, P. He, C. Liang and J. Tian. 2022. Diverse functions of GmNLA1 members in controlling phosphorus homeostasis highlight coordinate response of soybean to nitrogen and phosphorus availability. *Crop J.*, 11(4): 1251-1260.

(Received for publication 5 May 2024)