

## LEAD HARMS SEED GERMINATION AND GROWTH OF *ALBIZIA LEBBECK* (L.) BENTH. AND *PROSOPIS JULIFLORA* (SW.) DC.

**KHURSHID ANWAR<sup>1\*</sup>, AQSA NADEEM<sup>1</sup>, YAMIN BIBI<sup>1</sup>, FARWA IFTIKHAR<sup>1</sup>, NOSHIN ILYAS<sup>1</sup>, GHULAM YASEEN<sup>1</sup> AND MUHAMMAD KABIR<sup>2</sup>**

<sup>1</sup>*Department of Botany, PMAS, Arid Agriculture University, Rawalpindi, Pakistan*

<sup>2</sup>*Department of Biological Sciences, University of Sargodha, Sub-Campus Bhakkar, Pakistan*

<sup>\*</sup>*Corresponding author's email: khurshidbhatti0@gmail.com*

### Abstract

Lead produced significant ( $p<0.05$ ) effects on various growth parameters of *A. lebbeck* and *P. juliflora* i.e., root, shoot, seedling length, number of leaves, leaf area and seedling biomass. The seeds of *A. lebbeck* exhibited germination at 0ppm, 10pm, 20ppm and 30ppm in lab. Lead treatments at control to 10, 20, and 30ppm reduced the percentage of seed germination from 92% to 77% respectively in *P. juliflora*. Number of leaves, leaf area and biomass of both seedlings exhibited significant ( $p<0.05$ ) decrease when treatments of lead were increased from 20 ppm to 30 ppm. Lead treatments of 25, 50 and 75 ppm were also carried out in pot experiment. Lead treatments at 75 ppm demonstrated much reduction in growth as compared with control. The rise in absorption of lead directly effects shoot growth and root length is also reduced. But at 25 ppm shoot growth of *A. lebbeck* was better than control but shoot growth of *P. juliflora* was reduced. The shoot length was much reduced by increase in concentration of lead. The seedlings grown in control soils have better growth as compared with lead treated soils. The increase in concentration of lead decreases the seedling size. The seedlings which were treated with the 25ppm and 50 ppm were almost of same size but 75 ppm caused much reduction in both seedlings growth. Increased concentration of lead reduced the dry weight. General fallouts of different limits showed that lead is much toxic element, which caused much reduction in seed germination and growth of both *A. lebbeck* and *P. juliflora*.

**Key words:** *A. lebbeck*, *P. juliflora*, Lead, Biomass, Germination, Pot conditions.

### Introduction

Heavy metals in atmosphere disturb the morphological and physiological parameters of vegetation. The primary source of pollution is combustion of fossils fuel (Amin *et al.*, 2013). The green vegetables and plants take up heavy metals. After the uptake these toxic elements are transferred to different tissues of plants, increase in translocation of heavy metals enhances quantity of these metals in plants more than that of soil (Khan *et al.*, 2010). Heavy metals are not individually harmful for plants, if they are preceded in food chain they are also hazards for humans and other living organisms (Kumar *et al.*, 2013).

Lead is heavy metal with atomic number of 82, having atomic mass 207.2 amu and have low melting point. Lead (Pb) is one of the most harmful pollutants present in soil, taken in plants by roots. Lead pollution directly disturbs ecosystem by entering into the food chain of organisms. The major sources of Pb are primarily automobile emission, paints chips, fertilizers, Pb-acid batteries, insecticides, pesticides and other industrial wastes (Barrutia *et al.*, 2010). Pb is taken in plants by roots and sometimes by leaves (Amari *et al.*, 2017). Pb effects plants and food chain (He *et al.*, 2004; Belkhadi *et al.*, 2010; Gallego *et al.*, 2012; Saidi *et al.*, 2014). Plants grown on lead polluted soils like roadsides have decreased growth (Asati *et al.*, 2016). With increase uptake of lead by plants the ultra-structures of tissues, cells, chloroplast, mitochondria, nucleus, cell wall and cell membrane are altered in result different physiological processes of plants like photosynthesis, respiration and protein synthesis are effected which eventually lowers plant growth (Salazar & Pignata, 2014). High quantity of lead and cadmium effects metabolism rate of young *Phoenix dactylifera* (Abbas *et al.*, 2016). The High concentrations of

chromium and lead are also found to be stress full to the mustard seedling by decreasing growth (Al-Mahmud *et al.*, 2018). The content of chlorophyll decreases with increase in Pb (Tandon & Srivastva, 2015). Lead directly effects pigments, decreasing rate of photosynthesis i.e. eventually decreased plant growth in Pea plant (Pichhode & Nikhil, 2017). The activities of enzymes like superoxidase dismutase and peroxidases increases with increase in Pb while catalases activity is decreased eventually effect metabolism in plants (Yang *et al.*, 2011). To avoid phytotoxicity of different heavy metals like lead, plant have adopted mechanism like production of phytochelatins (PCs) and metallothioneins (MTs) (Lin *et al.*, 2012; Saidi *et al.*, 2014; Qing *et al.*, 2015). Selenium can reduce uptake of lead and cadmium (Wu *et al.*, 2016). The reed plants can uptake lead so they can be used to get rid of Pb from water (Khalid *et al.*, 2017).

*Albizia lebbeck* L., belongs to family Mimosaceae. *A. lebbeck* is native species to deciduous and semi deciduous forests in Asia. It is a leguminous tree, 10-20 m tall mostly and has widespread canopy. It can also be found in moist semi-evergreen or evergreen forests usually occurring scattered. It is also a good source of timber for furniture and other woodwork (Mazhar *et al.*, 2015).

*Prosopis juliflora* (Sw.) DC. belongs to family Fabaceae. It is very popular shrub, because it is involved in transmission of malaria especially during dry season. It is local specie of Mexico and Australia. This tree has height up to 12.5 meters (40ft) and its trunk is about 1.2 meters (3.9ft). It is used as forage and fodder for animals, wood is used in furniture. It is also used for environmental management. The aim of work was to investigate adverse effects of lead on different growth parameters of *A. lebbeck* and *P. juliflora*.

## Materials and Methods

The experiment was conducted to know effects of lead on different growth parameters of *Albizia lebbeck* (L.) Benth. and *Prosopis juliflora* (Sw.) DC. Healthy seeds of *Albizia lebbeck* and *Prosopis juliflora* were obtained from National Agriculture Research Center (NARC) Islamabad. For germination, seeds were purified and sterilized with sodium hypochlorite (0.5%) solution for 20 minutes. These seeds were then washed with distilled water. The seeds were water-logged for 12 hours. Petridishes were autoclaved to avoid pathogen attack. The seeds were then placed in petridishes with double layer of filter paper. The medium of germination for seeds was filter paper (Watman # 42). The control treatment received 5 ml of distilled water. The other petri dishes received 5 ml of solution of each lead treatment i.e. 10, 20, 30 ppm respectively. The control petri dish was kept moist by giving distill water when needed. The treated petri dishes were also kept moist by applying treatments when needed. The dishes were kept in the dark portion of the laboratory at room temperature (25 degree centigrade). Germination counts were noted day-to-day for a period of ten days. The pot experiment was laid out as completely randomized designed (CRD) with three treatments of lead contaminated soils in replicates. Lead treated soils and control soil were filled in 12 earthen pots having size (6 cm x 8 cm) length and breadth respectively. Each pot was filled with 50 gram of sample soil. The pots were given water and next day seeds were sown so that heat in soil was lost. Mechanical rubbing and cutting of seeds was done as pretreatment. Water was given daily as needs of seeds. The seed germination data was updated after four days. The reshuffling of pots was done every week to avoid effects of light or any other environmental factor. After two weeks germinated plants were shifted in disposable plastic glass pots. Each pot had one seedling. The pot soils samples received 25ppm, 50ppm and 75 ppm lead treatment respectively. The plants were collected after a month. Growth parameters like root length, shoot length, plant size was measured by meter-rule method with iron scale. Germination percentage and plant seedling growth parameters were determined (Iqbal & Rahmati, 1992). Data was analyzed by using analysis of variance (ANOVA) to check difference between control and treatments and Duncan's multiple range test ( $p<0.05$ ) using the SPSS Statistics (20.0 version) software.

## Results and Discussion

Overall it was observed that lead has adverse effects on different growth parameters of both *Albizia lebbeck* and *Prosopis juliflora*. The results presented that the seed germination, root growth, shoot growth, seedling growth, and dry weight of *A. lebbeck* were reduced in concentrations (10, 20, 30, 50 and 75 ppm) of lead as compared to control (Tables 1.1 & 1.2). Different growth parameters of *P. juliflora* were observed to reduce in lead concentrations (10, 20, 30, 50 and 75 ppm) as compared to control (Tables 2.1 & 2.2). Plants exposed to lead, even at low levels, growth reduction occurs (Kopittke *et al.*, 2010).

Lead directly prevents germination of seeds, growth of seedlings, length of parameters like root/shoot, tolerance index and biomass of root and shoot (Mishra & Tripathi, 2008). Same results are also observed in other

studies at various lead concentrations all vegetative parts showed negative growth effects in *Pisum Sativum*, in *Zea mays* (Çimrin *et al.*, 2007). The effects of lead are less significant ( $p<0.05$ ) on the seed germination of the *A. lebbeck*. Lead treatments at 10, 20, and 30ppm reduced the percentage of seed germination from 92 % to 67% (Table 1.1). Almost more than eighty percent seeds were germinated in all concentrations of lead. But there was much reduction in seed germination at 30ppm lead concentration (Table 1.1). Seeds of *P. juliflora* also showed significant ( $p>0.05$ ) reduction in seed germination as compared to control. Lead treatments at control to 10, 20, and 30ppm reduced the percentage of seed germination from 92 % to 77% respectively (Table 2.1). Germination is strongly inhibited by very low concentrations of Pb. Loss of seed germination due to lead is also reported in *Hordeum vulgare*, *Elsholtzia argyi*, *Spartina alterniflora*, *Pinus halepensis*, *Oryza sativa*, and *Zea may* (Islam & Hoque, 2014). The increase in concentrations of lead directly effects shoot growth. Shoot length of *A. lebbeck* was reduced from 4.78 cm to 3.35 cm with comparing control at 30 ppm treatment, respectively (Table 1.1). In lab. conditions shoot length of *P. juliflora* was reduced from 6.82 cm to 4.70 cm in control to 30 ppm treatment, respectively (Table 2.1). Even at low concentrations, lead effects all process of plant along with shoot growth in wheat (Li *et al.*, 2020). The growth of root was reduced as the quantity of lead increased. Root length of *A. lebbeck* was reduced from 2.15 cm to 1.75 cm in control to 30 ppm treatment, respectively (Table 1.1). Root elongation of *P. juliflora* was reduced from 2.95 cm to 1.85 cm in control to 30 ppm treatment, respectively (Table 2.1). Root elongation is significantly inhibited by lead in Mesquite (*Prosopis* ssp.) (Arias *et al.*, 2010). Dry weight of seedlings which were grown in lead treated soils was similar to control in *A. lebbeck* but dry weight of *P. juliflora* was reduced (Tables 1.1 & 2.1). Plant biomass is reduced by lead exposure (Singh *et al.*, 2010).

In pots, the shoot growth of both *A. lebbeck* and *P. juliflora* was reduced from 21.97 cm to 12.06 and from 21.91 cm to 12.25 cm in control to 75 ppm of lead, respectively (Tables 1.2 & 2.2). The control had better shoot growth than other lead treated soils. The root growth of both *A. lebbeck* and *P. juliflora* were also reduced from 10.67 cm to 9.64 cm and from 10.67 cm to 6.72 cm in control to 75 ppm of lead respectively (Tables 1.2 & 2.2). Lead toxicity is also responsible for bent, swollen, short and broad roots, responsible for increased number of secondary roots per unit root length (Haider *et al.*, 2006). The seedlings grown in control soils were well of size as compared to lead treated soils. Increased concentration of lead decreases the seedling length. The seedlings which were treated with 50ppm and 75ppm caused much reduction in seedling growth. The lead treatments 25, 50 and 75 ppm showed significantly reduce in dry weight. Same negative results of lead on different growth parameters are observed in *Pisum sativum* (Kevresan *et al.*, 2001) and *Cynodon dactylon* (Shua *et al.*, 2002) in *Lycopersicon esculentum* (Jaja & Odoemena, 2004), *Pomoea aquatic* (Gothberg *et al.*, 2004), *Phaseolus vulgaris* and *Lens culinaris* (Haider *et al.*, 2006).

**Table 1.1. Effects of different lead concentrations on different growth parameters of *A. lebbeck* in Lab. conditions.**

| Treatment of lead (ppm) | Germination (%) | Shoot length (Centimeters) | Root length (Centimeters) | Seedling length (Centimeters) | Dry Wt. of seedling (Milligrams) |
|-------------------------|-----------------|----------------------------|---------------------------|-------------------------------|----------------------------------|
| 0                       | 92.50 ± 2.50ab  | 4.78 ± 0.09b               | 2.15 ± 0.25b              | 6.23 ± 0.24b                  | 8.65 ± 0.15bc                    |
| 10                      | 82.50 ± 2.50a   | 4.59 ± 0.28a               | 2.38 ± 0.10a              | 7.45 ± 0.39a                  | 7.90 ± 0.25c                     |
| 20                      | 72.50 ± 2.50a   | 3.35 ± 0.12a               | 1.95 ± 0.15ab             | 5.25 ± 0.22ab                 | 6.80 ± 0.38b                     |
| 30                      | 67.50 ± 2.50a   | 4.43 ± 0.45ab              | 1.75 ± 0.19a              | 5.88 ± 0.45a                  | 4.08 ± 0.40a                     |

Values followed by the same letters in same column are not significantly different ( $p<0.05$ ) according to Duncan's Multiple Range Test

**Table 1.2. Effects of different lead concentrations on different growth parameters of *A. lebbeck* in pot experiment.**

| Treatment of lead (ppm) | Root length (Centimeters) | Shoot length (Centimeters) | Seedling length (Centimeters) | No. of leaves  | Fresh Wt. of Seedling (Grams) | Dry Wt. of Seedling (Milligrams) |
|-------------------------|---------------------------|----------------------------|-------------------------------|----------------|-------------------------------|----------------------------------|
| 0                       | 10.67 ± 1.53a             | 21.97 ± 1.39b              | 37.46 ± 1.40c                 | 36.00 ± 2.00c  | 1.00 ± 0.10a                  | 376 ± 74.00a                     |
| 25                      | 11.43 ± 1.78a             | 12.06 ± 1.40a              | 30.50 ± 1.24b                 | 23.00 ± 3.00bc | 0.76 ± 0.040a                 | 289.50 ± 70.00a                  |
| 50                      | 9.64 ± 0.25a              | 15.62 ± 0.89a              | 25.05 ± 1.62ab                | 30.00 ± 2.00ab | 0.76 ± 0.095a                 | 271 ± 31.00a                     |
| 75                      | 6.72 ± 0.63a              | 12.25 ± 0.85a              | 19.94 ± 1.66a                 | 15.00 ± 1.00a  | 0.82 ± 0.050a                 | 335 ± 55.00a                     |

Values followed by the same letters in same column are not significantly different ( $p<0.05$ ) according to Duncan's Multiple Range Test

**Table 2.1. Effects of different lead concentrations on different growth parameters of *A. lebbeck* in Lab. conditions.**

| Treatment of lead (ppm) | Germination (%) | Shoot length (Centimeters) | Root length (Centimeters) | Seedling length (Centimeters) | Dry wt. of seedling (Milligrams) |
|-------------------------|-----------------|----------------------------|---------------------------|-------------------------------|----------------------------------|
| 0                       | 92.50 ± 2.50b   | 6.82 ± 0.07b               | 2.95 ± 0.15b              | 9.82 ± 0.28b                  | 8.95 ± 0.05c                     |
| 10                      | 87.50 ± 2.50ab  | 5.28 ± 0.38a               | 2.20 ± 0.20ab             | 7.48 ± 0.59a                  | 8.10 ± 0.20bc                    |
| 20                      | 82.50 ± 2.50ab  | 5.18 ± 0.13a               | 1.85 ± 0.25a              | 7.04 ± 0.26a                  | 7.30 ± 0.40b                     |
| 30                      | 77.50 ± 2.50a   | 4.70 ± 0.35a               | 1.95 ± 0.18a              | 6.60 ± 0.47a                  | 5.00 ± 0.50a                     |

Values followed by the same letters in same column are not significantly different ( $p<0.05$ ) according to Duncan's Multiple Range Test

**Table 2.2. Effects of different lead concentrations on different growth parameters of *P. juliflora* in pot experiment.**

| Treatment of lead (ppm) | Root length (Centimeters) | Shoot length (Centimeters) | Seedling length (Centimeters) | No. of leaves  | Fresh wt. of Seedling (Grams) | Dry wt. of seedling (Milligrams) |
|-------------------------|---------------------------|----------------------------|-------------------------------|----------------|-------------------------------|----------------------------------|
| 0                       | 10.67 ± 1.53a             | 21.97 ± 1.39b              | 37.46 ± 1.40c                 | 36.00 ± 2.00c  | 1.00 ± 0.10a                  | 376 ± 74.00a                     |
| 25                      | 11.43 ± 1.78a             | 12.06 ± 1.40a              | 30.50 ± 1.24b                 | 23.00 ± 3.00bc | 0.76 ± 0.040a                 | 289.50 ± 70.00a                  |
| 50                      | 9.64 ± 0.25a              | 15.62 ± 0.89a              | 25.05 ± 1.62ab                | 30.00 ± 2.00ab | 0.76 ± 0.095a                 | 271 ± 31.00a                     |
| 75                      | 6.72 ± 0.63a              | 12.25 ± 0.85a              | 19.94 ± 1.66a                 | 15.00 ± 1.00a  | 0.82 ± 0.050a                 | 335 ± 55.00a                     |

Values followed by the same letters in same column are not significantly different ( $p<0.05$ ) according to Duncan's Multiple Range Test

In conclusion heavy metals, typically lead is not important element to plants, but it has ability to accumulate within various plant parts and harms all physiological developments. Lead presence in soil prevents germination of seeds. Atmospheric lead harms aerial parts of vegetation. Therefore, the Government and various environmental agencies are requested to control, heavy metals pollution particularly lead.

#### References

- Abbas, M.H., J.D. Neama and K. Al-Jabary. 2016. Biochemical responses to cadmium and lead stresses in date palm *Phoenix dactylifera* L. plants. *Adv. Agri. & Bot.*, 83: 92-110.
- Al-Mahmud, J., M. Hasanuzzaman, K. Nahar, M.B. Bhuyan and M. Fujita. 2018. Insights into citric acid-induced cadmium tolerance and phytoremediation in *Brassica juncea* L. coordinated functions of metal chelation, antioxidant defense and glyoxalase systems. *Ecotoxicol. & Environ. Safety*, 147: 990-1001.
- Amari, T., T. Ghnaya and C. Abdelly. 2017. Nickel, cadmium and lead phytotoxicity and potential of halophytic plants in heavy metal extraction. *S. Afr. J. Bot.*, 111: 99-110.
- Amin, H., B. Arain, F. Amin and M. Surhio. 2013. Phytotoxicity of chromium on germination, growth and biochemical attributes of *Hibiscus esculentus* L. *Amer. J. Plant Sci.*, 4: 2431-2439.
- Arias, J.A., J.R. Peralta-Videa, J.T. Ellzey, M. Ren, M.N. Viveros and J.L. Gardea-Torresdey. 2010. Effects of *Glomusdeserticola* inoculation on Prosopis: enhancing chromium and lead uptake and translocation as confirmed by X-ray mapping, ICP-OES and TEM techniques. *Environ. & Exp. Bot.*, 682: 139-148.
- Asati, A., M. Pichhode and K. Nikhil. 2016. Effect of heavy metals on plants: an overview. *Int. J. Appl. Innov. Eng. Manag.*, 5: 2319-4847.
- Barrutia, O., C. Garbisu, J. Hernández-Allica, J.I. García-Plazaola and J.M. Becerril. 2010. Differences in EDTA-assisted metal phytoextraction between metallicolous and non-metallicolous accessions of *Rumex acetosa* L. *Environ. Pollut.*, 1585: 1710-1715.

- Belkhadi, A., H. Hediji, Z. Abbes, I. Nouairi, Z. Barhoumi, M. Zarrouk and W. Djebali. 2010. Effects of exogenous salicylic acid pre-treatment on cadmium toxicity and leaf lipid content in *Linum usitatissimum* L. *Ecotox. & Environ. Saf.*, 735: 1004-1011.
- Cimrin, K.M., M. Turan and B. Kapur. 2007. Effect of elemental sulphur on heavy metals solubility and remediation by plants in calcareous soils. *Fresenius Environ. Bull.*, 169A-1113.
- Gallego, S.M., L.B. Pena, R.A. Barcia and C.E. Azpilicueta. 2012. Iannone MF, Rosales EP, Zawoznik MS, Groppa MD, Benavides MP Unravelling cadmium toxicity and tolerance in plants: Insight into regulatory mechanisms. *Environ. Exp. Bot.*, 83: 33-46.
- Gothberg, A., M. Greger, K. Holm and B.E. Bengtsson. 2004. Influence of nutrient levels on uptake and effects of mercury, cadmium, and lead in water spinach. *J. Environ. Qual.*, 33(4): 1247-1255.
- Haider, S., S. Kanwal, F.U. Ddin and R. Azmat. 2006. Phytotoxicity of Pb II. changes in chlorophyll absorption spectrum due to toxic metal Pb stress on *Phaseolus mungo* and *Lens culinaris*. *Pak. J. Biol. Sci.*, 9(11): 2062-2068.
- He, P.P., X.Z. Lv and G.Y. Wang. 2004. Effects of Se and Zn supplementation on the antagonism against Pb and Cd in vegetables. *Environ. Int.*, 302: 167-172.
- Iqbal, M.Z. and K. Rahmati. 1992. Tolerance of *Albizia lebbeck* to Cu and Fe application. *Ekológia, CSFR*, 114: 427-430.
- Islam, M.S. and M.F. Hoque. 2014. Concentrations of heavy metals in vegetables around the industrial area of Dhaka city, Bangladesh and health risk assessment. *Int. Food Res. J.*, 216.
- Jaja, E.T. and C.S.I. Odoemena. 2004. Effect of Pb, Cu and Fe compounds on the germination and early seedling growth of tomato varieties. *J. App. Sci. & Environ. Manag.*, 82: 51-53.
- Kevresan, S.N. Petrovic, M. Popovic and J. Kandrac. 2001. Nitrogen and protein metabolism in young pea plants as affected by different concentrations of nickel, cadmium, lead, and molybdenum. *J. Plant Nutr.*, 24: 1633-1644.
- Khalid, S., M. Shahid, N.K. Niazi, B. Murtaza, I. Bibi and C. Dumat. 2017. A comparison of technologies for remediation of heavy metal contaminated soils. *J. Geochem. Explor.*, 182: 247-268.
- Khan, S., S. Rehman., A.Z. Khan., A.M. Khan and M.T. Shah. 2010. Soil and vegetables enrichment with heavy metals from geological sources in Gilgit, northern Pakistan. *Ecotoxic. & Environ. Saf.*, 73: 1820-1827.
- Kopittke, P.M., F.P. Blamey, C.J. Asher and N.W. Menzies. 2010. Trace metal phytotoxicity in solution culture: a review. *J. Exp. Bot.*, 614: 945-954.
- Kumar, S., M.H Asif, D. Chakrabarty, R.D. Tripathi, R.S. Dubey and P.K. Trivedi. 2013. Expression of a rice Lambda class of glutathione S-transferase in *Arabidopsis* provides tolerance to heavy metal and other abiotic stresses. *J. Hazard. Mat.*, 6: 248-249.
- Li, L., Y. Zhang, J.A. Ippolito, W. Xing, K. Qiu and H. Yang. 2020. Lead smelting effects heavy metal concentrations in soils, wheat, and potentially humans. *Environ. Poll.*, 257: 113641.
- Lin, L., W. Zhou, H. Dai, F. Cao, G. Zhang and F. Wu. 2012. Selenium reduces cadmium uptake and mitigates cadmium toxicity in rice. *J. Hazar. Mater.*, 235: 343-351.
- Mazhar, H., M. Wajid, M.S. Aslam, M.S. Ahmad and A.S. Mamat. 2015. Ethnomedicinal survey of plants of Thal Desert Choubara district Layyah, Punjab, Pakistan. *Ind. Res. J. Pharm. and Sci.*, 6: 235-250.
- Mishra, V.K. and B.D. Tripathi. 2008. Concurrent removal and accumulation of heavy metals by the three aquatic macrophytes. *Bioreso. Technol.*, 9915: 7091-7097.
- Pichhode, M. and K. Nikhil. 2017. Carbon sequestration by different tree species at Malanjkhand, district Balaghat, Madhya Pradesh, *Ind. Amer. J. Eng. Res.*, 6(10): 6-13.
- Qing, X., X. Zhao, C. Hu, P. Wang, Y. Zhang, X. Zhang and C. Qu. 2015. Selenium alleviates chromium toxicity by preventing oxidative stress in cabbage *Brassica campestris* L. ssp. Pekinensis leaves. *Ecotoxic. & Environ. Saf.*, 114: 179-189.
- Saidi, I., Y. Chtourou and W. Djebali. 2014. Selenium alleviates cadmium toxicity by preventing oxidative stress in sunflower *Helianthus annuus* seedlings. *J. Plant Physiol.*, 1715: 85-91.
- Salazar, M.J. and M.L. Pignata. 2014. Lead accumulation in plants grown in polluted soils. Screening of native species for phytoremediation. *J. Geochem. Explor.*, 137: 29-36.
- Shua, W.S., Z.H. Yeb, C.Y. Lana, Z.Q. Zhanga and M.H. Wongb. 2002. Lead, zinc and copper accumulation and tolerance in populations of *Paspalum distichum* and *Cynodon dactylon*. *Environ. Pollut.*, 120: 445-453. <http://dx.doi.org/10.1016/S0269-74910200110-0>
- Singh, A., R.K. Sharma, M. Agrawal and F.M. Marshall. 2010. Risk assessment of heavy metal toxicity through contaminated vegetables from waste water irrigated area of Varanasi, India. *Trop. Ecol.*, 512: 375-387.
- Tandon, P.K. and P. Srivastava. 2015. Cadmium and lead induced biochemical effects in bean (*Dolichos lablab*. L) plants. *Res. Environ. Life Sci.*, 8(1): 129-130.
- Wu, Z., X. Yin, G.S. Bañuelos, Z.Q. Lin, Y. Liu, M. Li and L. Yuan. 2016. Indications of selenium protection against cadmium and lead toxicity in oilseed rape *Brassica napus* L. *Frontiers in Plant Sci.*, 7: 1875.
- Yang, Y., Z. Nan, Z. Zhao, S. Wang, Z. Wang and X. Wang. 2011. Chemical fractionations and bioavailability of cadmium and zinc to cole (*Brassica campestris* L.) grown in the multi-metals contaminated oasis soil, northwest of China. *J. Environ. Sci.*, 23(2): 275-281.

(Received for publication 11 April 2020)