

INFLUENCE OF SALICYLIC ACID SEED PRIMING ON GROWTH AND SOME BIOCHEMICAL ATTRIBUTES IN WHEAT GROWN UNDER SALINE CONDITIONS

MOHSINA HAMID¹, M. YASIN ASHRAF*, KHALIL-UR-REHMAN¹ AND M. ARASHAD²

Plant Stress Physiology and Biochemistry Lab, NIAB, Faisalabad, Pakistan

¹*Department of Biochemistry, University of Agriculture, Faisalabad, Pakistan.*

²*Cholistan Institute of Desert Studies, The Islamiya University, Bahawalpur, Pakistan.*

Abstract

A sand culture experiment was conducted to study the effect of salicylic acid seed priming on growth and some biochemical attributes in wheat (*Triticum aestivum* L.) grown under saline conditions. Wheat seeds of cv. Inqlab and S-24 were soaked in water and 100 mg L⁻¹ salicylic acid solution for 24 hours and sown in sand salinized with 0, 50 or 100 mM NaCl. Pots were irrigated with $\frac{1}{4}$ strength Hoagland's nutrient solution. Fourteen days old seedlings were harvested and growth parameters (shoot and root length, and shoot and root dry weights) were recorded and chlorophyll *a* and *b* contents and soluble sugars (reducing and non-reducing) in the leaves were estimated. Salt stress significantly reduced all growth parameters. However, salicylic acid treatment alleviated the adverse effect of salinity on growth. Salinity decreased the chlorophyll *a* and *b* content and chlorophyll *a/b* ratio in both the lines, but reduction in chlorophyll *a/b* ratio was lower in salt tolerant wheat line S-24, which could be a useful marker for selection of salt tolerant wheat. Salinity (NaCl) stress considerably increased the accumulation of reducing, non-reducing and total soluble sugars in the leaves of 14 days old wheat seedlings of both cultivars. Salt tolerant line S-24 accumulated higher sugar content which could also be a useful marker for salt tolerance in wheat.

Introduction

Salinity in the growth medium causes significant reduction in plant growth parameters like leaf area, leaf length, root and shoot dry weight (Hamada & Al-Hakimi, 2001; Ashrafuzzamana *et al.*, 2002), decrease carotenoid and induce reduction in chlorophyll and photosynthetic activity. Similarly, saline conditions decrease the activity of ribulose 1, 5-bisphosphate carboxylase (Rubisco) which results in reduction in formation of carbohydrates (El-Shihaby *et al.*, 2002).

One of the approaches to cope with the problem is to find out some bio-chemicals that might mitigate the effect of salinity on plants and enable them to grow normally and produce reasonable yields. Salicylic acid and related compounds have been reported to induce significant effects in various biological aspects in plants (Gill *et al.*, 2001; Wang & Li, 2006). These compounds inhibit certain processes and enhance others (Raskin, 1992). Different levels of acetylsalicylic acid function as anti-transpirant and inhibit the opening of stomata (Larque-Saavedra, 1979). Salicylic acid also reverses the closure of stomata caused by abscisic acid (Rai *et al.*, 1986). Exogenous application of salicylic acid improves the yield of crops (Singh & Kaur, 1980; Arfan *et al.*, 2007). Salicylic acid treatment retards ethylene synthesis; stimulates photosynthetic machinery and increase the content of chlorophyll (Leslie & Romani, 1988). Recently, it has been recognized that salicylic acid is required in the signal transduction for inducing systemic acquired resistance against pathogenic infections (Metraux *et al.*, 1990; Gaffney *et al.*, 1993; Vernooij *et al.*, 1994).

*Corresponding author E-mail: niabmyashraf@yahoo.co.uk

The objective of this study was to investigate whether salicylic acid could be used to mitigate the adverse effects of salinity on wheat seedlings and to identify the biochemical criteria conferring the tolerance to salinity stress in wheat.

Materials and Methods

A sand culture experiment was conducted in plastic pots in net house at NIAB, Faisalabad, Pakistan. Homogenous lots of healthy seeds of two wheat varieties, Inqlab and S-24 were sterilized with Sodium hypochloride solution (5%) for five minutes, washed thoroughly with distilled water before use. Before sowing seeds were soaked in Salicylic acid solution (100 mg L^{-1}) and distilled water for 24 hours. The silica sand thoroughly washed with distilled water and dried sand was filled in plastic pots (15 cm inner diam., and 15 cm deep) maintaining the weight of sand at 2 kg per pot.

Three levels of salinity (0, 50 and 100 mM NaCl) were developed in pots filled with sand before sowing. The seeds of two wheat varieties were soaked for 24 h in 0 and 100 mg L⁻¹ Salicylic acid. Ten seeds per pot containing 1/4 strength Hoagland nutrient solution (Hoagland and Arnon, 1950) and sand were sown. All the treatments were arranged in a completely randomized block design. The wheat seedlings were allowed to grow for 14 days. The roots and shoots of seedlings were collected and rinsed with distilled water. The seedlings were immediately dried in tissue paper. The fresh leaves were used for the estimation of chlorophyll, reducing, non-reducing and total soluble sugars. The dry weights of roots and shoots were recorded after drying in an oven at $70 \pm 2^\circ\text{C}$ for 72 h.

Chlorophyll was determined according to the method of Arnon (1949) and Davies (1976). Total soluble, reducing and non-reducing sugars were extracted in 0.2 M phosphate buffer with pH 7 and analyzed according to the method of Razi *et al.*, (1985).

All the collected data were subjected to analysis of variance and LSD worked out for comparison of means (Steel *et al.*, 1997).

Results

Salinity affected adversely the growth parameters such as shoot and root length, dry weight of shoot and root. In wheat cultivar Inqlab shoot length was severely reduced at 100 mM NaCl salinity (Table 1). Wheat cultivar S-24 performed better than Inqlab at the highest salinity level. Wheat cultivar Inqlab showed 49% reduction in shoot length at the highest salinity level which was 33.33% in S-24. Application of salicylic acid stimulated growth in both the cultivars under control and salinity stress. Cultivar S-24 showed more increase in shoot length than that of Inqlab.

The growth of root was also adversely affected by salinity in both the wheat cultivars (Table 1). In cultivar Inqlab, root length reduction was more prominent than that of S-24, under 50 and 100 mM NaCl. The salicylic acid treatment increased root length in both wheat cultivars. Salinity adversely affected the shoot and root dry weight of the two cultivar used in this study (Table 1). However, application of salicylic acid enhanced the shoot and root dry weight.

Root dry weight was also adversely affected by salinity (Table 1). Under control dry weight of root was 0.134 and 0.130 g plant⁻¹ in Inqlab and S-24, respectively. The dry weight of root was reduced by 18.7, 44.0% in Inqlab and 12.3 and 30.8% in S-24 under 50 and 100 mM NaCl stress, respectively. Salicylic acid treatment increased root dry

weight by 11.9, 18.4 and 33.7% in Inqlab and 7.7, 7.9 and 12.1% in S-24 under control and 50, 100 mM NaCl stress, respectively.

Salinity had an adverse effect on chlorophyll in wheat seedlings (Table 2). Both the salinity levels adversely affected Chlorophyll contents in both the wheat varieties however, reduction was more pronounced in Inqlab. Application of salicylic acid improved the chlorophyll 'a' content of wheat seedlings of both the cultivars under normal and saline conditions. Wheat variety S-24 performed better than that of Inqlab.

Like chlorophyll 'a' and 'b' content of wheat seedlings was also adversely affected by salinity (Table 2), the reduction in chlorophyll 'b' was more than that in chlorophyll 'a'. Severe reduction in chlorophyll 'b' was found at 100 mM NaCl in both the wheat cultivars. Salicylic acid application increased the content of chlorophyll 'b' in wheat seedlings of the two cultivars under study. Wheat line S-24 maintained higher chlorophyll b than Inqlab.

An interesting feature of chlorophyll 'a' and 'b' under stress is the shift of chlorophyll a/b ratio and results indicated that the chlorophyll a/b ratio increased with increase in salinity in both the cultivars. The increase in chlorophyll a/b ratio was more in Inqlab than S-24. Application of salicylic acid reduced the chlorophyll a/b ratio under normal as well as saline conditions in both the wheat varieties except in case of Inqlab at 100 mM NaCl where it significantly increased (Table 2).

Salinity affected significantly the non-reducing sugars in wheat seedlings (Table 3). Non-reducing sugars increased due salinity in both the wheat cultivars however, S-24 was successful in maintaining higher concentrations of non-reducing sugars than that of Inqlab (Table 3). Application of salicylic acid further increased the concentration of non-reducing sugars in seedlings of both the varieties under normal and salt stressed conditions.

Reducing sugars also influenced by salinity in both the wheat cultivars (Table 3). Reducing sugars significantly enhanced by in both the wheat cultivars, however, cultivar S-24 maintained high reducing sugars than Inqlab. Salicylic acid treatment further increased the concentration of reducing sugars in the both the wheat cultivars included in this study.

Salinity significantly affected the concentration of total soluble sugars in seedlings of the two cultivars (Table 3). Total sugars increased with increase in salinity in both the wheat cultivars and application of Salicylic acid was effective in further increasing the total sugars. Wheat cultivar S-24 maintained higher concentration of sugars than Inqlab.

Discussions

The results of the present study show that salinity reduced the growth and disturbed the plant metabolism. Seed priming with salicylic acid was beneficial in alleviating the adverse effects of salinity by enhancing growth, chlorophyll contents and sugars necessary for osmotic adjustment under adverse environmental conditions. Hamada & Al-Hakimi (2001) observed that soaking of wheat seed in salicylic acid before sowing was effective in reducing the stress effect of salinity on growth. Gutierrez *et al.*, (1998) reported increase in the growth of shoots and roots of soybean plants in response to salicylic acid treatment. Khodary (2004) observed that salicylic acid treated maize plants exhibited an increase in tolerance to salinity reflected in growth parameters like length, fresh and dry weight of shoot and root.

Table 1. Effect of Salicylic acid on growth characteristics of wheat seedlings of two varieties under NaCl stress.

Treatment		Shoot length cm		Root length cm		Shoot dry weight g plant ⁻¹		Root dry weight g plant ⁻¹		Shoot/root ratio	
SA mg L ⁻¹	NaCl mM	Inq.	S-24	Inq.	S-24	Inq.	S-24	Inq.	S-24	Inq.	S-24
0	0	29.6	25.5	34.3	31.9	0.264	0.260	0.134	0.130	1.970	2.00
	50	22.4	21.5	25.5	27.9	0.204	0.236	0.109	0.114	1.872	2.070
	100	15.1	17.0	17.2	19.7	0.105	0.130	0.065	0.09	1.615	1.444
	0	32.2	29.6	42.1	38.9	0.300	0.283	0.150	0.140	2.000	2.021
	50	23.8	25.7	30.8	33.0	0.229	0.253	0.129	0.123	1.775	2.056
	100	17.4	22.6	20.2	24.6	0.125	0.179	0.100	0.112	1.250	1.598
LSD (p>0.05)		4.62	5.04	0.053	0.009	0.009	0.052				

SA = Salicylic acid; Inq. = Inqlab; LSD = Least significant difference

Table 2. Effect of Salicylic acid on chlorophyll content of wheat seedlings of two varieties under NaCl stress.

Treatment		Chlorophyll 'a' mg g ⁻¹ F.Wt.		Chlorophyll b mg g ⁻¹ F.Wt.		a/b ratio	
SA mg L ⁻¹	NaCl mM	Inq.	S-24	Inq.	S-24	Inq.	S-24
0	0	0.874	0.885	0.539	0.564	1.621	1.569
	50	0.748	0.780	0.424	0.480	1.764	1.625
	100	0.549	0.656	0.278	0.345	1.974	1.901
	0	0.889	0.895	0.575	0.655	1.546	1.366
	50	0.765	0.798	0.560	0.640	1.366	1.247
	100	0.650	0.712	0.202	0.414	3.218	1.719
LSD (p>0.05)		0.119	0.075	0.075	0.075	ND	ND

SA = Salicylic acid; Inq. = Inqlab; LSD = Least significant difference; ND = Not determined

Table 3. Effect of Salicylic acid on non-reducing sugars, reducing sugars and total soluble sugars content of wheat seedlings of two varieties under NaCl stress.

Treatment		Non reducing sugar mg g ⁻¹ F.Wt.		Reducing sugars mg g ⁻¹ F.Wt.		Total soluble sugars mg g ⁻¹ F.Wt.	
SA mg L ⁻¹	NaCl mM	Inq.	S-24	Inq.	S-24	Inq.	S-24
0	0	1.626	1.618	0.453	0.448	2.079	2.066
	50	1.697	1.799	0.552	0.741	2.249	2.540
	100	1.706	2.018	0.653	0.914	2.359	2.932
	0	1.675	1.705	0.616	0.775	2.291	2.480
	50	2.014	2.570	0.745	0.880	2.759	3.450
	100	2.409	2.755	0.934	0.982	3.343	3.737
LSD (p>0.05)		0.465	0.131	0.131	0.315		

SA = Salicylic acid; Inq. = Inqlab; LSD = Least significant difference

Growth reduction due to salinity is well documented (Khan *et al.*, 1990; Saqib *et al.*, 1999; Ashraf & Harris, 2004; Ashraf *et al.*, 2005) and is mainly attributed to water deficit due to lowered water potential in the root zone, nutritional imbalance and specific ion toxicity arising from higher concentration of Na⁺ and Cl⁻ (Khan & Ashraf, 1988). A secondary effect of high concentration of Na⁺ and Cl⁻ in the rooting medium is the suppression of uptake of essential nutrients such as K⁺, Ca²⁺ and NO₃⁻ etc., (Ashraf, 2004; Gorham & Wyn Jones, 1993). Different physiological traits such as selectivity for potassium, exclusion of Na⁺ and Cl⁻ ions, osmotic adjustments by accumulation of organic solutes have all been related to salt tolerance of crop plants (Khan *et al.*, 1995; Ashraf *et al.*, 2002). There is also the possibility that NaCl treated plants utilize energy for the osmotic adjustment process (Ashraf *et al.*, 2006) at the expense of growth.

Regarding salicylic acid, people are not the only ones to benefit from Salicylic acid, the active ingredient is aspirin. Research has shown that spraying this naturally occurring compound on to some plants triggers natural defenses that keep harmful fungi, bacteria and viruses at bay. Plant scientists first encountered the phenomenon called systemic acquired resistance (SAR) in 1930s. Plants make Salicylic acid, particularly after encountering a pathogen, and use it as a key regulator or SAR expression of defense genes. But only recently have companies begun marketing Salicylic acid and other similar compounds as a way to activate SAR in crops.

Systemic acquired resistance (SAR) is the basic mechanism by which plants protect themselves. Treatment of wheat plants with Salicylic acid increased the level of cell division within the apical meristem of seedling roots which caused an increase in plant growth (Sakhabutdinova *et al.*, 2003). Phytohormones are known to play a key role in plant growth regulation. It was found that salicylic acid treatment caused accumulation of both Abscisic acid and indole acetic acid in wheat seedling. However, the Salicylic acid treatment did not influence cytokinin content.

It is supposed that the protective and growth promoting effects of Salicylic acid are due to phenomenon described above. The Salicylic acid treatment reduces the damaging action of salinity and water deficit on seedling growth and accelerates the restoration of growth processes. Treatment with Salicylic acid diminished the alteration of phytohormones level in wheat seedlings under salinity and water deficit. The Salicylic acid treatment prevented the decrease in Indole acetic acid and cytokinin content completely which reduces stress reduced inhibition of plant growth. The protective Salicylic acid action includes the development of antistress programs and acceleration of normalization of growth processes after removal of stress factors.

From the results presented here it might be concluded that NaCl salinity significantly reduced the growth of wheat seedlings (Shoot was affected more than root as is evident from root/shoot ratio) and chlorophyll content and increased the accumulation of soluble sugars in leaves. Wheat variety S-24 showed better performance than that of Inqlab in all parameters studied in this investigation. The above mentioned two criteria could be useful markers for selecting salt tolerant variety of wheat for breeding program for salt affected areas. Salicylic acid treatment of salt-stressed wheat plants stimulated their salt tolerance in both wheat varieties *via* accelerating their photosynthesis performance and carbohydrate metabolism so it can be utilized for enhancing the salt tolerance potential of wheat varieties and obtaining the economical wheat yield if cultivated on salt-affected soils.

References

Arfan, M., H.R. Athar and M. Ashraf. 2007. Does exogenous application of Salicylic acid through the rooting medium modulate growth and photosynthetic capacity in two differently adapted spring wheat cultivars under salt stress? *J. Plant Physiol.*, 164(6): 685-694.

Arnon, D.T. 1949. Copper enzymes in isolated chloroplasts poly phenoloxidase in *Beta vulgaris*. *Plant Physiol.*, 24: 1-15.

Ashraf, M. 2004. Some important physiological selection criteria for salt tolerance in plants. *Flora*, 199(5): 361-376.

Ashraf, M. and P.J.C. Harris. 2004. Potential biochemical indicators of salinity tolerance in plants. *Plant Sci.*, 166: 3-16.

Ashraf, M.Y. 2007. Variation in nutritional composition and growth performance of some halophytic species grown under saline conditions. *African Journal of Range & Forage Science*, 24: 19-23.

Ashraf, M.Y., N. Azhar and M. Hussain. 2006. Indole acetic (IAA) induced changes in growth, relative water contents and gas exchange attributes of barley (*Hordeum vulgare* L.) grown under water stress conditions. *Plant Growth Regulation*, 50: 85-90.

Ashraf MY, M. Ashraf and G. Sarwar. 2005. Response of Okra (*Hibiscus esculentus*) to drought and salinity stress In: *Vegetables: Growing Environment and Mineral Nutrition*. (Ed.): R. Dris. ISBN: 952-99555-1-0, WFL Publisher (Helsinki, Finland) pp. 166-177.

Ashraf, M.Y., R. Afaf and M.S. Qureshi, G. Sarwar and M.H. Naqvi. 2002. Salinity induced changes in α -amylase and protease activities and associated metabolism in cotton varieties during germination and early seedling growth stages. *Acta Physiologiae Plantarum*, 24(1): 37-44.

Ashrafuzzaman, M., M.A.H. Khan and S.M. Shahidullah. 2002. Vegetative growth of maize (*Zea mays*) as affected by a range of salinity. *Crop Res. Hisar.*, 24: 286-291.

Davies, B.H. 1976. Carotenoids, In: *Chemistry and Biochemistry of Plant Pigments*. (Ed.): T.W. Goodwin. (2nd Edition), Academic Press, London, 2: 38-165.

El-Shihaby, O.A., M.N.N. Alla, M.E. Younis and Z.M. Bastaway. 2002. Effect of kinetin on photosynthetic activity and carbohydrate content in waterlogged or sea-water treated *Vigna sinensis* and *Zea mays* plants. *Plant Biosyst.*, 136: 227-290.

Gaffney, T., L.Friederich, B. Vernooij, D. Negrotto, G. Nye, S. Uknas, E. Ward, H. Kessman and J. Ryals. 1993. Requirement of salicylic acid for the induction of systemic acquired resistance. *Science*, 261: 754-756.

Gill, P.K., A.D. Sharma, P. Singh and S.S. Bhullar. 2001. Effect of various abiotic stresses on the growth, soluble sugars and water relations of sorghum seedling growth in light and darkness. *Bulg. J. Plant Physiol.*, 27: 72-84.

Gorham, J. and R.G. Wyn Jones. 1993. Utilization of *Triticaceae* for improving salt tolerance in wheat. In: *Towards the Rational use of High Salinity Tolerant Plants*. (Eds.): H. Leith and A.A. Massoum. Kluwar Acad. Pub. The Netherlands. pp. 27-33.

Gutierrez-coronada, M.A., C. Trejo-Lopez and A. Larque-Saavedra. 1998 Effects of Salicylic acid on the growth of roots and shoots in soybean. *Plant Physiol. Biochem.*, 36: 653-665.

Hamada, A.M. and A.M.A. Al-Hakimi. 2001. Salicylic acid versus salinity-drought induced stress on wheat seedlings. *Rostl. Vyro.*, 47: 444-450.

Hoagland, D.R. and D.I. Arnon. 1950. The water culture method for growing plants without soil. *Calif. Agric. Exp. Stn. Cir.*, 347, p. 32.

Khan, A.H., M.Y. Ashraf and A.R. Azmi. 1990. Effect of NaCl on growth and nitrogen metabolism of sorghum. *Acta Physiol. Plant.*, 12: 233-238.

Khan, A.H. and M.Y. Ashraf. 1988. Effect of sodium chloride on growth and mineral composition of sorghum. *Acta Physiol. Plant.*, 10: 259-264.

Khan, A.H., M. Y. Ashraf, S.S.M. Naqvi, B. Khanzada and M. Ali. 1995. Growth, ion and solute contents of sorghum grown under NaCl and Na₂SO₄ salinity stress. *Acta Physiol. Plant.*, 17: 261-268.

Khodary, S.E.A. 2004. Effect of salicylic acid on the growth, photosynthesis and carbohydrate metabolism in salt stressed maize plants. *Int. J. Agri. Biol.*, 6:5-8.

Larque-Saavedra, A. 1979. Stomatal closure in response to acetylsalicylic acid treatment. *Z. Pflanzenphysiol.*, 93: 371-375.

Leslie, C.A. and R.J. Romani. 1988. Inhibition of ethylene biosynthesis by salicylic acid. *Plant Physiol.*, 88: 833-837.

Malcolm, C.V. 1993. The potential of halophytes for rehabilitation of degraded land. In: *Productive use of saline land*. (Eds.): N. Davidson and R. Galloway. ACIAR Proc. 42:8-11, Proc. Workshop, Perth, Western Australia.

Metraux, J.P., H. Singer, J. Ryals, E. Ward, M. Wyff-Panz, J. Gaudin, K. Rafehdorf, E. Fehmid, W. Blum and B. Inverardi. 1990. Increase in Salicylic acid at the onset of systemic acquired resistance in cucumber. *Science*, 250: 1004-1106.

Rai, V.K., S.S. Sharma and S. Sharma. 1986. Reversal of ABA-induced stomatal closure by phenolic compounds. *J. Exp. Bot.*, 37: 129-134.

Raskin, I. 1992. Role of Salicylic acid in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, 43: 439-463.

Razi, A., K. Matruda and A. Arslan. 1985. Water stress induced changes in concentration of proline and other solutes in growing regions. *J. Exp. Bot.*, 36: 1716-1725.

Sakhabutdinova, A.R., D.R. Fatkhutdinova, M.V. Bezrukova, F.M. Shakirova. 2003. Salicylic acid prevents the damaging action of stress factors on wheat plants. *Bulg. J. Plant Physiol.* (Special Issue), 314-319.

Saqib, M., R.H. Qurehs, J. Akhtar, S. Nawaz and M. Aslam. 1999. Effect of salinity and hypoxia on growth and ionic composition of different genotypes of wheat. *Pak. J. Soil. Sci.*, 17: 1-8.

Singh, S.K. and M. Kaur. 1980. Effect of growth regulators on podding and yield of mungbean (*Vigna radiata* L.). *Indian J. Plant Physiol.*, 23: 366-370.

Steel, R.G.D., J.H. Torrie and D. Dickey. 1997. *Principles and Procedures of Statistics*. McGraw Hill Book Co. Inc., New York.

Vernooij, B., L. Friedrich, A. Morse, R. Rest, R. Kolditz-Jawahar, E. Ward, S. Uknef, H. Kessman and J. Ryals. 1994. Salicylic acid is not the translocated signal responsible for inducing systemic acquired resistance but is required for signal transduction. *Plant Cell*, 6: 959-965.

Wang, L.J. and S.H. Li. 2006. Salicylic acid induced heat or cold tolerance in relation to Ca^{2+} homeostasis and antioxidant system in young grape plants. *Plant Sci.*, 170: 685-694.

(Received publication 24 February 2007)