Pak. J. Bot., 59(1): January 2027.

ARTICLE | DOI: http://dx.doi.org/10.30848/PJB2027-1(13)

EXOGENOUS GIBBERELLIC ACID (GA_3) ALLEVIATES LEAD-INDUCED PHYTOTOXICITY BY MODULATING GROWTH AND ANTIOXIDANT DEFENCE IN SPINACH (*SPINACIA OLERACEA* L.)

AYESHA MEHMOOD[1]*, ISHTIAQ AHMAD[2], MARYAM[3], ADEL A. REZK[4]*, HOSSAM S. EL-BELTAGI[4], BADER Y. ALSUBAIE[4], OTHMAN AL-DOSSARY[4], AHMED M. ISMAIL[5], KONSTANTIN KOROLEV[6] AND FARIDA SAFAROVA[7]

[1]Institute of Agro-Industry and Environment, Faculty of Agriculture and Environment, The Islamia University of Bahawalpur, 63100, Pakistan
[2]Department of Horticultural Sciences, The Islamia University of Bahawalpur, Pakistan
[3]Department of Environmental Sciences, The Govt. Sadiq College Women University, Bahawalpur 63100, Pakistan
[4]Agricultural Biotechnology Department, College of Agricultural and Food Science, King Faisal University, Al-Ahsa, 31982, Saudi Arabia
[5]Pests and Plant Diseases Unit, College of Agricultural and Food Sciences, King Faisal University, Al-Ahsa 31982, Saudi Arabia
[6]Department of Botany, Biotechnology and Landscape Architecture, School of Natural Sciences, Tyumen State University, Tyumen, 625003, Russia
[7]Department of Biology, Nakhchivan State University, Nakhchivan, Azerbaijan

*Corresponding author's email: ayeshamehmood957@gmail.com; arazk@kfu.edu.sa

Abstract

Lead (Pb) contamination is a serious threat to vegetable production as it adversely affects plant growth, disrupts physiological and oxidative balance. To assess the effects of Pb stress (20 mg L-1) and the mitigating role of gibberellic acid (GA3; 100 mg L-1) on Spinacia oleracea L., a pot experiment was conducted, laid out in a completely randomized design (CRD) with 5 replications. A significant reduction was observed in plant height, root length, fresh weight and dry weight by 34.6%, 20.7%, 25.3%, and 25.2%, respectively under Pb exposure, compared with the control ($p \leq 0.05$). Photosynthetic pigments also reduced with Pb stress, chlorophyll a (0.02 mg g^{-1} FW), chlorophyll b (0.28 mg g^{-1} FW), carotenoids (0.24 mg g^{-1} FW) and relative water content (56.2%), resulting in disruption of photosynthetic activity. Importantly, GA_3 application to plants grown under Pb stress significantly alleviated Pb-induced by restoring growth traits. Application of GA_3 significantly increased chlorophyll content, carotenoids and relative water content. Pb stress triggered antioxidant enzymatic activities, including superoxide dismutase, catalase, peroxidase, and ascorbate peroxidase, reflecting enhanced oxidative stress. GA_3 application under Pb stress further strengthened antioxidant defence, resulting in more efficient reactive oxygen species detoxification.

Key words: Heavy metal toxicity; Lead stress; Plant growth regulation; Leafy vegetable; Oxidative stress

Introduction

Spinach (*Spinacia oleracea* L.) is a popular leafy vegetable and contributes significantly to human nutrition (Iftikhar *et al*., 2023). Vegetables are key for maintaining our health and preventing chronic diseases; they are sources of important vitamins and minerals, as well as many bioactive compounds (Sarma & Bhavya, 2024). Spinach has received special attention as an important dietary constituent with exceptional nutritional composition in a balanced diet (Naseem *et al*., 2023). It contains vitamin A, vitamin C, vitamin K and numerous B vitamins and essential minerals (those include iron, calcium and magnesium). These nutrients are involved in systems including immune response, bone health and antioxidant defence systems. Protection against the oxidation is made by the phytochemicals like phenolics, carotenoids and organic acids (Tufail *et al*., 2025). Spinach helps reduce inflammation and the risk of cardiovascular disease by boosting immune system, moreover, it is also a rich source of dietary fibre and food availability for low-income households (Arooj *et al*., 2025). It is not only beneficial for health but also has economic significance, as it is cultivated extensively across the globe due to its widespread use in salad and cooked recipes (Isaza *et al*., 2025; Roberts & Moreau, 2016).

However, serious problems have been observed in spinach growth due to lead (Pb) toxicity which is a heavy metal (HM) and non-essential for plant growth but causes serious human health issues. Basically, the primary mechanism of lead uptake occurs through the roots from the soil; after entering the plant system, it disturbs various physiological processes (Gupta *et al*., 2024; Maryam *et al*., 2025; Teoh *et al*., 2024). Lead-contaminated soil poses serious risks to food safety and crop yield. According to international standards, permissible limits for lead contamination range from 140 to 400 mg/kg, moreover it depends on local standards for soil physical properties and permissible limits (Ahmad *et al*., 2020; Yi *et al*., 2022). Maximum limit of lead contamination ranges up to 0.3 mg/kg fresh weight that is acceptable for leafy vegetables including spinach, as set by Codex Alimentarius Commission at its 2024 session (Anon., 2024). Generally, lead, a toxic heavy metal, causes severe yield losses through

Received: 06-04-2026 | Revised: 26-07-2026 | Accepted: 02-08-2026 | Online: 25-08-2026

affecting photosynthetic activity and exerting oxidative stress (Ashar *et al*., 2024; Rahmdel *et al*., 2018). Studies have demonstrated that Pb contamination may have detrimental effects on plant morphology and chlorophyll contents, as controlled by oxidative damage reactions, such as increased production of lipid peroxides and reactive oxygen species (ROS) (Emanuil *et al*., 2020; Manzoor *et al*., 2025; Tanhan *et al*., 2025).

Heavy metals disrupt essential physiological processes, which decreases the plant germination and growth (Muhammad *et al*., 2024). Heavy metals in soil (such as lead) are transported quickly through the various plant tissues of spinach, where the maximum concentration has been observed in the roots in most instances due to their uptake tendency (Wang *et al*., 2025). Physiological stress induced by Pb exposure is lethal for spinach production, which signifies the need to develop strategies that could improve its tolerance against Pb toxicity (Bibi *et al*., 2024). Plants absorb Pb mainly through the roots from contaminated soils or water, which leads to the disturbance of ionic homeostasis and nutrient uptake. This blockage results in decreased chlorophyll content, stunted growth and production of lipid peroxidation, damaging the cell membranes and integrity of cells. Especially, Pb mimics essential nutrients and disrupts functions that depend on metal ions homeostasis in the plant body (Gupta *et al*., 2024). Pb imposes oxidative stress through the generation of ROS which causes cell disruption in plants in a chain reaction leading to lipid degradation and protein denaturation that finally ending with necrosis in plant tissues (Ilyas *et al*., 2024).

Several studies have explained how gibberellic acid (GA_3, CAS no. 77-06-5) manage antioxidant enzyme activity and increase oxidative damage resistance caused by lead ions. Physiologically, GA_3 enhances chlorophyll formation, which aids in photosynthesis, as well as controls numerous stress response genes (Vasyuk, 2025). Beyond reducing oxidative stress, GA_3 is helpful to improve the crop quality and yield, it is also considered as a promising tool in precision agriculture on sustainable basis. Mitigation of heavy metal toxicity can be addressed through judicious use of GA_3, through improving nutrient uptake on sustainable basis and enhancing tolerance the heavy metal stress (Ahmad *et al*., 2025).

Gibberellins (GAs) are a family of growth-promoting plant hormones, but the key biochemical and molecular signalling regulatory mechanisms controlling plant growth, developed from changes in hormone metabolism, remain largely unknown (Castro-Camba *et al*., 2022). It is believed that these hormones suppress the harmful effects of ROS formation resulting from exposure to heavy metals (Bücker-Neto *et al*., 2017). This is attributed to the activation of the synthesis of phytohormones and secondary metabolites by GAs, which stimulates stress tolerance pathways, including detoxification pathways for heavy metals chelation (Ghani *et al*., 2025). The significant improvement in chlorophyll content combined with increased biomass accumulation demonstrates the efficiency of GA_3 treatment to increase spinach productivity even in contaminated natural soil (Shahzad *et al*., 2024).

Studies have also demonstrated that GAs are involved in cellular mechanisms that alleviate the harmful impacts of HMs through the induction of adaptive growth strategies (Badar *et al*., 2022). However, the use of GAs has been found to sustain antioxidant enzyme activity, this antioxidant defence pathway is crucial in upholding plant structural integrity when confronted with Pb and is activated by gibberellin signalling to encourage the expression of genes (Li *et al*., 2026). Also, during the vegetative stage, GA_3 signalling has been linked to stress mitigation strategies, which also involve other phytohormones like abscisic acid. These elevated concentrations of abscisic acid (ABA), as customary, are a physiological response to abiotic cues that are capable of amplifying the positive role of GAs by inducing the expression of GA_3 biosynthesis genes or potentially in a synergistic fashion through the activation of growth-associated pathways (Mo *et al*., 2024; Hasddin et al., 2025).

The hypothesis was that GA_3 application could alleviate lead toxicity in spinach by enhancing the plant stress tolerance mechanism. Thus, the present study evaluated the effectiveness of GA_3 in enhancing growth and biomass accumulation of spinach under Pb stress.

Materials and Methods

A pot experiment was executed on *S. oleracea* L. to investigate its response to Pb stress and the role of GA_3 in growth regulation. Twenty plastic pots (59 cm in diameter, 38 cm in height) were filled with 10 kg of loamy soil. The soil samples were also analysed, and the results are shown in Table 1.

Table 1. Physico-chemical properties of soil before the experiment.

Soil texture	Silt loam
Saturation (%)	27
pH	8.0
Electrical conductivity ($mScm^{-1}$)	1.79
Organic matter (%)	0.37
P (mg/ kg)	3.47
K (mg/ kg)	114
Na (mg/ kg)	188
Cl (mg/ kg)	137

Experimental layout: A completely randomized design (CRD) was conducted with one GA_3 concentration (100 ppm) and one Pb concentration (20 ppm) via soil drenching , with deionized water spray as control, with 5 replications. Every single pot had 1 plant. Pb $(NO_3)_2$ was used as the Pb source, and GA_3 was dissolved in water.

Preparation of Pb solution: Pb levels were prepared in deionized water by dissolving the required amounts of Pb. 20 mg of Pb as nitrate salt (32 mg) was diluted to 1000 mL of distilled water for the preparation of a 20 ppm Pb^{2+} solution.

Preparation of GA_3: GA_3 levels were prepared in deionized water by dissolving the required amount of GA_3. 100 mg of GA_3 powder in 1000 mL of distilled water was used for the preparation of a 100-ppm solution.

Treatment application: After 15 days of germination, Pb stress was applied to the soil by adding a freshly prepared aqueous solution of lead salt at the required concentration. Thirty days after germination, the GA_3 treatment was performed using a solution of 100 ppm (100 mg GA_3 dissolved in 1000 mL deionized water as described above). After 60 days, the spinach leaves were collected, and morphological, physiological, and biochemical parameters were measured.

Morphological parameters: Morphological attributes were assessed by harvesting plant samples 40 days after sowing , followed by gentle rinsing with deionized water for removal of surface particulate matter. After harvesting plant samples were brought to the laboratory and roots and shoots were separated for further analysis. Digital balance (Ohaus PR64/E) was used to measure the fresh weight of plant samples. Finally, to remove dust particles distilled water was used for rinsing of plant samples and oven-dried at 70°C for three days (Haider *et al*., 2022).

Plant height (cm): The plant height was measured using measuring scale, plants were selected randomly from each treatment (n= 5 pots per treatment) and measurements were taken from the base of the plant at soil level up to tallest leaf of the plant.

Root length (cm): Using a scale, the length of the roots of randomly chosen plants was measured from the tip of the longest root to the point where the root and shoot join.

Fresh weight (g): Fresh weight was determined by carefully uprooting the plant, followed by washing to remove surface soil, blotting excess moisture with tissue paper, and weighing the plant samples using a digital electronic balance.

Dry weight (g): Dry weight of harvested plant sample was done by placing the plant samples in paper bags followed by placing in oven while maintaining the temperature at 70°C for 3 days till constant weight was attained and then weighed using a digital balance in g.

Physiological Parameters

Chlorophyll determination: Chlorophyll contents were determined by harvesting of fresh plant leaves (0.5 g) randomly from each treatment followed by washing with distilled water to remove dust particles from the sample. After washing the known weight of plant sample was ground by adding 80% acetone with a pestle and mortar. The ground material was centrifuged at 10000 × g for 10 minutes followed by separating the supernatant for further analysis. The extract was subjected to spectrophotometer absorbance at 480, 645, and 663 nm (Lichtenthaler and Wellburn, 1983). The following formulae were used (OD = optical density, subscript = wavelength).

$$\textbf{Chlorophyll (a)} = [(12.7 \times O.D_{663}) - (2.69 \times OD_{645})] \times V/1000 \times W \quad (1)$$

$$\textbf{Chlorophyll (b)} = [(22.9 \times OD_{645}) - (4.68 \times OD_{663})] \times V/1000 \times W \quad (2)$$

$$\textbf{Total chlorophyll} = [(20.2 \times OD_{645} + (8.02 \times OD_{663})] \times V/1000 \times W \quad (3)$$

$$\textbf{Carotenoids (Car)} = [OD_{480} + (0.114 \times OD_{663})] - (0.638 \times OD_{645}) \quad (4)$$

Where:

- OD663, OD645 = Optical density at 663 nm and 645 nm respectively
- 12.7 and 2.69 = coefficients of specific absorption
- V = Volume of extract (mL)
- W = Weight of fresh leaf sample (g)
- 1000 = converts µg to mg if needed, depending on units used.

Relative water content: Fully matured leaves from each treatment were gently separated from the lamina base followed by sealing in plastic labelled bags for analysis. After harvesting leaves were soaked in deionized water for 16 to 18 hours to obtain turgid weight (TW). Measurement of dry weight (DW) was done after 3 days oven dry at 70°C. RWC was calculated using the mathematical formula created by (Barrs and Weatherley, 1962).

Biochemical parameters: A homogenized mixture of 1 g plant sample was made by grinding in 5 mL of chilled extraction buffer (K_2HPO_4, pH 7.0) composed of 1 mM EDTA and 1% PVP (polyvinylpyrrolidone), followed by centrifugation at 15,000 × g for 15 minutes maintaining the temperature at 4°C. The supernatant was separated into layers for further biochemical analysis (You et al., 2019). Superoxide dismutase (SOD) activity was assessed as described by Bradford (1976). Catalase (CAT) activity was evaluated by monitoring the degradation of H_2O_2 at 240 nm, using an extinction coefficient (ε) of 0.0394 mM^{-1} cm^{-1}. Peroxidase (POD) activity was assessed through guaiacol oxidation at 470 nm, and ascorbate peroxidase (APX) activity was measured by the decline in ascorbate absorbance at 290 nm (ε = 2.8 mM^{-1} cm^{-1}).

Statistical analysis

Data regarding various morphological and biochemical attributes was recorded and arranged for further statistical analysis. For the assessment of significance among the treatments analyses were performed in Statistix 8.1. Treatment effects were evaluated using one-way analysis of variance (ANOVA), followed by the Least Significant Difference (LSD) test for mean comparison at the 5% significance level ($p \leq 0.05$).

Results

Effects of Pb stress and GA_3 on morphological traits of *S. oleracea*: Morphological performance of *S. oleracea* was severely affected due to Pb exposure as presented in Table 2, however application of GA_3 significantly improved growth ($p<0.05$). It was noticed that under control conditions, plants gained a height of 15.6 ± 0.92 cm, root length of 8.10 ± 0.94 cm, fresh weight of 40.14 ± 6.64 g and dry weight of 7.22 ± 0.25 g. Exogenous application of GA_3 significantly improved all growth traits, as maximum plant height was attained (21.0 ± 1.0 cm), longest roots (10.0 ± 0.15 cm) and the greatest biomass accumulation (50.20 ± 2.55 g fresh weight and 9.10 ± 0.16 g dry weight) as shown in Table 2. Compared with the control, GA_3 increased plant height by approximately 34.6%, root length by 23.5%, fresh weight by 25.1%, and dry weight by 26.0%.

In contrast, Pb stress severely reduced growth. Plant height declined to 10.2 ± 1.46 cm, root length to 6.42 ± 0.44 cm, fresh weight to 30.00 ± 1.52 g, and dry weight to 5.40 ± 0.33 g, all significantly lower than the control ($p<0.05$). In plants that did not receive GA_3, Pb exposure caused reductions of 34.6% in plant height, 20.7% in root length, 25.3% in fresh weight and 25.2% in dry weight, confirming the strong inhibitory effect of Pb on vegetative growth and biomass production.

Surprisingly, it was observed that combined application (Pb + GA_3) significantly mitigated Pb-induced growth suppression. Plants under Pb + GA_3 reached a height of 15.2 ± 1.07 cm and root length of 7.20 ± 0.28 cm, which were statistically comparable to the control for plant height (same letter "b") but remained lower than GA_3 alone. Similarly, fresh weight (39.76 ± 2.93 g) and dry weight (7.85 ± 0.32 g) were significantly higher than Pb treatment alone and approached control values. Compared with Pb-stressed plants, GA_3 supplementation under Pb improved plant height by 49.0%, root length by 12.1%, fresh weight by 32.5%, and dry weight by 45.4%. Conclusively, Pb stress significantly suppressed morphological traits in *S. oleracea*. However, application of GA_3 not only stimulated growth but also substantially alleviated Pb-induced growth inhibition.

Effect of Pb stress and GA_3 on photosynthetic pigments (mg g^{-1} FW) and relative water content (%) of *S. oleracea*: Lead (Pb) stress markedly impaired the physiological performance of *Spinacia oleracea* (Table 3). Plants exposed to Pb showed a significant reduction in chlorophyll a (0.28 mg g^{-1} FW), chlorophyll b (0.02 mg g^{-1} FW), and carotenoids (0.24 mg g^{-1} FW) compared with the control. Relative water content (RWC) also declined sharply to 36.20%, indicating severe disturbance of plant water status under Pb toxicity.

Application of GA_3 significantly improved the accumulation of photosynthetic pigments and water relations, with maximum values recorded for chlorophyll a (0.43 mg g^{-1} FW), chlorophyll b (0.09 mg g^{-1} FW), carotenoids (0.52 mg g^{-1} FW), and RWC (77.40%). Surprisingly, GA_3 application on plants receiving Pb stress restored these traits.

Effects of Pb stress and GA_3 on enzymatic activity of *S. oleracea*: A significant activation of the antioxidant defense system of *Spinacia oleracea* was observed due to Lead (Pb) stress (Fig. 1a–d). Superoxide dismutase (SOD) activity was increased from78 U mg^{-1} protein in control plants to 86 U mg^{-1} protein under Pb stress, indicating enhanced production of superoxide radicals. Exogenous application of GA_3 under Pb stress further increased SOD activity to about 90 U mg^{-1} protein, suggesting improved dismutation of superoxide ions into hydrogen peroxide.

Peroxidase (POD) activity also showed a significant increase, rising from 82 U mg^{-1} protein in control plants to 96 U mg^{-1} protein under Pb stress. The highest POD activity (100 U mg^{-1} protein min^{-1}) was recorded in Pb + GA_3–treated plants, reflecting enhanced hydrogen peroxide detoxification.

Catalase (CAT) activity (mmol H_2O_2 mg^{-1} protein) was increased markedly under Pb stress, reaching 11 mmol mg^{-1} protein compared with 6 mmol mg^{-1} protein in the control. GA_3 application further elevated CAT activity to 14 U mg^{-1} protein under Pb stress, indicating a stronger capacity for hydrogen peroxide scavenging.

Ascorbate peroxidase (APX) activity increased from 40 U mg^{-1} protein in control plants to 46 U mg^{-1} protein under Pb stress and reached a maximum value of 55 U mg^{-1} protein in Pb + GA_3–treated plants. This enhancement reflects a more efficient ascorbate–glutathione cycle under GA_3 supplementation.

Table 2. Effects of Pb stress and GA_3 on morphological parameters of *S. oleracea*.

Treatment	Plant height (cm)	Root length (cm)	Plant fresh weight (g)	Plant dry weight (g)
Control	15.6 ± 0.92b	8.10 ± 0.94ab	40.14 ± 6.64 b	7.22 ± 0.25ab
GA_3	21.0 ± 1.0a	10.0 ± 0.15a	50.20 ± 2.55a	9.10 ± 0.16a
Pb	10.2 ± 1.46c	6.42 ± 0.44b	30.00 ± 1.52c	5.40 ± 0.33b
Pb + GA_3	15.2 ± 1.07b	7.20 ± 0.28b	39.76 ± 2.93b	7.85 ± 0.32 b

Mean values sharing different letters are statistically significant at $p<0.05$ probability

Table 3. Effects of Pb stress and GA_3 on physiological parameters of *S. oleracea*.

Treatment	Chlorophyll a (mg g^{-1} FW)	Chlorophyll b (mg g^{-1} FW)	Carotenoids (mg g^{-1} FW)	Relative water content (%)
Control	0.36 ± 0.02b	0.04 ± 0.01b	0.41± 0.01a	66.00 ± 2.77 c
GA_3	0.43± 0.06a	0.09 ± 0.03a	0.52 ± 0.06a	77.40 ± 1.69a
Pb	0.28 ± 0.03c	0.02 ± 0.06c	0.24 ± 0.03a	36.20 ± 2.83d
Pb + GA_3	0.38 ± 0.02b	0.04 ± 0.12b	0.38 ± 0.02a	61.80 ± 2.97b

Mean values sharing different letters are statistically significant at $p<0.05$ probability



Fig. 1. **(A)** Impact of gibberellic acid (GA_3) on SOD (U mg^{-1} Protein) of *S. oleracea* under lead stress **(B)** Impact of gibberellic acid (GA_3) on POD (U mg^{-1} Protein) *S. oleracea* under lead stress **(C)** Impact of gibberellic acid (GA_3) on CAT (U mg^{-1} Protein) *S. oleracea* under lead stress **(D)**. Impact of gibberellic acid (GA_3) on APX (U mg^{-1} Protein) *S. oleracea* under lead stress

Discussion

Among the heavy metals lead (Pb) is recognized as a vital abiotic stressor that interferes with plant growth and development by disrupting nutrient uptake, damaging cellular structures and inducing oxidative stress (Gupta *et al*., 2024; Qiu *et al*., 2024), leading to limited vegetative growth, reduced biomass and impaired physiological processes (Meena *et al*., 2019). Results of current study revealed that Pb stress significantly reduced morphological traits including plant height, root length, fresh weight and dry weight of *Spinacia oleracea*, authenticating that Pb hampers cell division which indirectly restricts growth and development. These results are consistent with recent findings in other plant systems (Alia et al., 2015; Wang *et al*., 2023; Zhao *et al*., 2026), where Pb exposure led to deterioration in morphological and biomass traits due to inhibition of photosynthetic and metabolic activities (Haider *et al*., 2022) as Pb exposure interrupts the membrane integrity at cellular level indirectly disrupts the nutrient uptake results compromised chloroplast development.

Gibberellic acid (GA_3) has been extensively documented as a key phytohormone that promotes cell elongation, division, and overall plant growth (Camara *et al*., 2018). In the current work, GA_3 application significantly enhanced all morphological traits under both control and Pb stress conditions (Asif *et al*., 2025; Feijoo Leon et al., 2025). GA_3–treated plants attained greater heights, longer roots, and higher fresh and dry weights compared with controls, reflecting stimulation of growth processes. Similar promotive effects of GA_3 on plant growth under heavy metal stress have been reported in other studies; for example, combinations of GA_3 with rhizobacterial inoculation improved growth and chlorophyll content in Pb-stressed *Solanum melongena* (Ogugua *et al*., 2022), and GA_3 alleviated Cu stress in spinach by enhancing antioxidant responses and biomass. The growth promoting ability of GA_3 is biologically associated with cell division and elongation, in addition to this GA_3 activates relevant genes responsible for loosening and expansion of the cell wall.

Importantly, GA_3 significantly ameliorated Pb-induced growth suppression in *S. oleracea*. Morphological performance in Pb + GA_3–treated plants was significantly higher compared to plants with Pb only and, for plant height, statistically comparable to control levels (Srivastava & Srivastava, 2023). This indicates that GA_3 not only stimulates growth but also confers a protective effect under Pb toxicity, likely through enhanced hormonal balance, improved nutrient

acquisition, and modulation of stress responses (Shahzad *et al*., 2024). Enhanced antioxidant enzyme activities and improved physiological parameters observed in this study further support the notion that GA_3 aids in mitigating Pb stress by reducing oxidative damage and preserving cellular integrity, thereby maintaining growth dynamics even under adverse conditions.

The decline in chlorophyll content under Pb exposure suggests that Pb toxicity interfered with the photosynthetic apparatus, while the reduction in carotenoids indicates a possible disturbance in the plant's photoprotective capacity. Similarly, the lower RWC under Pb stress reflects impaired plant water relations. These observations are consistent with Pandey et al. (2009), who reported that Pb and Cd stress adversely affected photosynthetic pigments and physiological performance in spinach. GA_3 application improved these physiological responses, and its combined application with Pb partially alleviated the adverse effects of Pb toxicity. In addition, Pb exposure increased the activities of antioxidant enzymes, including SOD, POD, CAT and APX, indicating activation of the plant's antioxidant defense system in response to oxidative stress. The highest enzyme activities were observed under the combined Pb + GA_3 treatment, suggesting that GA_3 further stimulated antioxidant defense mechanisms under Pb stress. SOD converts superoxide radicals into hydrogen peroxide, while CAT and POD help eliminate hydrogen peroxide and protect cellular components from oxidative damage. Similar changes in antioxidant responses under heavy-metal stress have been reported in spinach by Emanuil et al. (2020). Overall, the improved physiological responses alongside enhanced antioxidant enzyme activities suggest that GA_3 may contribute to maintaining photosynthetic function, improving plant water status, and strengthening antioxidant defenses in spinach under Pb-induced stress (Wang et al., 2025).

Conclusions

This study shows that lead (Pb) stress markedly restricts growth and disrupts physiological and biochemical processes in *Spinacia oleracea*, as evidenced by reduced biomass, lower photosynthetic pigment levels, poor water status, and enhanced oxidative stress. Exogenous application of gibberellic acid (GA_3) not only promoted plant growth under normal conditions but also effectively alleviated Pb-induced damage by improving morphological traits, maintaining chlorophyll content and relative water content, and strengthening the antioxidant defense system. The coordinated enhancement of growth and redox balance under GA_3 treatment suggests that this phytohormone plays an important protective role against Pb phytotoxicity. Overall, GA_3 emerges as a promising and practical approach to improve spinach tolerance to Pb stress, although further studies under field conditions and assessments of Pb accumulation in edible tissues are needed before broader agricultural application.

Author's Contributions: conceptualization: Ayesha Mehmood, Ishtiaq Ahmad and Maryam; Formal Analysis: Maryam, validation: Adel A. Rezk, Hossam S. El-Beltagi and Bader Y. Alsubaie; Acquisition of funds: Othman Al-Dossary and Ahmed M. Ismail; writing draft: Konstantin Korolev; Reviewing: Farida Safarova

Funding Statement: The authors extend their appreciation to the Researchers Supporting Project number (KFU262628), King Faisal University, Al-Ahsa, 31982, Saudi Arabia, for the financial support that enabled the successful execution of this research.

Acknowledgements

Authors extend their appreciation to the Deanship of Scientific Research, Vice Presidency for Graduate Studies and Scientific Research, King Faisal University, Saudi Arabia, for funding this research work (KFU262628).

Conflict of Interest: There is no conflict of interest related to current study.

References

Abubakar, A.S., Y. Wu, F. Chen, A. Zhu, P. Chen, K. Chen, X. Qiu, X. Huang, H. Zhao, J. Chen and G. Gao. 2023. Comprehensive analysis of WUSCHEL-related homeobox gene family in *Boehmeria nivea* indicates its potential role in adventitious root development. *Biology*, 12(12): 1475.

Ahmad, A., S. Hussain, S.M. Rao, A. Asghar, M. Irshad, M. Aslam, S.S. Shafqat, M. Javed, M.S. Bashir and H.S. Kausar. 2020. Comparative studies of lead and heavy metals concentrations in Pakistan soil and its toxic effects. *Pol. J. Environ. Stud*., 29: 2533.

Ahmad, I., M.R. Bozhuyuk, A. Maryam, R. Ahmad and M.A. Anjum. 2025. Applications of plant growth regulators in vegetables. In: *Sustainable and Innovative Vegetable Production in Times of Climate Change*, p. 311.

Hasddin, Ulyasniati, Jamal Mukaddas, Marjani, Mirad, Mustafa Kan, Abdul Aman Ega and Musadia Afa. 2025. Sustainable agriculture: The role of biostimulants in enhancing crop growth and resilience. *J. Glob. Innov. Agric. Sci.* 13(2): 1688. https://doi.org/10.22194/JGIAS/25.1688

Anonymous. 2024. FAO/WHO. *Codex Alimentarius: General Standard for Contaminants and Toxins in Food and Feed*. 17th Session of the Codex Committee on Contaminants in Foods (CCCF17).

Feijoo Leon, A. D., Guerrero Salinas, K. F., Conrrado, C., Cajamarca Carrazco, D., Huebla Concha, V. H. and Brito, M. 2025. The role of acetylsalicylic acid in forage production: Scientific evidence and human dimensions of agricultural innovation. *J. Glob. Innov. Agric. Sci.* 13(4): 1257-1265. https://doi.org/10.22194/JGIAS/25.1835

Barrs, H.D. and Weatherley, P.E. 1962. A re-examination of the relative turgidity technique for estimating water deficits in leaves. *Aust. J. Biol. Sci.,* 15: 413-428. DOI: 10.1071/BI9620413.

Iacobellis, I., Lisi, A., Vacca, M., Apa, C.A., Celano, G., Mancini, L., Minervini, F., Calasso, M. and De Angelis, M. (2024). Nutritional, biochemical, and functional properties of spinach leaf-enriched dough: A healthier alternative to conventional pasta. *Foods* 13(22): 3608.

Bücker-Neto, L., Paiva, A.L.S., Machado, R.D., Arenhart, R.A. and Margis-Pinheiro, M. (2017). Interactions between plant hormones and heavy metals responses. *Genetics and*

Molecular Biology 40(1): 373-386. Ashar, Z., E. Syam'un, F. Ulfa and I.R. Saleh. 2024. Effect of *Tithonia diversifolia* compost and *Trichoderma* sp. coating on potato (*Solanum tuberosum* L.) growth and yield. *J. Glob. Innov. Agric. Sci*., 12: 1093.

Badar, Z., A. Shanableh, A. El-Keblawy, K. Mosa, L. Semerjian, A. Mutery and M. Semreen. 2022. Assessment of uptake, accumulation and degradation of paracetamol in spinach (*Spinacia oleracea* L.) under controlled laboratory conditions. *Plants*, 11: 1626.

Bibi, A., F. Rasul, S. Shahzad, R. Sakrabani, W.U.D. Din, P. McKenna and M. Sajid. 2024. Toxicity, bioaccumulation and mitigating strategies of heavy metals stress on morpho-physiology of spinach. *Discover Plants*, 1: 1.

Emanuil, N., M. Akram, S. Ali, M. El-Esawi, M. Iqbal and M. Alyemeni. 2020. Peptone-induced physio-biochemical modulations reduce cadmium toxicity and accumulation in spinach (*Spinacia oleracea* L.). *Plants*, 9: 1806.

Camara, M.C., Vandenberghe, L.P.S., Rodrigues, C., de Oliveira, J., Faulds, C., Bertrand, E. and Soccol, C.R. (2018). Current advances in gibberellic acid (GA_3) production, patented technologies and potential applications. *Planta,* 248(5): 1049-1062.

Ghani, M., M. Abbas, B. Ali, R. Aziz, R. Qadri, A. Noor and S. Ali. 2021. Alleviating role of gibberellic acid in enhancing plant growth and stimulating phenolic compounds in carrot (*Daucus carota* L.) under lead stress. *Sustainability*, 13: 12329.

Ghani, M.A., M. Azam, B. Ali, M. Azam, W. Qadri, K. Zia, M.M. Jahangir, F. Shireen and M.M. Abbas. 2025. Salicylic acid-induced thermotolerance and antioxidant enzyme modulation in sponge gourd (*Luffa cylindrica*) seedlings. *J. Hortic. Sci. Technol*., 8: 125.

Lichtenthaler, H.K. and Wellburn, A.R. 1983. Determinations of total carotenoids and chlorophylls a and b of leaf extracts in different solvents. *Biochem. Soc. Trans.,* 11: 591-592. Gupta, M., V. Dwivedi, S. Kumar, A. Patel, P. Niazi and V. Yadav. 2024. Lead toxicity in plants: Mechanistic insights into toxicity, physiological responses of plants and mitigation strategies. *Plant Signal. Behav*., 19: 1.

Haider, F.U., X. Wang, U. Zulfiqar, M. Farooq, S. Hussain, T. Mehmood, M. Naveed, Y. Li, C. Liqun, Q. Saeed, I. Ahmad and A. Mustafa. 2022. Biochar application for remediation of organic toxic pollutants in contaminated soils: An update. *Ecotoxicol. Environ. Saf*., 248: 114322.

Iftikhar, M., M. Rashid, Z. Yousaf, A. Tufail, N. Zahra, H. Ahmad and A. Hannan. 2023. Taxonomic evaluation of *Spinacia oleracea* L. accessions by morphological and anatomical markers. *J. Plant Environ*., 5: 135.

Ilyas, M.Z., K.J. Sa, M.W. Ali and J.K. Lee. 2024. Toxic effects of lead on plants: Integrating multi-omics with bioinformatics to develop Pb-tolerant crops. *Planta*, 259: 18.

Isaza, C., A.M. Aleman-Trejo, C.F. Ramirez-Gutierrez, J.P. Zavala de Paz, J.A. Rizzo-Sierra and K. Anaya. 2025. Spinach (*Spinacia oleracea* L.) growth model in indoor controlled environment using Agriculture 4.0. *Sensors*, 25: 1684.

Pandey, N., Pathak, G.C., Pandey, D.K. and Pandey, R., 2009. Heavy metals, Co, Ni, Cu, Zn and Cd, produce oxidative damage and evoke differential antioxidant responses in spinach. *Brazil. J. Plant Physiol.*, 21: 103-111.

Shahzad, M.A., Younis, U., Ehsan, A., Alarfaj, A.A., Alharbi, S.A. and Ansari, M.J. (2024). Impact of gibberellic acid (GA_3), quantum dot biochar, and rhizosphere bacteria on fenugreek plant growth and stress responses under lead stress. *Scientific Reports* 14(1): 29612.You, L., Song, Q., Wu, Y., Li, S., Jiang, C., Chang, L., Yang, X. and Zhang, J. 2019. Accumulation of glycine betaine in transplastomic potato plants expressing choline oxidase confers improved drought tolerance. *Planta,* 249: 1963-1975.

Manzoor, M., K. Korolev and R. Ahmad. 2025. Zinc oxide nanoparticles: Abiotic stress tolerance in fruit crops focusing on sustainable production. *Phyton (Int. J. Exp. Bot.)*, 94: 1401.

Maryam, M., M.F. El-Mogy, I. Jan, I. Naz, I. Ahmad, R. Ahmad and M.T. Altaf. 2025. Nanoparticle innovations for mitigating metal toxicity in plants. *Phyton (Int. J. Exp. Bot.)*, 94: 623.

Meena, V., M.L. Dotaniya, J.K. Saha, H. Das and A.K. Patra. 2019. Impact of lead contamination on agroecosystem and human health. In: *Lead in Plants and the Environment*, pp. 67-82.

Muhammad, H.M.D., S. Naz, R. Ahmad, E. Ali, M. Anwar, M.A. Altaf, S. Alansi, A.A. Alsahli and S. Abou Fayssal. 2024. Melatonin triggers salinity tolerance in pansy (*Viola tricolor*) by regulation of defense system. *J. King Saud Univ. Sci*., 36: 103286.

Naseem, A., S. Akhtar, T. Ismail, M. Qamar, Dur-e-Shahwar, W. Sattar, T. Saeed, T. Esatbeyoglu, E. Bartkiene and J.M. Rocha. 2023. Effect of growth stages and lactic acid fermentation on anti-nutrients and nutritional attributes of spinach (*Spinacia oleracea*). *Microorganisms*, 11: 2343.

Tufail, T., Fatima, S., Bader Ul Ain, H., Ikram, A., Noreen, S., Rebezov, M., Al-Farga, A., Saleh, R. and Shariati, M.A. (2025). Role of phytonutrients in the prevention and treatment of chronic diseases: A concrete review. *ACS Omega* 10(13): 12724-12755.Ogugua, U.V., Kanu, S.A. and Ntushelo, K. (2022). Gibberellic acid improves growth and reduces heavy metal accumulation: A case study in tomato (*Solanum lycopersicum* L.) seedlings exposed to acid mine water. *Heliyon* 8(12): e12399.

Qiu, L., C. Yan, T. Munir, Y. Wang, E. Wang, R. Li, X. Wu, Y. Huang and B. Li. 2024. Comparing struvite, K-struvite and hydroxyapatite for the remediation of lead and cadmium contaminated soil. *Sustain. Horiz*., 10: 100084.

Rahmdel, S., M. Rezaei, J. Ekhlasi, S.H. Zarei, M. Akhlaghi, S.M. Abdollahzadeh, R. Sefidkar and S.M. Mazloomi. 2018. Heavy metals (Pb, Cd, Cu, Zn, Ni, Co) in leafy vegetables collected from production sites: Their potential health risk to the general population in Shiraz, Iran. *Environ. Monit. Assess*., 190: 650.

Roberts, J.L. and R. Moreau. 2016. Functional properties of spinach (*Spinacia oleracea* L.) phytochemicals and bioactives. *Food Funct*., 7: 3337.

Mo, W., Zheng, X., Shi, Q., Zhao, X., Chen, X., Yang, Z. and Zuo, Z. 2024. Unveiling the crucial roles of abscisic acid in plant physiology: Implications for enhancing stress tolerance and productivity. *Front. Plant Sci.,* 15: 1437184.

Castro-Camba, R., Sánchez, C., Vidal, N. and Vielba, J.M. (2022). Plant development and crop yield: The role of gibberellins. *Plants* 11(19): 2650.Sarma, U. and T.R. Bhavya. 2024. Dietary phytonutrients in common green leafy vegetables and the significant role of processing techniques on spinach: A review. Food Prod. Process. *Nutr*., 6: 10.

Shahzad, M.A., U. Younis, A. Ehsan, A.A. Alarfaj, S.A. Alharbi and M.J. Ansari. 2024. Impact of gibberellic acid (GA_3), quantum dot biochar and rhizosphere bacteria on fenugreek plant growth and stress responses under lead stress. *Sci. Rep*., 14: 29612.

Li, Z., Liu, X., Meng, W., Shangguan, J., Zhang, J., Ali, I., Yao, N., Zhang, M., Ahmad, N. and Liu, X. (2026). Gibberellic

acid-induced regulation of antioxidant–flavonoid channels provides protection against oxidative damage in safflower under salinity stress. *Plants* 15: 267.

Asif, M., Md., Imran, S., Paul, N.C., Rabbi, R.H.Md., Jahan, N., Sarker, P., Najmol Hoque, Md., Jahan Sumi, M., Asaduzzaman, Md., Rehman, S.U., Brestic, M., Bárek, V., Skalicky, M., Hossain, A. and Rhaman, M.S. (2025). An overview and current progress of gibberellic acid-mediated abiotic stress alleviation in plants. *Plant, Soil Environ.,* 71(7): 453-479.

Srivastava, D. and N. Srivastava. 2023. Molecular mechanism of lead toxicity and tolerance in plants. In: *Lead Toxicity: Challenges and Solution*, pp. 247-286.

Tanhan, P., K. Imsilp, N. Lansubsakul and W. Thong-asa. 2025. Evaluation of lead and cadmium bioaccumulations related to oxidative responses in the tissues of two wild mud crabs, *Scylla olivacea* and *Scylla paramamosain*, from Pattani Bay, Thailand. *Asian J. Agric. Biol*., e2025054.

Teoh, Y.J., W.H. Cheng, W. Wider, K. Krishnan, C.K. Yap, Q. Chen, L. Jiang, T.M.T. Lei, T. Tanee, R. Sudmoon and S.Y. Lee. 2024. Heavy metal accumulation in paddy plants and health risks: Insights from southern peninsular Malaysia and global research trends. *Asian J. Agric. Biol*., 2024135.

Vasyuk, V.A. 2025. The effect of gibberellic acid (GA_3) on growth, photosynthetic pigments and metal biosorption in the water fern *Salvinia natans* (L.) All. under zinc stress. *Biotechnol. Acta*, 18: 23.

Wang, J., H. Sun, H. Mou and S. Yang. 2024. Protein and lysine improvement harnessed by a signal chain of red light-emitting diode light in *Chlorella pyrenoidosa*. *Bioresour. Technol*., 414: 131620.

Wang, N., Z. Liu, L. Nan and Y. Luo. 2025. Effects of different concentrations of lead and cadmium stress on spinach growth and physiology. *Bangladesh J. Bot*., 54: 855-864.

Wang, Y.X., T.F. Yu, C.X. Wang, J.T. Wei, S.X. Zhang, Y.W. Liu, J. Chen, Y.B. Zhou, M. Chen, Y.Z. Ma, J.H. Lan, J.C. Zheng, F. Li and Z.S. Xu. 2023. Heat shock protein TaHSP17.4, a TaHOP interactor in wheat, improves plant stress tolerance. *Int. J. Biol. Macromol*., 246: 125694.

Yi, J., H. Li, Y. Zhao, M. Shao, H. Zhang and M. Liu. 2022. Assessing soil water balance to optimize irrigation schedules of flood-irrigated maize fields with different cultivation histories in the arid region. Agric. *Water Manag*., 265: 107543Bradford, M.M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.,* 72: 248-254.

Gupta, M., V. Dwivedi, S. Kumar, A. Patel, P. Niazi and V.K. Yadav. 2024. Lead toxicity in plants: Mechanistic insights into toxicity, physiological responses of plants and mitigation strategies. *Plant Signal. Behav.*, 19(1): 2365576. https://doi.org/10.1080/15592324.2024.2365576

Zhao, M., Liu, W., Feng, W., Zhang, C., Zhang, J., Xu, L., Ma, X., Zou, G. and Chen, Y. 2026. Shape-dependent polyurethane microplastics reprogram root architecture and detoxification modules to suppress cadmium uptake in a hyperaccumulator. *J. Environ. Chem. Eng.* 14(4): 123400. https://doi.org/10.1016/j.jece.2026.123400

Alia, N., Sardar, K., Said, M., Salma, K., Sadia, A., Sadaf, S., Toqeer, A. and Miklas, S. (2015). Toxicity and bioaccumulation of heavy metals in spinach (*Spinacia oleracea*) grown in a controlled environment. *Int. J. Environ. Res. Public Health* 12: 7400-7416.